

X. — A NEW MYNES (NYMPHALIDAE) FROM BOEROE.  
BY J. H. JURRIAANSE AND G. VOLBEDA, ROTTERDAM. —  
(WITH PLATE I).

*Mynes talboti* spec. nov.

Male. Forewing upperside: yellowish-white (XVII 25' f), 1) costa and apical part black with bluish tinge. Upper and lower discoidal nervure black till end of cell. Outer margin black, 1 mm. wide. Sub-marginal band from black apex till anal angle greenish black (XLVII 33''''b) about 4 mm. wide, blackened on veins 1, 2 and 3.

Hindwing upperside: whitish blue-green, (XLVIII 37''''b) lightest at basal part and abdominal margin. A black border from costal margin till anal angle, narrowest at vein 3 about 5 mm. wide. A thin white marginal line from vein 3 till about half way abdominal margin.

Forewing underside: white, yellowish at base. A black marginal line from base via apex till anal angle, about 1 mm. wide. A curved dark brown (XLVI 13''''k) band about 5 mm. wide, from cell above the lower discoidal nervure, touching the terminal margin between veins 2, 3 and 4, and bending a little inwards, ending on vein 1. In interspace 6, about 5 mm. from cell-apex an oblong whitish spot followed by a smaller oblong yellow spot. Under those, in interspace 5 a more or less prominent yellowish streak. In interspace 3 and 4 a sub-marginal orange-red (II 11. i.) more or less rectangular spot, divided by vein 4.

One ♂ shows this red spot very pronounced in interspace 3 only, leaving a few red scales in interspace 4.

Apex white and yellow, with black veins. Interspace 8 with a few black scales, interspace 6 about 1 mm. longer than 7 and a little triangular part in interspaces 4 and 5.

Hindwing underside: brownish-black (XLVI 13''''k). A triangular red spot (II 11. i.) from basis above costal-nervure. A post-discal row of more or less prominent black lunules turned inwards from apex till half way abdominal margin, between the veins. An olive yellow (XXX 23') sub-terminal band from apex till anal angle, bordered by a line of four black spots till vein 4. In interspace 1, 2 and 3 terminal black, more or less lunule-shaped spots.

---

1) The figures given behind the names of colors refer to Ridgway, Color standards and color nomenclature, Washington, 1912.

In interspace *1a* and *1b* an abdominal yellow (XVI 21') space with a triangular black marginal spot a little under the end of post-discal row of lunules. A diffused whitish line from apex till vein 4. A black line of the same width and length and a clear white terminal line from apex till half way abdominal margin, bordered with black. Cilia white.

Thorax: upperside slate-blue, underside black.


Abdomen: upperside blackish, underside whitish yellow. Length of forewing 35 mm.

3 ♂♂ — Island Boeroe — Wae Katin. Dec. 1920—Jan. 1922.

In collection Jurriaanse. Type numbers 538—540. One ♂, number 538, offered to the Leyden Museum.

Rotterdam, August 1922.

---



MYNES TALBOTI ♂

ISL. BOEROE.

See page 127.