

Fossils of the type Maastrichtian

J.W.M. Jagt, S.K. Donovan & E.A. Jagt-Yazykova

Jagt, J.W.M., Donovan, S.K. & Jagt-Yazykova, E.A. Fossils of the type Maastrichtian. In: Jagt, J.W.M., Donovan, S.K. & Jagt-Yazykova, E.A. (eds.), *Fossils of the type Maastrichtian (Part 1)*. Scripta Geologica Special Issue, 8: 1-4, Leiden, February 2012.

John W.M. Jagt, Natuurhistorisch Museum Maastricht, de Bosquetplein 6-7, NL-6211 KJ Maastricht, the Netherlands (john.jagt@maastricht.nl); Stephen K. Donovan, NCB Naturalis, Department of Geology, Postbus 9517, NL-2300 RA Leiden, the Netherlands (Steve.Donovan@ncbnaturalis.nl); Elena A. Jagt-Yazykova, Uniwersytet Opolski, Zakład Paleobiologii, Katedra Biosystematyki, ul. Oleska 22, PL-45-052 Opole, Poland (eyazykova@uni.opole.pl).

The present special issue is the first in a series to be devoted to the macrofossil content of the uppermost Cretaceous chalks and biocalcarenites in the type area of the Maastrichtian Stage, the youngest time slice of the Mesozoic Era, dated 70.6 ± 0.6 at the base and 65.5 ± 0.3 Ma at the top (www.stratigraphy.org; see also Husson *et al.*, 2011). The time is right to have such a series, because macrofossils from the area are immensely popular with professional and amateur palaeontologists alike. In fact, new taxa are being discovered almost every month; these either present new records for the area of species known from elsewhere or undescribed forms in need of a name. This may come as some surprise, because several smaller, albeit prolific, quarries in southern Limburg have been closed down in recent years and quarrying activities at the ENCI-HeidelbergCement Group quarry (Maastricht) are to cease on July 1, 2018. In addition, on a daily basis, we receive queries from both collectors and fellow professionals on how to track down literature items on particular fossils from the type Maastrichtian. In general, papers on most biota are seriously outdated, and older works often are hard to come by and suffer from a lack of stratigraphical detail. Other data are widely dispersed in the literature and time-consuming library searches would be needed to get hold of these. Other complicating factors are that not all papers are in English, or appeared in local or regional journals, with a limited distribution.

Although macrofossils of all kinds from the type Maastrichtian are renowned, there has always been a strong bias towards vertebrates (mosasaurs, turtles, sharks, rays and bony fish), echinoids and ammonites. It is not surprising, therefore, to see a plethora of recent papers on these groups (e.g., Herman, 1977; Kennedy, 1987; van der Ham *et al.*, 1987, 2006; Mulder, 2003; Schulp, 2006; Brouwers *et al.*, 2012), while others are more or less ignored. They do deserve better, though.

In an attempt to bring several of the lesser-loved fossil groups to the attention of collectors, a series of smaller contributions in the Dutch journal *Natuurhistorisch Maandblad*, entitled, 'Opmerkelijke Luiks-Limburgse Krijtfossielen', has been running since 1995. In addition, a profusely illustrated catalogue of fossils from the ENCI-HeidelbergCement Group quarry was published in 1998 by Jagt *et al.*; this was rapidly sold out, illustrating the need for such overviews. Fortunately, interest in other types of macrofossils has increased in recent years. For instance, thecideid and craniid brachiopods, monographed in 1959 and 1969, respectively (Backhaus, 1959; Krugtzer, 1969), are now being revised, and other articulate genera and species have been erected in

recent years (Simon, 2003, 2004a, b, 2005, 2007a, b, 2011; Simon & Owen, 2001). Additional groups have also witnessed a surge of interest, including crabs (Collins *et al.*, 1995; van Bakel *et al.*, 2003; Fraaije, 1996a, b, 1997, 2002a, b, 2003; Fraaije & van Bakel, 1998; Jagt *et al.*, 2010), brachiurans (Fraaije *et al.*, 2003), cirripedes (Jagt, 2011), crinoids, ophiuroids and asteroids (Jagt, 1999, 2000), and terrestrial plants and associated fungi (van der Ham *et al.*, 2001, 2003a, b, 2004, 2010, 2011; van der Ham & Dortangs, 2005). But the large majority, including bivalves, gastropods, nautiloids, sponges, scleractinian corals and trace fossils, are still waiting for a detailed taxonomic treatment.

All of these drawbacks we wish to counter by issuing a kind of identification guide to the macrofossil content of these strata, in several parts, to be published on a regular basis. Each special issue will comprise a certain number of peer-reviewed papers. Rather than publish a pocket-sized field guide, like the admirable ones issued by the Palaeontological Association (London), we have opted for such a series, because it offers authors the opportunity to erect new taxa and, in time, revise or modify chapters for successive issues.

The present volume contains a revision of the stratigraphy of all strata of Maastrichtian age in the type area of that stage and five chapters on macrofossils. These treat larger benthic foraminifera, sabellid and serpulid polychaete worms, scaphopod molluscs, coleoid cephalopods and ray-finned fish. Contributions to the present issue were reviewed by L. Cavin, J. Kriwet, C.P. Palmer and the present editors.

References

- Backhaus, E. 1959. Monographie der cretacischen Thecideidae (Brach.). *Mitteilungen aus dem Geologischen Staatsinstitut Hamburg*, **28**: 5-90.
- Bakel, B.W.M. van, Jagt, J.W.M., Fraaije, R.H.B. & Coole, Y. 2003. New data on the crab *Binkhorstia ubaghssii* (Late Maastrichtian; NE Belgium, SE Netherlands). *Contributions to Zoology*, **72**: 85-89.
- Brouwers, G.L.L.M., Jagt, J.W.M. & Mulder, E.W.A. (eds.). 2012. Fossiele cephalopoden van Nederland. *Staringia*, **13**: 1-227.
- Collins, J.S.H., Fraaye, R.H.B. & Jagt, J.W.M. 1995. Late Cretaceous anomurans and brachyurans from the Maastrichtian type area. *Acta Palaeontologica Polonica*, **40**: 165-210.
- Fraaije, R.H.B. 1996a. A new Tethyan migrant: *Cretachlorodius enciensis* n. gen., n. sp. (Crustacea, Decapoda), from the Maastrichtian type area. *Journal of Paleontology*, **70**: 293-296.
- Fraaije, R.H.B. 1996b. Two new crabs, *Graptocarcinus maastrichtensis*, and *Caloxanthus kuypersi* (Crustacea, Decapoda), from the type Maastrichtian of The Netherlands. *Journal of Paleontology*, **70**: 463-465.
- Fraaije, R.H.B. 1997. Late Cretaceous swimming crabs: radiation, migration, competition, and extinction. *Acta Geologica Polonica*, **46** (for 1996): 269-278.
- Fraaije, R.H.B. 2002a. The first record of albuneid crabs (Crustacea, Decapoda) from the Cretaceous. *Bulletin of the Mizunami Fossil Museum*, **29**: 69-73.
- Fraaije, R.H.B. 2002b. New calappid crabs (Crustacea, Decapoda, Brachyura) from the Late Maastrichtian of the Netherlands. *Journal of Paleontology*, **76**: 913-917.
- Fraaije, R.H.B. 2003. Evolution of reef-associated decapod crustaceans through time, with particular reference to the Maastrichtian type area. *Contributions to Zoology*, **72**: 119-130.
- Fraaije, R.H.B. & van Bakel, B.W.M. 1998. New raninid crabs (Crustacea, Decapoda, Brachyura) from the late Maastrichtian of the Netherlands. *Geologie en Mijnbouw*, **76**: 293-299.
- Fraaije, R.H.B., Schram, F.R. & Vonk, R. 2003. *Maastrichtiocaris rostratus* new genus and species, the first Cretaceous cycloid. *Journal of Paleontology*, **77**: 386-388.
- Ham, R.W.J.M. van der & Dortangs, R.W. 2005. Structurally preserved ascomycetous fungi from the Maastrichtian type area (NE Belgium). *Review of Palaeobotany and Palynology*, **136**: 48-62.

- Ham, R.W.J.M. van der, Dortangs, R.W., Herngreen, G.F.W. & van der Burgh, J. 2003a. *Brachiphyllum patens* (Miquel) comb. nov. (Cheirolepidiaceae?): remarkable conifer foliage from the Maastrichtian type area (Late Cretaceous, NE Belgium, SE Netherlands). *Review of Palaeobotany and Palynology*, **127**: 77-97.
- Ham, R.W.J.M. van der, de Wit, W., Zuidema, G. & van Birgelen, M. 1987. Zeeëgels uit het Krijt en Tertiair van Maastricht, Luik en Aken. Een atlas van de zeeëgels uit het Campaniëen, Maastrichtien en Daniën van Zuid-Limburg en aangrenzende delen van België en Duitsland. *Publicaties van het Natuurhistorisch Genootschap in Limburg*, **36**: 1-92.
- Ham, R., Indeherberge, L., Defour, E. & Meuris, R. 2006. Zee-egels uit het vuursteeneluvium van Hallembaye (Montagne Saint-Pierre). *Starlingia*, **12**: 1-59.
- Ham, R.W.J.M. van der, Jagt, J.W.M., Renkens, S. & van Konijnenburg-van Cittert, J.H.A. 2010. Seed-cone scales from the upper Maastrichtian document the last occurrence in Europe of the Southern Hemisphere conifer family Araucariaceae. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **291**: 469-473.
- Ham, R.W.J.M. van der & van Konijnenburg-van Cittert, J.H.A. 2003b. Rare conifers from the type area of the Maastrichtian (Upper Cretaceous, southeast Netherlands). *Scripta Geologica*, **126**: 111-119.
- Ham, R.W.J.M. van der, van Konijnenburg-van Cittert, J.H.A. & Burgh, J. van der. 2001. Taxodiaceous conifers from the Maastrichtian type area (Late Cretaceous, NE Belgium, SE Netherlands). *Review of Palaeobotany and Palynology*, **116**: 233-250.
- Ham, R.W.J.M. van der, van Konijnenburg-van Cittert, J.H.A., Kieft, B.N. & Walsmit Sachs, A. 2011. *Mosacaulis spinifer* gen. et sp. nov.: an enigmatic Maastrichtian plant. *Review of Palaeobotany and Palynology*, **168**: 51-67.
- Ham, R.W.J.M. van der, van Konijnenburg-van Cittert, J.H.A. & Nieuwenhuis, E.A.P.M. 2004. *Cunninghamites ubaghsii* (Taxodiaceae?) from the Maastrichtian type area (Late Cretaceous, SE Netherlands) rediscovered. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **74**: 89-96.
- Herman, J. 1977. Les sélaciens des terrains néocrétacées & paléocènes de Belgique & des contrées limitrophes. Éléments d'une biostratigraphie intercontinentale. *Mémoires pour l'Explication des Cartes géologiques et minières de la Belgique*, **15**: 1-401.
- Husson, D., Galbrun, B., Laskar, J., Hinov, L.A., Thibault, N., Gardin, S. & Locklair, R.E. 2011. Astronomical calibration of the Maastrichtian (Late Cretaceous). *Earth and Planetary Science Letters*, **305**: 328-340.
- Jagt, J.W.M. 1999. Late Cretaceous-Early Palaeogene echinoderms and the K/T boundary in the southeast Netherlands and northeast Belgium – Part 1: Introduction and stratigraphy. Part 2: Crinoids. *Scripta Geologica*, **116**: 1-57; 59-255.
- Jagt, J.W.M. 2000. Late Cretaceous-Early Palaeogene echinoderms and the K/T boundary in the southeast Netherlands and northeast Belgium – Part 3: Ophiuroids, with a chapter on: Early Maastrichtian ophiuroids from Rügen (northeast Germany) and Møn (Denmark) by Manfred Kutscher and John W.M. Jagt; Part 4: Echinoids; Part 5: Asteroids; Part 6: Conclusions. *Scripta Geologica*, **121**: 1-179; 181-375; 377-503; 505-577.
- Jagt, J.W.M. 2011. Charles Darwin and Joseph de Bosquet – brothers in barnacles: how diminutive crustaceans helped shape a theory. *Cretaceous Research*, **32**: 597-605.
- Jagt, J.W.M., Fraaije, R.H.B., van Bakel, B.W.M. & Artal, P. 2010. *Necrocarcinus ornatissimus* Forir, 1887, and *Prehepatus wernerii* Fraaye & Collins, 1987 (upper Maastrichtian, The Netherlands) revisited, with notes on other Cretaceous dynomenid crabs (Decapoda, Brachyura). *Crustaceana Monographs*, **11**: 173-195.
- Jagt, J.W.M., Leloux, J. & Dhondt, A.V. (eds.) 1998. Fossielen van de St. Pietersberg. *Grondboor & Hamer*, **52** [Limburgnummer 9B: Fossielen van de St. Pietersberg]: 93-157.
- Kennedy, W.J. 1987. The ammonite fauna of the type Maastrichtian with a revision of *Ammonites colligatus* Binkhorst, 1861. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **56** (for 1986): 151-267.
- Kruytzer, E.M. 1969. Le genre *Crania* du Crétacé supérieur et du Post-Maastrichtien de la province de Limbourg néerlandais (Brachiopoda, Inarticulata). *Publicaties van het Natuurhistorisch Genootschap in Limburg*, **19**: 4-42.

- Mulder, E.W.A. 2003. On latest Cretaceous tetrapods from the Maastrichtian type area. *Publicaties van het Natuurhistorisch Genootschap in Limburg*, **44**: 1-188.
- Schulp, A.S. 2006. On Maastricht mosasaurs. *Publicaties van het Natuurhistorisch Genootschap in Limburg*, **45**: 1-140.
- Simon, E. 2003. A new late Maastrichtian rhynchonellide brachiopod from the Kunrade Limestone facies near Maastricht (southern Limburg, The Netherlands). *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **73**: 137-148.
- Simon, E. 2004a. A new Late Maastrichtian species of *Cyanoia* (Terebratulida, Brachiopoda) from Belgium and The Netherlands. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **74**: 105-118.
- Simon, E. 2004b. A reappraisal of some large Late Maastrichtian brachiopods from Kunrade (southern Limburg, The Netherlands). *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **74**-Supplement: 121-137.
- Simon, E. 2005. New Lower Maastrichtian brachiopods (Gulpen Formation, Vijlen Member) from southern Limburg (The Netherlands). *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **75**: 127-165.
- Simon, E. 2007a. A new Late Maastrichtian species of *Gisilina* (Brachiopoda, Chlidonophoridae) from the Maastricht area (The Netherlands, Belgium) first illustrated by Faujas de Saint-Fond. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **77**: 131-140.
- Simon, E. 2007b. A new Late Maastrichtian species of *Isocrania* (Brachiopoda, Craniidae) from The Netherlands and Belgium. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **77**: 141-157.
- Simon, E. 2011. The late Maastrichtian cancellothyridid brachiopod *Terebratulina chrysalis* (Von Schlotheim, 1813) from the type Maastrichtian (southern Limburg, the Netherlands) and elsewhere in Europe. In: Jagt, J.W.M., Jagt-Yazykova, E.A. & Schins, W.J.H. (eds.), A tribute to the late Felder brothers – pioneers of Limburg geology and prehistoric archaeology. *Netherlands Journal of Geosciences*, **90**: 111-127.
- Simon, E. & Owen, E.F. 2001. A first step in the revision of the genus *Cretirhynchia* Pettitt, 1950. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **71**: 53-118.