



Naturalis Repository

Hyperdominance in the Amazonian Tree Flora

Hans ter Steege et al.

Downloaded from

<https://doi.org/10.1126/science.1243092>

Article 25fa Dutch Copyright Act (DCA) - End User Rights

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with consent from the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available following a reasonable period after the work was first published, provided that reference is made to the source of the first publication of the work.

This publication is distributed under the Naturalis Biodiversity Center 'Taverne implementation' programme. In this programme, research output of Naturalis researchers and collection managers that complies with the legal requirements of Article 25fa of the Dutch Copyright Act is distributed online and free of barriers in the Naturalis institutional repository. Research output is distributed six months after its first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and copyrights owner(s) of this work. Any use of the publication other than authorized under this license or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the department of Collection Information know, stating your reasons. In case of a legitimate complaint, Collection Information will make the material inaccessible. Please contact us through email: collectie.informatie@naturalis.nl. We will contact you as soon as possible.

Hyperdominance in the Amazonian Tree Flora

Hans ter Steege* *et al.*

READ THE FULL ARTICLE ONLINE

<http://dx.doi.org/10.1126/science.1243092>



Cite this article as H. ter Steege *et al.*, *Science* **342**, 1243092 (2013). DOI: 10.1126/science.1243092

Introduction: Recent decades have seen a major international effort to inventory tree communities in the Amazon Basin and Guiana Shield (Amazonia), but the vast extent and record diversity of these forests have hampered an understanding of basinwide patterns. To overcome this obstacle, we compiled and standardized species-level data on more than half a million trees in 1170 plots sampling all major lowland forest types to explore patterns of commonness, rarity, and richness.

Methods: The ~6-million-km² Amazonian lowlands were divided into 1° cells, and mean tree density was estimated for each cell by using a loess regression model that included no environmental data but had its basis exclusively in the geographic location of tree plots. A similar model, allied with a bootstrapping exercise to quantify sampling error, was used to generate estimated Amazon-wide abundances of the 4962 valid species in the data set. We estimated the total number of tree species in the Amazon by fitting the mean rank-abundance data to Fisher's log-series distribution.

Results: Our analyses suggest that lowland Amazonia harbors 3.9×10^{11} trees and ~16,000 tree species. We found 227 "hyperdominant" species (1.4% of the total) to be so common that together they account for half of all trees in Amazonia, whereas the rarest 11,000 species account for just 0.12% of trees. Most hyperdominants are habitat specialists that have large geographic ranges but are only dominant in one or two regions of the basin, and a median of 41% of trees in individual plots belong to hyperdominants. A disproportionate number of hyperdominants are palms, Myristicaceae, and Lecythidaceae.

Discussion: The finding that Amazonia is dominated by just 227 tree species implies that most biogeochemical cycling in the world's largest tropical forest is performed by a tiny sliver of its diversity. The causes underlying hyperdominance in these species remain unknown. Both competitive superiority and widespread pre-1492 cultivation by humans are compelling hypotheses that deserve testing. Although the data suggest that spatial models can effectively forecast tree community composition and structure of unstudied sites in Amazonia, incorporating environmental data may yield substantial improvements. An appreciation of how thoroughly common species dominate the basin has the potential to simplify research in Amazonian biogeochemistry, ecology, and vegetation mapping. Such advances are urgently needed in light of the >10,000 rare, poorly known, and potentially threatened tree species in the Amazon.

FIGURES AND TABLES IN THE FULL ARTICLE

Fig. 1. A map of Amazonia showing the location of the 1430 ATDN plots that contributed data to this paper.

Fig. 2. A rank-abundance diagram of 4962 tree species extrapolated to estimate the size of the Amazon tree flora.

Fig. 3. Characteristics of hyperdominant tree species of the Amazon.

Fig. 4. Proportions of hyperdominance by region and forest type.

Fig. 5. Distribution maps of three hyperdominant Amazon tree species.

Table 1. Population characteristics of the 20 most abundant tree species of the Amazon.

Table 2. Hyperdominance by region and forest type.

SUPPLEMENTARY MATERIALS


Supplementary Text

Figs. S1 to S12

Tables S1 to S3

Appendices S1 to S4

References (53–67)



A map of Amazonia showing the location of the 1430 Amazon Tree Diversity Network (ATDN) plots that contributed data to this paper. The white polygon marks our delimitation of the study area and consists of 567 1° grid cells (area = 6.29 million km²). Orange circles indicate plots on terra firme; blue squares, plots on seasonally or permanently flooded terrain (várzea, igapó, swamps); yellow triangles, plots on white-sand podzols; gray circles, plots only used for tree density calculations. Background is from Visible Earth. CA, central Amazonia; EA, eastern Amazonia; GS, Guyana Shield; SA, southern Amazonia; WAN, northern part of western Amazonia; WAS, southern part of western Amazonia. More details are shown in figs. S1 to S3.

Lists of authors and affiliations are available in the full article online.

*Corresponding author. E-mail: hans.tersteege@naturalis.nl

Hyperdominance in the Amazonian Tree Flora

Hans ter Steege,^{1,2*} Nigel C. A. Pitman,^{3,4} Daniel Sabatier,⁵ Christopher Baraloto,⁶ Rafael P. Salomão,⁷ Juan Ernesto Guevara,⁸ Oliver L. Phillips,⁹ Carolina V. Castilho,¹⁰ William E. Magnusson,¹¹ Jean-François Molino,⁵ Abel Monteagudo,¹² Percy Núñez Vargas,¹³ Juan Carlos Montero,^{14,11} Ted R. Feldpausch,^{9,15} Eurídice N. Honorio Coronado,^{16,9} Tim J. Killeen,¹⁷ Bonifacio Mostacedo,¹⁸ Rodolfo Vasequez,¹² Rafael L. Assis,^{11,19} John Terborgh,³ Florian Wittmann,²⁰ Ana Andrade,²¹ William F. Laurance,²² Susan G. W. Laurance,²² Beatriz S. Marimon,²³ Ben-Hur Marimon Jr.,²³ Ima Célia Guimarães Vieira,²⁴ Iêda Leão Amaral,²⁵ Roel Brien, ⁹ Hernán Castellanos,²⁶ Dairon Cárdenas López,²⁷ Joost F. Duivenvoorden,²⁸ Hugo F. Mogollón,²⁹ Francisca Dionízia de Almeida Matos,¹¹ Nállarett Dávila,³⁰ Roosevelt García-Villacorta,^{31,32} Pablo Roberto Stevenson Diaz,³³ Flávia Costa,¹¹ Thaise Emilio,¹¹ Carolina Levis,¹¹ Juliana Schietti,¹¹ Priscila Souza,¹¹ Alfonso Alonso,³⁴ Francisco Dallmeier,³⁴ Alvaro Javier Duque Montoya,³⁵ Maria Teresa Fernandez Piedade,¹¹ Alejandro Araujo-Murakami,³⁶ Luzmila Arroyo,³⁶ Rogerio Gribel,³⁷ Paul V. A. Fine,⁸ Carlos A. Peres,³⁸ Marisol Toledo,³⁹ Gerardo A. Aymard C.,⁴⁰ Tim R. Baker,⁹ Carlos Cerón,⁴¹ Julien Engel,⁴² Terry W. Henkel,⁴³ Paul Maas,¹ Pascal Petronelli,⁴⁴ Juliana Stropp,⁴⁵ Charles Eugene Zartman,¹¹ Doug Daly,⁴⁶ David Neill,⁴⁷ Marcos Silveira,⁴⁸ Marcos Ríos Paredes,⁴⁹ Jerome Chave,⁵⁰ Diógenes de Andrade Lima Filho,¹¹ Peter Møller Jørgensen,⁵¹ Alfredo Fuentes,^{52,51} Jochen Schöngart,²⁰ Fernando Cornejo Valverde,⁵³ Anthony Di Fiore,⁵⁴ Eliana M. Jimenez,⁵⁵ Maria Cristina Peñuela Mora,⁵⁵ Juan Fernando Phillips,⁵⁶ Gonzalo Rivas,⁵⁷ Tinde R. van Andel,¹ Patricio von Hildebrand,⁵⁶ Bruce Hoffman,¹ Eglée L. Zent,⁵⁸ Yaduvinder Malhi,⁵⁹ Adriana Prieto,⁶⁰ Agustín Rudas,⁶⁰ Ademir R. Ruschell,⁶¹ Natalino Silva,⁶² Vincent Vos,⁶³ Stanford Zent,⁵⁸ Alexandre A. Oliveira,⁶⁴ Angela Cano Schutz,³³ Therany Gonzales,⁶⁵ Marcelo Trindade Nascimento,⁶⁶ Hirma Ramirez-Angulo,⁶⁷ Rodrigo Sierra,⁶⁸ Milton Tirado,⁶⁸ María Natalia Umaña Medina,³³ Geertje van der Heijden,^{69,70} César I. A. Vela,⁷¹ Emilio Vilanova Torre,⁶⁷ Corine Vriesendorp,⁴ Ophelia Wang,⁷² Kenneth R. Young,⁷³ Claudia Baider,^{64,74} Henrik Balslev,⁷⁵ Cid Ferreira,¹¹ Italo Mesones,⁸ Armando Torres-Lezama,⁷⁶ Ligia Estela Urrego Giraldo,³⁵ Roderick Zagt,⁷⁷ Miguel N. Alexiades,⁷⁸ Lionel Hernandez,²⁶ Isau Huamantupa-Chuquimaco,⁷⁹ William Milliken,⁸⁰ Walter Palacios Cuenca,⁸¹ Daniela Pauletto,⁸² Elvis Valderrama Sandoval,^{83,84} Luis Valenzuela Gamarra,¹² Kyle G. Dexter,⁸⁵ Ken Feeley,^{86,87} Gabriela Lopez-Gonzalez,⁹ Miles R. Silman⁸⁸

The vast extent of the Amazon Basin has historically restricted the study of its tree communities to the local and regional scales. Here, we provide empirical data on the commonness, rarity, and richness of lowland tree species across the entire Amazon Basin and Guiana Shield (Amazonia), collected in 1170 tree plots in all major forest types. Extrapolations suggest that Amazonia harbors roughly 16,000 tree species, of which just 227 (1.4%) account for half of all trees. Most of these are habitat specialists and only dominant in one or two regions of the basin. We discuss some implications of the finding that a small group of species—less diverse than the North American tree flora—accounts for half of the world's most diverse tree community.

Much remains unknown about the Amazonian flora, the richest assemblage of plant species on Earth. Tree inventories carried out over the past two decades have helped improve our understanding of regional-scale patterns of distribution and abundance in Amazonian tree communities, but similar advances at the basin-wide scale remain scarce. Scientists still do not know how many tree species occur in the Amazon (*1*), how many tree species have been recorded to date in the Amazon, how those species are distributed across the basin, and in what regions or forest types they are rare or common. So uncertain are patterns at the largest scales that even the simplest question of all—what is the most common tree species in the Amazon?—has never been addressed in the scientific literature, much less answered.

In practical terms, this lack of basic information means that the largest pool of tropical carbon on Earth remains a black box for ecologists. Also, given that abundance and frequency are the primary currencies of conservation status, there is essentially no information on which Amazonian tree species face the most severe threats of extinction and where to protect them. Although ecologists have an ever-larger toolbox of methods to extrapolate from local surveys to larger-scale patterns, well-known barriers to sampling, identifying, and putting valid names on tropical trees have long been assumed to make Amazon-wide extrapolations impossible.

We challenged those assumptions with a wide-ranging assessment of the composition and biogeography of Amazonian tree communities (*2*).

We compiled stem density and species abundance data from 1170 tree inventory plots across the Amazon (Fig. 1), well distributed among all regions and major forest types (table S1 and figs. S1 to S3), to generate basin-wide estimates of the abundance, frequency, and spatial distribution of thousands of Amazonian tree species.

Results

A Rank-Abundance Distribution for Amazonian Trees

The plots contained a total of 4962 valid species, 810 genera, and 131 families of trees [free-standing stems ≥ 10 cm in diameter at breast height (dbh)]. By using stem density and species abundance data collected in the individual plots, we constructed a spatial model that yielded estimated basin-wide population sizes for every valid species in the data set. The rank-abundance distribution (RAD) of these data (Fig. 2) offers four important insights regarding Amazonian tree communities.

First, it provides the most precise estimates yet of two numbers that have been debated for decades: How many trees and how many tree species occur in the ~ 6 -million- km^2 landscape of Amazonia (*1, 3–5*). Our estimate of tree density yielded a total of 3.9×10^{11} individual trees and a median tree density of 565 trees/ha (fig. S4). Assuming that our population size estimates for the common species are reasonable (fig. S5) and that Fisher's log-series model fits our data (table S2 and figs. S6 and S7) (*1*), we estimated the total number of tree species in the Amazon to be about 16,000 (Fig. 2). A second estimate based on the Fisher's alpha scores of all plots yielded a similar figure: 15,182 species (fig. S8).

Second, the RAD suggests that just 227 (1.4%) of the estimated 16,000 species account for half of all individual trees in Amazonia. We refer to these species, all of which have estimated populations of $>3.7 \times 10^8$ trees, as hyperdominant species (see a list of the 20 most abundant species in Table 1 and a full list in appendix S1). These hyperdominant species form the basis of the tree communities in individual plots as well, accounting for a median of 41% of trees (range = 0 to 94%, fig. S9) and 32% of species (range = 0 to 78%) per plot (fig. S9).

Third, all species ranking in abundance from 5000 to 16,000 are very rare. These species in the tail of the RAD have total populations of $<10^6$ individuals and together account for just 0.12% of all trees in Amazonia. Although some of these species may be treelets or climbers rarely reaching tree stature or “vagrants” spilling over from extra-Amazonian biomes such as the Cerrado and Andes, thousands must be Amazonian endemics that run a high risk of going extinct, even before they can be found and described by biologists. The rarest 5800 species have estimated population sizes of <1000 , which is sufficient to classify those that are endemic as globally threatened (*6*). Together, these taxa (the rarest 36% of species)

account for just 0.0003% of all trees in Amazonia. Given the extreme unlikelihood of locating a fertile individual of one of these species, we believe that discovering and describing the unknown portion of Amazonian biodiversity will be a long-term struggle with steeply diminishing returns and not an easy linear process (7). Indeed, the RAD suggests that floras of even well-collected areas may remain half-finished for decades. For example, our model predicts that ~4500 tree species occur in the Guianas (fig. S10), but centuries of collecting there have yielded just half that number (8). Some of these species may be present among the unidentified species of our plots or as undescribed specimens in herbaria (9), but the majority may yet have to be collected.

Fourth, there are strong similarities between theoretical models of tree species richness in the Amazon (1) and our distribution of species abundances based on empirical data. For example, Hubbell *et al.* (1) used a log-series distribution to predict that the most common species in the Amazon should account for 1.39% of all trees. This is very close to our estimate for the most common species in our data set, the palm *Euterpe precatoria* (1.32%). Our empirical estimate of

Fisher's alpha for the Amazon (fig. S8) is also extremely close to Hubbell *et al.*'s modeled prediction [754 versus 743 in (1)]. Although these strong correlations between predictions and our data set suggest that the log series may offer useful insights on the most poorly known tree species in the Amazon (e.g., the number of undescribed taxa), they should not necessarily be interpreted as evidence for any one theory of how these tree communities are structured (10, 11).

Hyperdominant Patterns in Regions, Forest Types, and Taxonomic Groups

We examined species' geographic ranges and abundances by plots, regions, and forest types to explore how hyperdominant species differ from other taxa, as a first step toward understanding what makes them so successful. Hyperdominant species have larger ranges than other taxa (Fig. 3A) and reach greater maximum relative abundances in plots (Fig. 3B). Most hyperdominant species (121 out of 227) are habitat specialists (Fig. 3C) [i.e., they show a strong preference for one of the five major Amazonian forest types: terra firme (53 spp.), várzea (26), white-sand forest (16), swamps (14), and igapó (12)]. Likewise, most are only

dominant within one or two forest types. When the study area was divided into six regions (Guiana Shield and northwest, southwest, south, east, and central Amazonia), most hyperdominant species (73%) were found to be dominant within only one or two regions (Table 2).

It is thus important to emphasize that, although the Amazonian RAD is dominated by a small suite of species, most of those species are only dominant in certain forest types and in certain regions of the basin. Just one species qualified as dominant in all six regions (*Eschweilera coriacea*), no species were dominant in all five forest types, and only four species were dominant in four forest types (*E. precatoria*, *Oenocarpus bataua*, *Licania apetala*, and *Euterpe oleracea*). Much more representative of the 227 hyperdominant species are taxa like *Siparuna decipiens* (112th largest population size overall), only dominant in terra firme forests in southwest Amazonia, and *Eperua falcata* (13th), only dominant in the Guiana Shield. Indeed, 58% of hyperdominant species qualify as both dominant in one or two regions and dominant in one or two forest types.

Within each region, an even smaller number of species (75 to 163) typically accounts for

¹Naturalis Biodiversity Center, Leiden, Netherlands. ²Ecology and Biodiversity Group, Utrecht University, Netherlands. ³Center for Tropical Conservation, Nicholas School of the Environment, Duke University, Durham, NC 27708, USA. ⁴The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA. ⁵Institut de Recherche pour le Développement, UMR Architecture, Fonctionnement et Évolution des plantes, Montpellier, France. ⁶Institut National de la Recherche Agronomique, UMR Ecologie des Forêts de Guyane, French Guiana. ⁷Ministério da Ciência, Tecnologia e Inovação/Museu Paraense Emílio Goeldi–Cordenadoria de Botânica, Belém, Brazil. ⁸Department of Integrative Biology, University of California, Berkeley, CA 94720–3140, USA. ⁹School of Geography, University of Leeds, Leeds LS2 9JT, UK. ¹⁰Embrapa Roraima, Boa Vista, RR, Brazil. ¹¹Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brazil. ¹²Jardín Botánico de Missouri, Oxpampa, Peru. ¹³Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru. ¹⁴BOLFOR (Bolivia Sustainable Forest Management Project), Cuarto Anillo, esquina Av. 2 de Agosto, Casilla 6204, Santa Cruz, Bolivia. ¹⁵College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4RJ, UK. ¹⁶Instituto de Investigaciones de la Amazonia Peruana, Av. José A. Quiñones km. 2.5, Iquitos, Peru. ¹⁷World Wildlife Fund (WWF), Washington, DC 20037, USA. ¹⁸Universidad Autónoma Gabriel René Moreno, Facultad de Ciencias Agrícolas, Santa Cruz, Bolivia. ¹⁹Department of Ecology and Natural Resource Management, Norwegian University of Life Sciences (UMB), Aas, Norway. ²⁰Max Planck Institute for Chemistry, Biogeochemistry, Mainz, Germany. ²¹Instituto Nacional de Pesquisas da Amazônia, Projeto Dinâmica Biológica de Fragmentos Florestais, Manaus, AM, Brazil. ²²Centre for Tropical Environmental and Sustainability Science (TESS) and School of Marine and Tropical Biology, James Cook University, Cairns, Queensland, Australia. ²³Universidade do Estado de Mato Grosso, Nova Xavantina, MT, Brazil. ²⁴Museu Paraense Emílio Goeldi, Belém, PA, Brazil. ²⁵Projeto TEAM (Tropical Ecology Assessment and Monitoring)–Manaus, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, AM, Brazil. ²⁶Universidad Nacional Experimental de Guayana, Puerto Ordaz, Bolívar, Venezuela. ²⁷SINCHI (Instituto Amazónico de Investigaciones Científicas) Herbario Amazónico Colombiano, Calle 20 No. 5, 44 Bogotá, Colombia. ²⁸Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, Netherlands. ²⁹Endangered Species Coalition, 8530 Geren Road, Silver Spring, MD 20901, USA. ³⁰Universidade Estadual de Campinas, Campinas,

São Paulo, Brazil. ³¹Institute of Molecular Plant Sciences, University of Edinburgh, Mayfield Rd, Edinburgh EH9 3JH, UK. ³²Royal Botanic Garden of Edinburgh, 20a Inverleith Row, Edinburgh EH3 5LR, UK. ³³Laboratório de Ecologia de Bosques Tropicales y Primatología, Universidad de los Andes, Bogotá DF, Colombia. ³⁴Smithsonian Conservation Biology Institute, National Zoological Park MRC 0705, Washington, DC 20013, USA. ³⁵Universidad Nacional de Colombia, Departamento de Ciencias Forestales, sede Medellín, Colombia. ³⁶Museo de Historia Natural Noel Kempff Mercado, Santa Cruz, Bolivia. ³⁷Instituto de Pesquisas Jardim Botânico do Rio de Janeiro, Rio de Janeiro, RJ, Brazil. ³⁸School of Environmental Sciences, University of East Anglia, Norwich, UK. ³⁹Instituto Boliviano de Investigación Forestal, Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia. ⁴⁰UNELLEZ (Universidad Nacional Experimental de los Llanos Occidentales Ezekiel Zamora)–Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT), estado Portuguesa, 3350 Venezuela. ⁴¹Herbario Alfredo Paredes (QAP), Universidad Central del Ecuador, Ap. Postal 17.01.2177, Quito, Ecuador. ⁴²CNRS, UMR Ecologie des Forêts de Guyane, French Guiana. ⁴³Department of Biological Sciences, Humboldt State University, Arcata, CA 95521, USA. ⁴⁴La Recherche Agronomique pour le Développement (CIRAD), UMR Ecofog, Kourou, French Guiana. ⁴⁵Land Resource and Management Unit, Joint Research Centre of the European Commission, Via Enrico Fermi 2749, TP 440, I-21027 Ispra (VA), Italy. ⁴⁶New York Botanical Garden, Bronx, New York, NY 10458–5126, USA. ⁴⁷Universidad Estatal Amazónica, Puyo, Ecuador. ⁴⁸Museu Universitário, Universidade Federal do Acre, Rio Branco, AC, Brazil. ⁴⁹Servicios de Biodiversidad EIRL, Iquitos, Peru. ⁵⁰CNRS and Université Paul Sabatier, UMR 5174 EDB, 31000 Toulouse, France. ⁵¹Missouri Botanical Garden, Post Office Box 299, St. Louis, MO 63166–0299, USA. ⁵²Herbario Nacional de Bolivia, Casilla 10077 Correo Central, La Paz, Bolivia. ⁵³Andes to Amazon Biodiversity Program, Madre de Dios, Peru. ⁵⁴Department of Anthropology, University of Texas at Austin, Austin, TX 78712, USA. ⁵⁵Grupo de Ecología de Ecosistemas Terrestres Tropicales, Universidad Nacional de Colombia Sede Amazonia, Leticia, Amazonas, Colombia. ⁵⁶Fundación Puerto Rastrojo, Cra 10 No. 24-76 Oficina 1201, Bogotá, Colombia. ⁵⁷Wildlife Ecology and Conservation and Quantitative Spatial Ecology, University of Florida, Gainesville, FL 32611, USA. ⁵⁸Laboratory of Human Ecology, Instituto Venezolano de Investigaciones Científicas, Ado 20632, Caracas 1020-A, Venezuela. ⁵⁹Environmental Change Institute, School of Ge-

ography and the Environment, University of Oxford, Oxford, UK. ⁶⁰Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia. ⁶¹Embrapa Amazônia Oriental, Belém, PA, Brazil. ⁶²UFRA (Universidade Federal Rural da Amazônia), Belém, PA, Brazil. ⁶³Universidad Autónoma del Beni, Riberalta, Bolivia. ⁶⁴Universidade de São Paulo, Instituto de Biociências, Departamento Ecologia, Cidade Universitária, São Paulo, SP, Brazil. ⁶⁵Amazon Center for Environmental Education and Research Foundation, Jirón Cusco No. 370, Puerto Maldonado, Madre de Dios, Peru. ⁶⁶Laboratório de Ciências Ambientais, Universidade Estadual do Norte Fluminense, Campos dos Goytacazes, RJ 28013–620, Brazil. ⁶⁷INDEFOR (Research Institute for Forestry Development), Universidad de los Andes, Mérida, Venezuela. ⁶⁸Geoinformática y Sistemas, Cia. Ltda. (GeolS), Quito, Ecuador. ⁶⁹Department of Biological Sciences, University of Wisconsin–Milwaukee, Milwaukee, WI 53202, USA. ⁷⁰Smithsonian Tropical Research Institute, Apartado Postal 0843-03092, Panama City, Panama. ⁷¹Facultad de Ciencias Forestales y Medio Ambiente, Universidad Nacional de San Antonio Abad del Cusco, Jr. San Martín 451, Puerto Maldonado, Madre de Dios, Peru. ⁷²Northern Arizona University, Flagstaff, AZ 86011, USA. ⁷³Geography and the Environment, University of Texas, Austin, TX 78712, USA. ⁷⁴The Mauritius Herbarium, Agricultural Services, Ministry of Agro-Industry and Food Security, Reduit, Mauritius. ⁷⁵University of Aarhus, Aarhus, Denmark. ⁷⁶Universidad de los Andes, Mérida, Venezuela. ⁷⁷Tropenbos International, Wageningen, Netherlands. ⁷⁸School of Anthropology and Conservation, Marlowe Building, University of Kent, Canterbury, Kent CT2 7NR, UK. ⁷⁹Herbario CUZ, Universidad Nacional San Antonio Abad del Cusco, Cusco, Peru. ⁸⁰Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK. ⁸¹Universidad Técnica del Norte/Herbario Nacional del Ecuador, Quito, Ecuador. ⁸²Serviço Florestal Brasileiro, Santarém, PA, Brazil. ⁸³Department of Biology, University of Missouri–Saint Louis, R 102 Research, St. Louis, MO 63121, USA. ⁸⁴Facultad de Biología, Universidad Nacional de la Amazonia Peruana, Pevás Sta cdra, Iquitos, Peru. ⁸⁵School of Geosciences, University of Edinburgh, 201 Crew Building, King's Buildings, Edinburgh EH9 3JN, UK. ⁸⁶Department of Biological Sciences, Florida International University, Miami FL 33199, USA. ⁸⁷Fairchild Tropical Botanic Garden, Coral Gables FL 33156, USA. ⁸⁸Biology Department and Center for Energy, Environment and Sustainability, Wake Forest University, Winston-Salem, NC 27106, USA.

*Corresponding author. E-mail: hans.tersteeg@naturalis.nl

50% of all individual trees, and most of these regional dominants are also hyperdominant species (Fig. 4A). For example, the data suggest that half of all individual trees in southwest Amazonia belong to just 76 species, 50 of which are also hyperdominant species. The same pattern holds for forest types, which are individually dominated by 25 to 195 species (Fig. 4B). Half of all individual trees in white-sand forests belong to just 25 species, 15 of which are also hyperdominant species. Because most hyperdominant species are only dominant in one or two regions or forest types, in any single region or forest type the majority of the 227 hyperdominant species are not locally dominant.

Given these results, it seems likely that the basinwide patterns of dominance we describe here arise in part from regional-scale patterns of dominance described previously at various sites in upper Amazonia (12, 13). There is substantial compositional overlap between Pitman *et al.*'s (12) "oligarchies" in Peru and Ecuador and our hyperdominant species, even though those authors' plots represent just 2.1% of the full Ama-

zon Tree Diversity Network (ATDN) data set and only include terra firme forests. Sixty-eight "oligarchs" of (12) are on the list of 227 hyperdominant species, including 8 of the top 10 most common hyperdominants. The 250 oligarchic species in (12) account for 26.9% of all trees in Amazonia, according to the RAD in Fig. 2. These results suggest that the regional-scale and Amazon-wide patterns derive from similar processes.

Hyperdominants are more frequent in some families (table S3). Arecaceae, Myristicaceae, and Lecythidaceae have many (~four to five times) more hyperdominant species than expected by chance, whereas Myrtaceae, Melastomataceae, Lauraceae, Annonaceae, and Rubiaceae have fewer, probably because many of their species are shrubs or treelets that do not reach our 10-cm-diameter cutoff. In Fabaceae, the most abundant and most diverse family in the data set, the observed number of hyperdominant species is not significantly different from the expected.

We observed a negative relationship between the number of species in a genus and the frequency of hyperdominant species (fig. S11). This

pattern has been observed in several plant communities worldwide, and scientists have yet to determine whether it is ecologically informative or an artifact of rank-based taxonomy (14, 15). The 227 hyperdominant species belong to 121 genera, and 68 of these contain more hyperdominants than expected by chance (appendix S3). The highest number of hyperdominant species is found in moderately diverse *Eschweilera* (52 species overall; 2.4 hyperdominant species expected versus 14 observed), also the most abundant genus in the ATDN data set (5.2% of all stems). Given that the families and genera mentioned here dominate Amazonian forests, it remains a key goal to determine why some achieve dominance with a large number of mostly rare species (e.g., *Inga*, Sapotaceae) whereas others do so with a small number of common species (palms), differences that may result from variation in speciation and extinction rates (14–17). Although genetics data may reveal some hyperdominant species to be species complexes, there is not yet enough knowledge on how widespread such complexes are, where they are located along our RAD,




Fig. 1. A map of Amazonia showing the location of the 1430 ATDN plots that contributed data to this paper. The white polygon marks our delimitation of the study area [with subregions after (33)] and consists of 567 1°-grid cells (area = 6.29 million km²). Orange circles indicate plots on terra firme; blue squares, plots on seasonally or permanently flooded terrain

(várzea, igapó, and swamps); yellow triangles, plots on white-sand podzols; gray circles, plots only used for tree density calculations. Background is from Visible Earth (52). CA, central Amazonia; EA, eastern Amazonia; GS, Guyana Shield; SA, southern Amazonia; WAN, northern part of western Amazonia; WAS, southern part of western Amazonia. More details are shown in figs. S1 to S3.

and to what degree they could alter the patterns described here [(18) and references therein].

Discussion

Exploring Potential Causes for Hyperdominance

We found no evidence that two key functional traits for trees, seed mass and wood density,

vary consistently with hyperdominance. The 227 hyperdominant species include both shade-tolerant, typically large-seeded climax species with dense wood (e.g., *Chlorocardium rodiei*, *Clathrotropis* spp., and *Eperua* spp.) and shade-intolerant, small-seeded pioneers with light wood (e.g., *Cecropia* spp., *Jacaranda copaia*, and *Laetia*

procera). Given that most hyperdominant species attain very high local densities (>60 trees/ha) somewhere in the plot network, we predict that they will be found to be disproportionately resistant to pathogens, specialist herbivores, and other sources of frequency-dependent mortality (19, 20).




Fig. 2. A rank-abundance diagram of 4962 tree species extrapolated to estimate the size of the Amazon tree flora. The mean estimated Amazon-wide population sizes of 4962 tree species are shown as a solid line, and the dotted line is an extrapolation of the distribution used to estimate the total number of tree species in Amazonia.

Table 1. Population characteristics of the 20 most abundant tree species of the Amazon. Mean estimated population sizes of the 20 most abundant tree species in Amazonia and the empirical abundance and frequency data on

which the estimates were based. Median values for the 207 other hyperdominant species and for the 4735 other valid species in the data set are provided for comparison. Data on all species can be found in appendix S1.

Species	Mean estimated population in the Amazon	SD estimated population (%)	No. trees in data set	% of all plots where present	Maximum abundance recorded (trees/ha)
<i>Euterpe precatoria</i>	5.21×10^9	9.9	5903	32.7	168
<i>Protium altissimum</i>	5.21×10^9	18.0	5889	15.6	128
<i>Eschweilera coriacea</i>	5.00×10^9	5.6	9047	47.9	28
<i>Pseudolmedia laevis</i>	4.30×10^9	8.9	5285	36.1	121
<i>Iriartea deltoidea</i>	4.07×10^9	13.1	8405	18.5	169
<i>Euterpe oleracea</i>	3.78×10^9	17.5	8572	7.4	397
<i>Oenocarpus bataua</i>	3.71×10^9	10.7	4767	29.9	108
<i>Trattinnickia burserifolia</i>	2.78×10^9	29.4	3023	10	125
<i>Socratea exorrhiza</i>	2.68×10^9	10.8	863	28.6	82
<i>Astrocaryum murumuru</i>	2.41×10^9	11.2	5748	16.7	325
<i>Brosimum lactescens</i>	2.28×10^9	10.0	2234	28.2	106
<i>Protium heptaphyllum</i>	2.13×10^9	32.2	1365	11.3	169
<i>Eperua falcata</i>	1.95×10^9	15.8	1898	10.9	266
<i>Hevea brasiliensis</i>	1.91×10^9	15.5	6031	14.8	179
<i>Eperua leucantha</i>	1.84×10^9	32.3	1453	1.4	282
<i>Helicostylis tomentosa</i>	1.79×10^9	25.6	1948	36.5	89
<i>Attalea butyracea</i>	1.78×10^9	16.2	2561	5.8	73
<i>Rinorea guianensis</i>	1.69×10^9	18.6	1243	13.7	182
<i>Licania heteromorpha</i>	1.57×10^9	14.4	2483	35	173
<i>Metrodorea flavida</i>	1.55×10^9	14.7	1326	7.7	128
Median of other hyperdominant species	5.79×10^8		808	11.4	60
Median of non-hyperdominant species	1.11×10^7		15	0.5	5

Widespread pre-1492 cultivation by humans is a compelling hypothesis to explain hyperdominance (21). Numerous hyperdominant species are widely used by modern indigenous groups (*Hevea brasiliensis*, *Theobroma cacao*, and many palms), and some are associated with pre-Columbian settlements (*Attalea butyracea*, *A. phalerata*, *Mauritia flexuosa*) (22–26). On the other hand, most hyperdominant species are not commonly cultivated; many of the most commonly used hyperdominants (palms) belong to a family that appears to have been dominant in tropical South America since the Paleocene (27), and large portions of the Amazon Basin do not appear to have been heavily cultivated before 1492 (28).

Testing the Validity of the Model Predictions

A fundamental assumption of our analyses is that the population-size estimates generated by

the loess model were reasonably accurate for the most abundant species. This assumption is disputable for a few reasons: (i) The data set is very small compared with the community to which it was extrapolated; (ii) tree plots were not distributed randomly across the study area; (iii) trees were identified by many different research teams; and (iv) no environmental data were used by the model, even though many species in the ATDN data set are known to respond to environmental heterogeneity in the study area. A fifth problem makes the assumption especially difficult to test: (v) the fact that a basinwide population size has not been empirically determined for any Amazonian tree species, which precludes a comparison between projected and observed values. Here, we address these shortcomings by attempting to quantify the error that each could introduce into our results.




Fig. 3. Characteristics of hyperdominant tree species of the Amazon. (A) Hyperdominant species (red) have larger geographic ranges than other species (gray), (B) reach higher maximum relative abundances in individual plots, and (C) are more likely to be habitat specialists.

Table 2. Hyperdominance by region and forest type. The number of hyperdominant species that are also dominant in individual forest types and regions. Most hyperdominants only dominate a single forest type, and most are dominant in one or two regions.

		No. forest types where dominant						
		0	1	2	3	4	5	total
No. regions where dominant	0	3	3	0	0	0	0	6
	1	18	47	8	0	0	0	73
	2	12	65	12	3	0	0	92
	3	2	17	4	1	1	0	25
	4	0	9	3	5	0	0	17
	5	0	6	1	4	2	0	13
	6	0	1	0	0	0	0	1
total		35	148	28	13	3	0	227

To test how sampling intensity and the geographic distribution of plots (problems i and ii) affected the estimated population sizes of hyperdominant species, we recorded the frequency with which the 227 hyperdominants qualified as hyperdominant in the 500 runs of the bootstrap exercise described in the methods section. Most species (137, 60% of the total) qualified as hyperdominants in 90 to 100% of runs, whereas 207 species (91.2%) qualified as hyperdominants in more than half of runs (fig. S12A). Median (fig. S12B) and mean (fig. S12C) ranks for the 500 runs showed high stability.

In bootstrap runs for which a given hyperdominant species did not qualify among the top 227 species, it rarely qualified as rare. The lowest median rank observed for a hyperdominant species in the 500 bootstrap runs was 275, and hyperdominant species never ranked lower than 1000th (i.e., ranks 1000 to 4790). These analyses provide strong evidence that the identities and estimated population sizes of the hyperdominant species remain stable and predictable with varying levels of sampling intensity and geographic bias.

Taxonomic and identification problems (problem iii) are widespread in Amazonian tree inventories. However, two independent lines of evidence suggest that resolving these problems will not fundamentally alter the patterns described for hyperdominant species.

First, we observed a consistent relationship in the ATDN data set between the abundance of a species and the likelihood that it had been identified with a valid name. The percentage of identified species in individual plots was significantly higher than that of unidentified species-level taxa (87 versus 13% stems/ha, analysis of variance, $F_{5,22774} = 22,774$, $P < 0.001$). Furthermore, very common morphospecies are very infrequent in the ATDN data set. Only 48 of the 1170 ATDN plots contained a morphospecies that accounted for >10% of all individuals, and only 10 plots contained a morphospecies that reached >20%. Given that all 227 hyperdominants reach high local relative abundances (Fig. 3B), these numbers suggest that very few currently unidentified species will eventually qualify as hyperdominant species.

Second, we see strong evidence that taxonomic and identification problems are less severe in hyperdominant species than in other species, in the form of a strong positive correlation between the abundance of a species in the field, the number of specimens in herbaria, and the number of fertile specimens (i.e., specimens with flowers or fruits) collected during field work. Common species are better represented in herbaria than rare species, because individual collectors are more likely to encounter them (29). Common species are also more likely than rare species to be collected fertile during the establishment of tree plots. For example, in 25 ATDN plots established in eastern Ecuador (30), we found that hyperdominant species were more likely than other species to be collected fertile (27.8 versus 17.7%). Botanists trying to identify a hyperdominant species thus

have both a higher likelihood of matching their field specimens with museum specimens and a broader range of morphological features to facilitate identification.

The model we used to estimate population sizes was a loess function, parameterized exclusively with plot location and observed species abundances in plots and no environmental data (problem iv). This is a very different approach from the most commonly used class of species distribution modeling: maximum entropy modeling or Maxent (31, 32). Maxent uses presence-only data fitted to environmental variables of confirmed locations to produce a map of habitat suitability. In a Maxent model, a species known to occur under a given set of environmental conditions is predicted to occur in all environmentally similar areas, even when those areas are outside of the species' known range. Because Amazonian tree species are known to respond strongly to environmental variation, an earlier version of our model included climatic data. That version, however, routinely predicted significant populations of species in regions of the Amazon where a large number of ATDN plots and other plant collection efforts had consistently failed to record those species (i.e., type I errors were common). Modeling with only latitude and longitude as predictive variables is a more conservative option, because it ensures that such errors will be made at a much lower frequency and that species will never be predicted far from confirmed records (Fig. 5). For the same reason, we used a span of 0.2; at higher span values, species ranges extended too far into areas with no known occurrence. Varying span values from 0.2 to 0.5 did not strongly affect population size estimates.

It is not possible to compare estimated population sizes with measured population sizes (problem v), because the latter do not exist for any Amazonian tree species. However, it is possible to compare the population sizes estimated by the loess model with population sizes estimated by using a different method based on the measured extent of Amazonian forest types. The estimated population of *Mauritia flexuosa* is 1.5 billion stems. If we assume that one hectare of monodominant *M. flexuosa* swamp contains 565 *M. flexuosa* trees, then our 1.5-billion-stem estimate suggests that there are <3 million ha of monodominant *M. flexuosa* swamps in the entire basin. This appears reasonable, because the largest block of largely monodominant *M. flexuosa* stands in the basin (the Pastaza Fan) measures ~2.2 million ha. A similar test for white sands and podzol using *E. falcata* and *E. leucantha* (lumped together) was carried out. Together the model estimates that 3.9 billion trees in the greater Amazon belong to these species. If we assume that one hectare of white-sand [podzols and albic arenosols (33)] forest contains on average 150 stems that belong to these species, then the model suggests that there are roughly 26 million ha of white-sand and podzol forest in the greater Amazon. The extent of podzols in the greater Amazon has been estimated as 17 million ha (34). The estimate of podzols and arenosols (fig. S2) is 34 million ha (33).

We know of one study that attempted to estimate populations of trees over a large area in the Amazon Basin based on forest inventories of trees >30 cm dbh (35). The most abundant species in central western Amazonia (blocks: Roraima-Boa Vista, Manaus, and Rio Purus; total forest area 623,139 km²) was *E. coriacea*, with an esti-

mated population of 193 million individuals (this compares to roughly 800 million trees with >10 cm dbh), followed by *Goupia glabra* (93 million individuals, or 370 million trees with >10 cm dbh). Rollet concluded that *E. coriacea* is likely the most common tree species in the Brazilian Amazon. Although our data suggest that two other species have higher total population sizes (*E. precatória* and *Protium altissimum*), a difference caused by our much larger study area (~10×) and lower diameter cutoff (four times as many trees ha⁻¹), our estimate of *E. coriacea* (~5000 million) is of a similar order of magnitude (193 million × 10 × 4 = 7000 million). It is also worth noting that, in the forest inventories used by Rollet, other *Eschweilera* species were pooled more often with *E. coriacea* than in our inventories [see (36, 37) for a discussion on this].

Practical Implications

The finding that Amazonia is dominated by just 227 tree species has important practical implications. It suggests that roughly half of all fruits, flowers, pollen, leaves, and biomass in the world's most diverse forest belong to a very small suite of species, which must therefore account for a large proportion of Amazonian ecosystem services, including water, carbon and nutrient cycling. Our data also suggest that it may be possible to forecast a substantial proportion of the tree community composition and structure of unstudied sites in Amazonia with a purely spatial model. Although no one should underestimate the importance of the >10,000 rare and poorly known tree species in the Amazon (38), an appreciation of how thoroughly common species dominate the basin has the potential to greatly simplify




Fig. 4. Proportions of hyperdominance by region and forest type. (A) Proportions of the trees in each region belonging to species that are regionally dominant, hyperdominant, or neither. **(B)** Proportions of the trees in each forest

type belonging to species that are dominant in that forest type, hyperdominant, or neither. White integers show the number of species in each compartment. IG, igapó; PZ, podzol; SW, swamp; TF, terra firme; VA, várzea.

research in Amazonian biogeochemistry, plant and animal ecology, and vegetation mapping.

Materials and Methods

The ATDN network (39) comprises 1430 tree inventory plots distributed across the Amazon Basin and Guiana Shield, hereafter Amazonia (Fig. 1). Plots were established between 1934 and 2011 by hundreds of different botanists, some working in basinwide or global networks (39–42). Analyses of tree density were performed by using the 1346 plots with trees with ≥ 10 cm dbh that remained after plots with outlying density values (<100 or >1000 individuals/ha), poorly defined areas, or a different diameter cutoff level were removed.

Analyses of composition were performed with a subset of 1170 plots in which all 639,639 free-standing trees with ≥ 10 cm dbh had been iden-

tified with a valid name at the species (86.6%), genus (96.9%), or family (98.9%) level before our study. Most plots (852) measured 1 ha; 253 were smaller, 61 were larger, and 4 were plotless samples (point-centered quarter) for which the sampled area was unknown but the number of trees was equivalent to that typically found in 0.5 to 1 ha. We did not compare specimens or reidentify trees from these plots but resolved major nomenclatural issues (i.e., synonyms and misspellings) in the existing data sets by cross-checking all names with the TROPICOS database (43), via the Taxonomic Name Resolution Service [TNRS (44) (version October 2011)]. We made two adjustments to the names given in TROPICOS (supplementary text). *Rollinia* was merged with *Annona*, because phylogenetic analysis has revealed it to be nested inside that genus (45). Similarly, *Crepidospermum* and *Tetragastris*

are nested in *Protium* (46) and were merged into that genus. For the small proportion of names whose validity could not be determined with those tools, we used *The Plant List* (47). Lianas, bamboos, tree ferns, and tree-sized herbs were excluded from all analyses. Varieties and subspecies were ignored (i.e., all individuals were assigned to the species level). Although some individuals may be misidentified, we assume that this error is within acceptable limits, especially for common species (see discussion above).

The total number of trees ≥ 10 cm dbh in Amazonia was estimated as follows. First, the study area was divided into 567 1° -grid cells (DGCs; Fig. 1). We constructed a loess regression model for tree density (stems ha^{-1}) on the basis of observed tree density in 1195 plots, with latitude, longitude, and their interaction as independent variables. The span was set at 0.5 to yield a relatively




Fig. 5. Distribution maps of three hyperdominant Amazon tree species. Distribution maps estimated by the spatial loess model for three Amazonian hyperdominant species: (A) *E. falcata*, ranked 13th in abundance overall and with an eastern distribution; (B) *Iriarteia deltoidea*, ranked fifth overall and with a western distribution; and (C) *E. coriacea*, ranked third overall and with a pan-Amazonian distribution. Black dots are tree plots where the species has been recorded, and dot size indicates the relative abundance of the species in the plot. Red dots are plots where the species has not been recorded. Shading in DGCs indicates the loess spatial average. For *E. falcata*, the relative abundance in individual plots ranged from 0 to 73.28%, and the loess spatial average in individual grid cells ranged from 0 to 11.27%. Comparable numbers for *I. deltoidea* are 0 to 38.47% and 0 to 12.17% and, for *E. coriacea*, 0 to 21.52% and 0 to 15.01%.

smooth average. The model was used to estimate average tree density in each DGC (D_{DGC} , stems ha^{-1}). The total number of trees in each DGC (N_{DGC}) was then calculated by multiplying D_{DGC} by 1,232,100 ha (the area of a DGC close to the equator—the deviation from this area is just 2.8% at 14°S and 1.1% at 8°N, our latitudinal range). Both empirical (plot data) and interpolated tree densities are illustrated in fig. S4.

The total number of trees belonging to each species in Amazonia was estimated as follows. Abundances of all valid species were converted to relative abundances for each plot: $RA_i = n_i/N$, where n_i = the number of individuals of species i and N = the total number of trees in the plot (including unidentified trees).

For each of the 4962 species with a valid name, we constructed a loess model for RA_i , with latitude, longitude, and their interaction as independent variables and a span of 0.2. We used only spatially independent variables, because test runs including environmental variables commonly led to predictions of species occurrences in well-sampled areas where they had never been recorded in plots. For a similar reason (i.e., to keep predictions spatially conservative), a smaller span was used than in the tree density analysis. Negative predicted abundances were set to 0. The loess model of a species predicted relative abundance in each DGC, yielding a map of its predicted variation in relative abundances across Amazonia. The total population size of each species was calculated by multiplying its relative abundance in each DGC by the total number of trees in that DGC and then summing these products for all DGCs.

To reduce the impact of individual plots and quantify uncertainty in the above procedure, we carried out a bootstrap exercise. This involved randomly drawing 1000 plots from the 1170-plot data set (with replacement) and calculating the population sizes of all species as described above. This was repeated 500 times, and the 500 population estimates per species were used to calculate mean estimated population size and 95% confidence intervals (i.e., mean \pm 1.96 SD).

To estimate range size, we used the same data and methods as (48), standardized with TNRS and updated with specimen records from SpeciesLink (49). Species not found in this database were left out of the range size analysis ($n = 842$). Worldwide species diversity of genera was estimated by counting accepted species in (47). Seed mass and wood density data were obtained from sources described in (36).

Habitat preference was analyzed by means of Indicator Species Analysis, a permutation test that calculates indicator values for each species based on their frequency and relative abundance (50) in the five forest types (igapó, terra firme, swamp, várzea, and white-sand forest).

To analyze regional-level dominance, we divided Amazonia into six regions and created a RAD for each region by summing population sizes in the DGCs they contained. RADs were also constructed for each forest type by sum-

ming the individuals of each species in all plots of a given forest type and calculating the average density of each species in that forest type. The forest-type RADs thus have their basis not in population estimates in DGCs but in the raw abundance data in our plots. A species was considered dominant in a given region or forest type if it appeared in the list of species comprising the upper 50% percentile of the respective RAD.

All analyses were carried out with the R software platform (51). For Indicator Species Analysis, we used the package labdsv. All other permutation tests were custom written.

References and Notes

1. S. P. Hubbell *et al.*, Colloquium paper: How many tree species are there in the Amazon and how many of them will go extinct? *Proc. Natl. Acad. Sci. U.S.A.* **105** (suppl. 1), 11498–11504 (2008). doi: [10.1073/pnas.0801915105](https://doi.org/10.1073/pnas.0801915105); pmid: [18695228](https://pubmed.ncbi.nlm.nih.gov/18695228/)
2. For a general description as well as other supplementary materials, see *Science* Online.
3. G. A. Black, T. H. Dobzhansky, C. Pavan, Some attempts to estimate species diversity and population density of trees in Amazonian forests. *Bot. Gaz.* **111**, 413 (1950). doi: [10.1086/335612](https://doi.org/10.1086/335612)
4. J. M. Pires, T. Dobzhansky, G. A. Black, An estimate of the number of species of trees in an Amazonian forest community. *Bot. Gaz.* **114**, 467 (1953). doi: [10.1086/335790](https://doi.org/10.1086/335790)
5. M. J. G. Hopkins, Modelling the known and unknown plant biodiversity of the Amazon Basin. *J. Biogeogr.* **34**, 1400–1411 (2007). doi: [10.1111/j.1365-2699.2007.01737.x](https://doi.org/10.1111/j.1365-2699.2007.01737.x)
6. IUCN (International Union for Conservation of Nature), *IUCN Red List Categories and Criteria: Version 3.1* (IUCN, Gland, Switzerland, 2001), www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria.
7. M. J. Costello, R. M. May, N. E. Stork, Can we name Earth's species before they go extinct? *Science* **339**, 413–416 (2013). doi: [10.1126/science.1230318](https://doi.org/10.1126/science.1230318); pmid: [23349283](https://pubmed.ncbi.nlm.nih.gov/23349283/)
8. P. Haripersaud, H. ter Steege, J.-J. de Granville, H. Chevillotte, M. Hoff, Species abundance, distribution and diversity in time and space after centuries of collecting in the Guianas. *Taxon* **59**, 592 (2010).
9. D. P. Bebber *et al.*, Herbaria are a major frontier for species discovery. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 22169–22171 (2010). doi: [10.1073/pnas.1011841108](https://doi.org/10.1073/pnas.1011841108); pmid: [21135225](https://pubmed.ncbi.nlm.nih.gov/21135225/)
10. B. J. McGill *et al.*, Species abundance distributions: Moving beyond single prediction theories to integration within an ecological framework. *Ecol. Lett.* **10**, 995–1015 (2007). doi: [10.1111/j.1461-0248.2007.01094.x](https://doi.org/10.1111/j.1461-0248.2007.01094.x); pmid: [17845298](https://pubmed.ncbi.nlm.nih.gov/17845298/)
11. R. J. Warren 2nd, D. K. Skelly, O. J. Schmitz, M. A. Bradford, Universal ecological patterns in college basketball communities. *PLOS ONE* **6**, e17342 (2011). doi: [10.1371/journal.pone.0017342](https://doi.org/10.1371/journal.pone.0017342); pmid: [21408063](https://pubmed.ncbi.nlm.nih.gov/21408063/)
12. N. C. A. Pitman *et al.*, Dominance and distribution of tree species in upper Amazonian terra firme forests. *Ecology* **82**, 2101–2117 (2001). doi: [10.1890/0012-9658\(2001\)082\[2101:DADOTS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[2101:DADOTS]2.0.CO;2)
13. N. C. A. Pitman, M. R. Silman, J. W. Terborgh, Oligarchies in Amazonian tree communities: A ten-year review. *Ecography* **36**, 114–123 (2013). doi: [10.1111/j.1600-0587.2012.00083.x](https://doi.org/10.1111/j.1600-0587.2012.00083.x)
14. F. Domínguez Lozano, M. W. Schwartz, Patterns of rarity and taxonomic group size in plants. *Biol. Conserv.* **126**, 146–154 (2005). doi: [10.1016/j.biocon.2005.04.024](https://doi.org/10.1016/j.biocon.2005.04.024)
15. M. W. Schwartz, D. Simberloff, Taxon size predicts rates of rarity in vascular plants. *Ecol. Lett.* **4**, 464–469 (2001). doi: [10.1046/j.1461-0248.2001.00241.x](https://doi.org/10.1046/j.1461-0248.2001.00241.x)
16. J. E. Richardson, R. T. Pennington, T. D. Pennington, P. M. Hollingsworth, Rapid diversification of a species-rich genus of neotropical rain forest trees. *Science* **293**, 2242–2245 (2001). doi: [10.1126/science.1061421](https://doi.org/10.1126/science.1061421); pmid: [11567135](https://pubmed.ncbi.nlm.nih.gov/11567135/)
17. T. L. Couvreur, F. Forest, W. J. Baker, Origin and global diversification patterns of tropical rain forests: Inferences from a complete genus-level phylogeny of palms. *BMC Biol.* **9**, 44 (2011). doi: [10.1186/1741-7007-9-44](https://doi.org/10.1186/1741-7007-9-44); pmid: [21679405](https://pubmed.ncbi.nlm.nih.gov/21679405/)
18. S. Cavers, C. W. Dick, Phylogeography of Neotropical trees. *J. Biogeogr.* **40**, 615–617 (2013). doi: [10.1111/jbi.12097](https://doi.org/10.1111/jbi.12097)
19. D. H. Janzen, Herbivores and the number of tree species in tropical forests. *Am. Nat.* **104**, 501 (1970). doi: [10.1086/282687](https://doi.org/10.1086/282687)
20. S. A. Mangan *et al.*, Negative plant-soil feedback predicts tree-species relative abundance in a tropical forest. *Nature* **466**, 752–755 (2010). doi: [10.1038/nature09273](https://doi.org/10.1038/nature09273); pmid: [20581819](https://pubmed.ncbi.nlm.nih.gov/20581819/)
21. C. L. Erickson, in *The Handbook of South American Archaeology*, H. Silverman, W. H. Isbell, Eds. (Springer, New York, 2008), pp. 157–183.
22. W. Balée, D. G. Campbell, Evidence for the successional status of liana forest (Xingu river basin, Amazonian Brazil). *Biotropica* **22**, 36 (1990). doi: [10.2307/2388717](https://doi.org/10.2307/2388717)
23. C. Levis *et al.*, Historical human footprint on modern tree species composition in the Purus-Madeira interfluvium, central Amazonia. *PLOS ONE* **7**, e48559 (2012). doi: [10.1371/journal.pone.0048559](https://doi.org/10.1371/journal.pone.0048559); pmid: [23185264](https://pubmed.ncbi.nlm.nih.gov/23185264/)
24. D. A. Posey, Indigenous management of tropical forest ecosystems: The case of Kayapó Indians of the Brazilian Amazon. *Agrofor. Syst.* **3**, 139–158 (1985). doi: [10.1007/BF00122640](https://doi.org/10.1007/BF00122640)
25. C. M. Peters, in *Imperfect Balance: Landscape Transformation in the Pre-Columbian Americas*, D. Lentz, Ed. (Columbia Univ. Press, New York, 2000), pp. 203–223.
26. E. Montoya *et al.*, Forest–savanna–morichal dynamics in relation to fire and human occupation in the southern Gran Sabana (SE Venezuela) during the last millennia. *Quat. Res.* **76**, 335–344 (2011). doi: [10.1016/j.yqres.2011.06.014](https://doi.org/10.1016/j.yqres.2011.06.014)
27. C. Gomez-Navarro, C. Jaramillo, F. Herrera, S. L. Wing, R. Callejas, Palms (Arecaceae) from a Paleocene rainforest of northern Colombia. *Am. J. Bot.* **96**, 1300–1312 (2009). doi: [10.3732/ajb.0800378](https://doi.org/10.3732/ajb.0800378); pmid: [21628279](https://pubmed.ncbi.nlm.nih.gov/21628279/)
28. C. H. McMichael *et al.*, Sparse pre-Columbian human habitation in western Amazonia. *Science* **336**, 1429–1431 (2012). doi: [10.1126/science.1219982](https://doi.org/10.1126/science.1219982); pmid: [22700926](https://pubmed.ncbi.nlm.nih.gov/22700926/)
29. H. ter Steege, P. P. Haripersaud, O. S. Bánki, F. Schieving, A model of botanical collectors' behavior in the field: Never the same species twice. *Am. J. Bot.* **98**, 31–37 (2011). doi: [10.3732/ajb.1000215](https://doi.org/10.3732/ajb.1000215); pmid: [21613082](https://pubmed.ncbi.nlm.nih.gov/21613082/)
30. N. C. A. Pitman, thesis, Duke University, Durham, NC (2000).
31. S. J. Phillips, R. P. Anderson, R. E. Schapire, Maximum entropy modeling of species geographic distributions. *Ecol. Modell.* **190**, 231–259 (2006). doi: [10.1016/j.ecolmodel.2005.03.026](https://doi.org/10.1016/j.ecolmodel.2005.03.026)
32. S. J. Phillips, M. Dudík, Modeling of species distributions with Maxent: New extensions and a comprehensive evaluation. *Ecography* **31**, 161–175 (2008). doi: [10.1111/j.0906-7590.2008.5203.x](https://doi.org/10.1111/j.0906-7590.2008.5203.x)
33. C. A. Quesada *et al.*, Soils of Amazonia with particular reference to the RAINFOR sites. *Biogeosciences* **8**, 1415–1440 (2011). doi: [10.5194/bg-8-1415-2011](https://doi.org/10.5194/bg-8-1415-2011)
34. J. M. Adeney, thesis, Duke University, Durham, NC (2009).
35. B. Rollet, Tree populations in natural tropical forests. A case study: Brazilian Amazonia. *Bois For. Trop.* **236**, 43 (1993).
36. H. ter Steege *et al.*, Continental-scale patterns of canopy tree composition and function across Amazonia. *Nature* **443**, 444–447 (2006). doi: [10.1038/nature05134](https://doi.org/10.1038/nature05134); pmid: [17006512](https://pubmed.ncbi.nlm.nih.gov/17006512/)
37. P. M. Fearnside, Wood density for estimating forest biomass in Brazilian Amazonia. *For. Ecol. Manage.* **90**, 59–87 (1997). doi: [10.1016/S0378-1127\(96\)03840-6](https://doi.org/10.1016/S0378-1127(96)03840-6)
38. D. Moullot *et al.*, Rare species support vulnerable functions in high-diversity ecosystems. *PLOS Biol.* **11**, e1001569 (2013). doi: [10.1371/journal.pbio.1001569](https://doi.org/10.1371/journal.pbio.1001569); pmid: [23723735](https://pubmed.ncbi.nlm.nih.gov/23723735/)

39. H. ter Steege *et al.*, Amazon Tree Diversity Network, <http://web.science.uu.nl/Amazon/ATDN/>.
40. G. Lopez-Gonzalez, S. L. Lewis, M. Burkitt, O. L. Phillips, ForestPlots.net: A web application and research tool to manage and analyse tropical forest plot data. *J. Veg. Sci.* **22**, 610–613 (2011). doi: [10.1111/j.1654-1103.2011.01312.x](https://doi.org/10.1111/j.1654-1103.2011.01312.x)
41. RAINFOR, <http://www.rainfor.org/>.
42. Tropical Ecology Assessment and Monitoring Network, www.teamnetwork.org/.
43. Missouri Botanical Garden, www.tropicos.org.
44. iPlant Collaborative, Taxonomic Name Resolution Service version 2.0, <http://tnrs.iplantcollaborative.org>.
45. P. J. M. Maas, L. Y. T. Westra, H. Rainer, A. Q. Lobão, R. H. J. Erkens, An updated index to genera, species, and infraspecific taxa of Neotropical Annonaceae. *Nord. J. Bot.* **29**, 257–356 (2011). doi: [10.1111/j.1756-1051.2011.01092.x](https://doi.org/10.1111/j.1756-1051.2011.01092.x)
46. P. V. A. Fine, D. C. Daly, G. Villa Muñoz, I. Mesones, K. M. Cameron, The contribution of edaphic heterogeneity to the evolution and diversity of Burseraceae trees in the western Amazon. *Evolution* **59**, 1464–1478 (2005). PMID: [16153032](https://pubmed.ncbi.nlm.nih.gov/16153032/)
47. Royal Botanic Gardens, Missouri Botanical Gardens, The Plant List, www.theplantlist.org/.
48. K. J. Feeley, M. R. Silman, Extinction risks of Amazonian plant species. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 12382–12387 (2009). doi: [10.1073/pnas.0900698106](https://doi.org/10.1073/pnas.0900698106); PMID: [19617552](https://pubmed.ncbi.nlm.nih.gov/19617552/)
49. Centro de Referência em Informação Ambiental, SpeciesLink <http://splink.cria.org.br/>.
50. M. Dufrene, P. Legendre, Species assemblages and indicator species: The need for a flexible asymmetrical approach. *Ecol. Monogr.* **67**, 345 (1997).
51. R Development Core Team, R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, Vienna, Austria, 2012); available online at www.R-project.org.
52. NASA, Visible Earth, <http://visibleearth.nasa.gov/>.

Acknowledgments: This paper is the result of the work of hundreds of different scientists and research institutions in the Amazon over the past 80 years. Without their hard work this analysis would have been impossible. This work was supported by Alberta Mennega Stichting; ALCOA Suriname; Banco de la República; Center for Agricultural Research in Suriname; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Plano Nacional de Pós-Graduação); Conselho Nacional de Desenvolvimento Científico e Tecnológico of Brazil (CNPq) projects Programa de Pesquisas Ecológicas de Longa Duração (PELD) (558069/2009-6), Programa de Apoio a Núcleos de Excelência da Fundação de Amparo à Pesquisa do Estado do Amazonas (PRONEX-FAPEAM) (1600/2006), Áreas Úmidas, and MAUA; PELD (403792/2012-6), PPBio, CENBAM, Universal (479599/2008-4), and Universal 307807-2009-6; Fundação de Amparo À Pesquisa Do Estado Do Amazonas (APEAM) projects DCR/2006, Hidroveg with FAPESP, and PRONEX with CNPq; FAPESP; Colciencias; Duke University; Ecopetrol; FEPIIM 044/2003; the Field Museum; Conservation International/DC (TEAM/INPA Manuas), Gordon and Betty Moore Foundation; Guyana Forestry Commission; Investissement d'Avenir grant of the French Agence Nationale de la Recherche

(ANR) (Centre d'Étude de la Biodiversité Amazonienne ANR-10-LABX-0025); Margaret Mee Amazon Trust; Miquel fons; National Geographic Society (7754-04, 8047-06 to P.M.J.); Netherlands Foundation for the Advancement of Tropical Research WOTRO grants WB85- 335 and W84-581; Primate Conservation Incorporated; Programme Écosystèmes Tropicaux (French Ministry of Ecology and Sustainable Development; Shell Prospecting and Development Peru; Smithsonian Institution's Biological Diversity of the Guiana Shield Program; Stichting het van Eeden-fonds; the Body Shop; the Ministry of the Environment of Ecuador; TROBIT; Tropenbos International; NSF (NSF-0743457 and NSF-0101775 to P.M.J.); USAID; Variety Woods Guyana; WWF-Brazil; WWF-Guianas; Xllème Contrat de Plan Etat Région-Guyane (French Government and European Union); and grants to RAINFOR from the European Union, UK Natural Environment Research Council, the Gordon and Betty Moore Foundation, and U.S. National Geographic Society. O.L.P. is supported by a European Research Council Advanced Grant and a Royal Society Wolfson Research Merit Award. A summary of the data is given in appendix 1. Plot metadata are given in appendix 4.

Supplementary Materials

www.sciencemag.org/content/342/6156/1243092/suppl/DC1
Supplementary Text
Figs. S1 to S12
Tables S1 to S3
Appendices S1 to S4
References (53–67)

11 July 2013; accepted 19 August 2013
10.1126/science.1243092