

NOTE XXXIV.

SYNOPSIS AND ALPHABETICAL LIST
OF THE DESCRIBED SPECIES OF THE COLEOPTEROUS
GENUS HELOTA McL.

BY

C. RITSEMA Cz.

SYNOPSIS OF SPECIES. ¹⁾

- I. Base of elytra coloured as apex.
- A. Elytra with four convex flavous spots.
- a. Pronotum rugose, with raised patches.
- α. Elytra regularly punctate-striate all over.
- ×. Upper surface bright metallic green, flavous elytral spots large.
- *. Apical half of antennae pitchy black; narrowed front-portion of head finely punctured *Vigorsii*. ²⁾
- **.. Antennae entirely pale fulvous; narrowed front-portion of head strongly punctured *scintillans*.
- ××. Upper surface coppery red or dark bronze, flavous elytral spots small.
- †. Upper surface and elytral epipleurae bronze coloured.
- *. Sides of prothorax reddish testaceous. *Faeae*.
- **.. " " " coloured as the disc. *Vandepolli*.
- ††. Upper surface and elytral epipleurae coppery red.

1) For a summary of the principal divisions with reference to the pages, see p. 232.

2) As to the author-names I refer to the alphabetical list of species.

- .. Lateral streaks of pronotum irregularly confluent punctured, leaving some smooth interspaces. — ♂. Apices of elytra conjointly rounded, with a spine at the suture. — ♀. Apices of elytra not or slightly prolonged.
- ». ♂. Anterior tibiae with a small compressed dilatation a little *before* the apex of the inner margin or without it. — ♀. Apices of elytra slightly prolonged, last ventral segment rounded posteriorly.
- »». ♂. Anterior tibiae with a small dilatation, hindlegs of normal length. — ♀. Apices of elytra dehiscent. *Servillei*.
♂. Anterior tibiae without dilatation, hindlegs very slender and elongate. — ♀. Apices of elytra not dehiscent *longipes*.
- »»». ♂. Anterior tibiae with a very conspicuous compressed dilatation *at* the end of the inner margin. — ♀. Apices of elytra conjointly rounded, last ventral segment broadly truncate posteriorly. *Fairmairei*.
- . Lateral streaks of pronotum regularly densely punctured, without smooth interspaces. — ♂. Apices of elytra faintly bidentate near the suture (about as in *Vigorsii*). — ♀. Apices of elytra acutely prolonged, last ventral segment truncate in a bisinuate manner. *Oberthüri*.
- β. Elytra punctured in rows near the suture, the remaining portion irregularly punctured; flavous elytral spots small.
- §. Space between the suture and the line which unites the centres of the flavous

- ⊙. Raised thoracical patches impunctate or nearly so. (Large or moderately large species: $18\frac{1}{2}$ mm. — $12\frac{1}{2}$ mm.).
 - ⊙. ♀. Apices of elytra rather bluntly prolonged; last ventral segment as long as the 3rd and 4th taken together. — ♂. Middle of metasternum with a tomentose spot, sutural spine very distinct. *Gorhami*.
 - ⊙. ♀. Apices of elytra very acutely prolonged; last ventral segment decidedly shorter than the 3rd and 4th taken together. — (♂ unknown). *caudata*.
- ⊙⊙. Raised thoracical patches punctured. (Smaller species: $8\frac{1}{2}$ mm. — $9\frac{1}{4}$ mm.).
 - ×. Sides of prothorax finely crenulate (each with 12—14 crenulations); under surface and legs dark pitchy brown *sinensis*.
 - ××. Sides of prothorax coarsely crenulate (each with 8 or 9 crenulations); middle of sterna and the abdomen pale fulvous; femora pale fulvous with green tips, tibiae and tarsi alternately pitchy and pale fulvous. *thibetana*.
- b. Pronotum more finely and evenly punctured, without raised patches.
 - α. Prosternum entirely fulvous.
 - ×. Legs partly fulvous, partly metallic green.
 - †. Tips of femora and the tibiae metallic green.
 - *. Anterior and posterior elytral spots situated between the same striae, viz. between the 3rd and 7th *Guerinii*.
 - **.

- ,. Anterior spots between the 4th and 6th, the posterior ones between the 3rd and 6th striae *curvipes*.
 ». Anterior spots between the 4th and 7th, the posterior ones between the 3rd and 7th striae *ocellata*.
 »». Anterior spots between the 4th and 7th, the posterior ones between the 3rd and 6th striae.
 ». Outer margin of elytra entire. . . *Dohertyi*.
 »». Apical third of outer margin of elytra serrulate *serratipennis*.
 ††. Tips of femora metallic green; tibiae pitchy, marked with brown-red on the apical half; the spots on the elytra between the 3rd and 6th striae . . . , *rotundata*.
 ××. Legs fulvous, without metallic green, and, especially in the ♂, very slender and elongate. *laevigata*.
 β. Prosternum fulvous in the middle, lateral portions metallic green.
 §. Tibiae entirely metallic green.
 *. Shape narrow (long. $7\frac{1}{2}$ — $8\frac{1}{2}$ mm.; lat. $2\frac{1}{4}$ — $2\frac{1}{2}$ mm.); coppery bronze; anterior elytral spots between the 4th and 6th, the posterior ones between the 3rd and 6th striae. *cereopunc-*
 ***tata*. Shape broader (long. 8 mm.; lat. 3 mm.); bronze green; elytral spots between the 3rd and 6th striae. *brevis*.
 §§. Tibiae metallic green at the base, the rest fulvous.
 ⊖. Anterior and posterior elytral spots between the same striae, viz. between the 3rd and 7th.
 ×. Prothorax quadrate. *Gestroi*.
 ××. » narrowed to the front (trapezoidal).

ALPHABETICAL LIST OF SPECIES.

Helota

- MacLeay. *Annulosa javanica*, 1825, p. 42; *Ed. Lequien*, 1833, p. 151.
- africana* Olliff. *Ann. a. Mag. Nat. Hist.* (5) Angola. XIII, 1884, p. 479. — *Aid Ident. Ins.* t. 153, f. 3.
- Boysii* Rits. *Notes Leyd. Mus.* XI, 1889, India. p. 189.
- brevis* Rits. *Notes Leyd. Mus.* XIII, 1891, Borneo. p. 199.
- caudata* Rits. *Notes Leyd. Mus.* XI, 1889, China. p. 102.
- cereo-punctata* Lewis. *Ent. Mo. Mag.* XVII, 1881, p. 255. — *Aid Ident. Ins.* t. 133, f. 1.
- costata* Rits. *Notes Leyd. Mus.* XI, 1889, Zanzibar. p. 108.
- culta* Olliff. *Cist. Ent.* III, 1883, pp. 55 India bor. and 101; t. 3, f. 2.
- curvipes* Oberth. *Coleopt. Novit.* I, 1883, Himalaya. p. 60. — Rits. *Notes Leyd. Mus.* XI, 1889, p. 105. — id. *Ann. Mus. Civ. Genova*, XXX, 1891, p. 889.
- difficilis* Rits. *Ann. Mus. Civ. Genova*, XXX, Burma. 1891, p. 896.
- Dohertyi* Rits. *Ann. Mus. Civ. Genova*, XXX, " 1891, p. 898.
- dubia* Rits. *Ann. Mus. Civ. Genova*, XXX, " 1891, p. 901.
- Fairmairei* Rits. *Notes Leyd. Mus.* XI, 1889, Sikkim. p. 101.
- Feari* Rits. *Ann. Mus. Civ. Genova*, XXX, Burma. 1891, p. 886.
- fulvitaris* Rits. *Notes Leyd. Mus.* XI, 1889, Darjeeling. p. 107.

Notes from the Leyden Museum. Vol. XIII.

- fulviventris** Kolbe. *Arch. f. Naturgesch.* LII, 1, Korea.
1886, p. 182; t. 11, f. 25. — Rits. *Notes Leyd. Mus.* XI, 1889, p. 104.
- gemmata** Gorh. *Trans. Ent. Soc. London*, 1874, Japan
p. 448 (pars) ¹). — Reitter. *Verhandl. naturf. Ver. Brünn*, XIV, 1876, p. 65;
t. 1, f. 1—5. — Harold. *Abhandl. naturw. Ver. Bremen*, V, 1876, p. 119. —
Rits. *Notes Leyd. Mus.* XI, 1889, p. 104. —
id. *Ann. Mus. Civ. Genova*, XXX, 1891,
p. 888.
- Gestroi** Rits. *Ann. Mus. Civ. Genova*, XXX, Burma.
1891, pp. 893 and 900.
- Gorhami** Olliff. *Cist. Ent.* III, 1883, pp. 53 China.
and 56. — Kolbe. *Arch. f. Naturgesch.*
LII, 1, 1886, p. 181. — Rits. *Notes Leyd.*
Mus. XI, 1889, p. 102.
- Guerinii** Hope. *Coleopt. Manual*, III, 1840, Travancore.
p. 188. — Olliff. *Cist. Ent.* III, 1883,
pp. 54, 56 and 100 (*without synonyms*);
t. 3, f. 1. — Rits. *Notes Leyd. Mus.* XI,
1889, p. 105.
- guineensis** Rits. *Notes Leyd. Mus.* XI, 1889, Assinia.
p. 108.
- immaculata** Rits. *Ann. Mus. Civ. Genova*, XXX, Burma.
1891, p. 895.
- Kolbei** Rits. *Notes Leyd. Mus.* XI, 1889, China.
p. 103. — id. *Ann. Mus. Civ. Genova*,
XXX, 1891, p. 889.
- laevigata** Oberth. *Coleopt. Novit.* I, 1883, Darjeeling.
p. 59. — Olliff. *Cist. Ent.* III, 1884,
p. 100. — *Aid Ident. Ins.* t. 144, f. 1.

1) A very badly drawn figure of this species is published on plate 133 (fig. 2) of Waterhouse's *Aid to the Identification of Insects*. — If the larva described and figured by Sidney Olliff (*Cist. Ent.* III, p. 52; pl. 3, f. 8) belongs to this or to the foregoing species I cannot say, Mr. Lewis having captured both species.

- longipes Rits. *Notes Leyd. Mus.* XI, 1889, Sikkim.
p. 101.
- notata Rits. *Ann. Mus. Civ. Genova*, XXX, Burma.
1891, p. 900.
- Oberthüri Rits. *Notes Leyd. Mus.* XI, 1889, Darjeeling.
p. 100.
- ocellata Rits. *Notes Leyd. Mus.* III, 1881, Java.
p. 79. — id. *l. c.* XI, 1889, p. 105.
- pusilla Oberth. *Coleopt. Novit.* I, 1883, Darjeeling.
p. 60. — Olliff. *Cist. Ent.* III, 1884,
p. 101. — *Aid Ident. Ins. t.* 144, f. 2.
- rotundata Rits. *Ann. Mus. Civ. Genova*, XXX, Burma.
1891, p. 891.
- scintillans Olliff. *Cist. Ent.* III, 1884, p. 99. Java.
- semifulva Rits. *Notes Leyd. Mus.* III, 1881, "
p. 80. — Olliff. *Cist. Ent.* III, 1883,
p. 57. — *Aid Ident. Ins. t.* 153, f. 4.
- serratipennis Rits. *Ann. Mus. Civ. Genova*, Burma.
XXX, 1891, p. 890.
- Servillei Hope. *Coleopt. Manual*, III, 1840, Poonah.
p. 187; t. 3, f. 4. — Rits. *Notes Leyd.*
Mus. XI, 1889, p. 100.
- sinensis Olliff. *Cist. Ent.* III, 1883, p. 54; China.
t. 3, f. 3.
- thibetana Westw. *Ann. a. Mag. Nat. Hist.* Simlah.
VIII, 1842, p. 123 ¹⁾. — Rits. *Stett.*
Ent. Zeit. 1876, p. 19. — Olliff. *Cist.*
Ent. III, 1883, pp. 54 and 56. — Rits.
Notes Leyd. Mus. XI, 1889, p. 110.
- Mellii* Westw. *Cab. Orient. Ent.* 1848,
p. 86; t. 41, f. 8. — Rits. *Notes Leyd.*
Mus. XI, 1889, p. 110.
- Vandepolli Rits. *Notes Leyd. Mus.* XIII, 1891, Borneo.
p. 197.

1) Published in October 1841.

- Vigorsii Mac L. *Annul. Javan.* 1825, p. 43; Java.
 t. 1, f. 9; *Ed. Lequien*, 1833, p. 152;
 t. 5, f. 4. — *Genera des Coléopt. Atlas*,
 t. 131, f. 2.
 var. Rits. *Ann. Mus. Civ. Genova*, XXX, Labuan.
 1891, p. 885. — id. *Notes Leyd. Mus.*
 XIII, 1891, p. 197, note.

SUMMARY OF THE PRINCIPAL DIVISIONS,
 WITH REFERENCE TO THE PAGES.

- p. 223. I. Base of elytra coloured as apex.
 A. Elytra with four convex flavous spots.
 a. Pronotum rugose, with raised patches.
 α. Elytra very regularly punctate striate all over.
 p. 224. β. Elytra punctured in rows near the suture,
 the remaining portion irregularly punctured.
 p. 226. b. Pronotum more finely and evenly punctured,
 without raised patches.
 α. Prosternum entirely fulvous.
 p. 227. β. Prosternum fulvous in the middle, lateral
 portions metallic green.
 p. 228. B. Elytra without flavous spots.
 II. Base of elytra fulvous, apex metallic green
 or black.
 A. Apical portion of elytra metallic green; pro-
 notum entirely fulvous. (Asiatic species).
 B. Apical portion of elytra black; pronotum ful-
 vous at the sides, black along the middle.
 (African species).

Leyden Museum, August 1891.