

## NOTE XIV.

## A NEW MYODITES FROM LIBERIA

(COLEOPTERA: RHIPIPHORIDAE)

DESCRIBED BY

**C. RITSEMA Cz.***Myodites fasciatipennis*, nov. spec.

Length 5 mm. — Black; the plumose antennae pale fulvous, the tip of the flabellae slightly infusate; the legs pale fulvous, the extreme tip of the femora and tibiae and the basal third of the posterior femora infusate; the elytra dark pitchy brown, narrowly margined with pale fulvous and provided with an ill-defined fulvous band (narrowest in the middle) beyond the base; wings hyaline, with a pale fulvous band before the apex, the nervules pale yellowish, the costa and those in the fulvous band darker.

Head and thorax opaque, owing to the very dense punctation, and covered with a short grey pubescence; the vertex conically elevated, the pronotum with three small smooth spots forming a triangle with its top directed towards the front margin of the thorax, the anterior spot impressed and somewhat larger than the basal ones; a trace of a mesial line is present at the base of the pronotum; the scutellum is glossy and impunctate, and provided with a central fovea. The elytra are subshining, strongly punctured, the shoulders glossy and impunctate; their inner margin is very faintly concave, their outer margin strongly and regularly curved, their tip obliquely rounded towards

Notes from the Leyden Museum, Vol. XXXV.

the suture. The abdomen densely punctured, the margins of the segments glossy and impunctate. The legs strongly punctured, especially the posterior pair; the intermediate tarsi are the longest; the basal joint of the posterior tarsi is compressed, broadest about the middle of the length, which is due to the much convex upper side.

Hab. Robertsport (Liberia). — The described specimen has been captured by Mr. Jackson Demery. It is in the Leyden Museum.

Leyden Museum, July 1913.