

NOTE XXVII.

ON TWO NEW SPECIES OF BIRDS FROM
SOUTH CELEBES

BY

J. BÜTTIKOFER.

Amongst the numerous birds, collected by the Dutch botanist Teysmann in the district of Macassar during the year 1878, the Leyden Museum is in possession of a specimen of the genus *Cinnyris*, which I am not able to identify with any known species and which I introduce here under the name of

Cinnyris Teysmanni.

Entire crown, sides of head, hind neck, mantle, back and rump, wing-coverts and outer edge of quills uniform earthy brown with a slight tinge of olive, upper tail-coverts purplish brown, tail-feathers purplish black, the outer webs narrowly edged with metallic green; chin and throat metallic purplish blue, flanked on both sides with a broad metallic green stripe, forming a moustachial streak which is produced down to the sides of the chest; the latter is separated from the breast by a narrow maroon-brown cross-bar which is somewhat paler than in *C. asiaticus*; breast, flanks, abdomen and under tail-coverts black with a purplish gloss, pectoral tufts bright yellow without any trace of orange-red, under wing-coverts and thighs dusky black; bill of the same size as in *C. asiaticus* and *C. zenobia*, but rather less strongly bent, black;

Notes from the Leyden Museum, Vol. XV.

feet black. Wing 5,3 cm.; tail 3,5; tarsus 1,5, bill from front 1,8.

This bird agrees, as to the color of the under-surface, very much with *C. asiaticus*, only is the maroon-brown bar across the chest somewhat paler, the abdomen and under tail-coverts are less strongly glossed with purple and the orange-red feathers in the pectoral tufts are wanting. But the bird cannot be an immature *C. asiaticus*, showing no marks of any metallic gloss on its upper-surface while the lower surface entirely presents the metallic plumage of the fully adult stage. In the transitional stages of *C. asiaticus* the upper-surface, which is much paler than in our Celebean bird, always shows strong marks of metallic blue, especially on rump and lesser wing-coverts, long before the lower-surface has assumed its full metallic plumage. This bird is rather to be placed in Captain Shelley's *Cyrtostomus* or olive-backed Asiatic group, in which the green-backed *C. zenobia* from the Moluccas would be its nearest ally.

The second species, also from the district of Macassar, belongs to the genus *Dicaeum*, and *D. Mackloti* Müll. may be considered to be its nearest ally. From this latter it differs, however, in its smaller size and in the upper-surface having a steel-blue instead of a well-pronounced purplish gloss.

I propose to name this species

Dicaeum splendidum.

Entire head, hind neck and sides of neck, mantle, back and wing-coverts steel-blue, the latter somewhat glossed with purplish, rump and upper tail-coverts scarlet, tail-feathers glossy steel-blue; chin white, throat and fore-neck scarlet, encircled by a dull black stripe which is beginning as a moustachial streak, running down along the sides of the neck and forming a rather broad black band across the chest, continued by a black stripe along

the centre of the breast. Breast, flanks, abdomen, thighs, under tail-coverts and under wing-coverts creamy white, edge of wing dull black; bill and feet black.

D. splendidum: Wing 5,2 cm.; tail 2,7; tarsus 1,8; culmen 0,9.

D. Mackloti: Wing 5,5 cm.; tail 3; tarsus 1,8; culmen 1.

Leyden Museum, March 1893.