

NOTE XXXIX.

A NEW ORIENTAL SPECIES OF THE COLEOPTEROUS
GENUS CHELONARIUM

DESCRIBED BY

C. RITSEMA Cz.

Chelonarium dorsale, n. sp.

Length 7 mm. — Elongate elliptical, much more broadly rounded in front than behind, convex (above and beneath) in the transverse as well as in the longitudinal direction so that the insect is highest just before the middle of its length. Subshining, pitchy brown, darker above than beneath and than the legs and antennae, the tarsi and the short antennal joints pale rusty red; the whole insect covered with a tomentose pubescence and moreover, especially on the upper surface, with long erect hairs; the colour of the entire pubescence is greyish yellow with the exception of an ill-defined transverse spot on the highest portion of the back where the hairs are black, which spot is very conspicuous when the insect is seen sideways.

Head entirely hidden under the prothorax, strongly and densely punctured. — Prothorax distinctly broader than long, slightly narrower than the elytra, very broadly rounded and sharply edged in front, the front-margin slightly upturned, the sides constricted before the posterior angles so as to make the latter divergent; the basal margin bisinuate, the middle-lobe subtruncate with rounded angles; the pronotum equally and rather densely covered with very distinct punctures. Scutellum as long as broad, with strongly curved

sides and rounded tip. — Elytra rather finely and irregularly punctured, having here and there the appearance of being faintly rugose, and showing traces of faint costae; the shoulders are prominent in consequence of a shallow impression along the base and a narrow deeper one along the basal portion of the lateral margins. — The under surface is very densely punctured.

Hab. Java occid. — Captured by Mr. J. D. Pasteur, who presented four specimens to the Leyden Museum.

Together with these four specimens, and from the same sources, I received two others which agree perfectly with the described ones, but they are smaller (measuring 6 mm. in length), the prothorax is more narrowly rounded in front and consequently of a narrower shape, and the pubescence is denser and of a more yellow colour. Most probably these two specimens are males, the four larger ones females.

If the above described species, which is very closely allied to *Ch. unifasciatum* Reitt. from East Sumatra (Notes Leyd. Mus. Vol. VIII, p. 219), did not possess the spot of black pubescence on the middle of the back, I should have meant to have Macleay's *Ch. villosum* before me, the short description of which (see Notes Leyd. Mus. Vol. XI, p. 47) being for the rest pretty well applicable to my insect.

Leyden Museum, October 1891.