BIBLIOGRAPHY OF PACIFIC AND MALESIAN PLANT MAPS OF PHANEROGAMS

Fourth Supplement*

M.M.J. VAN BALGOOY

ACANTHACEAE

Acanthus — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 60.
Partial (2 spp.); Australia; localities indicated; occasional.

A. ilicifolius L. & **A. ebracteatus** Vahl — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 69, fig. 6.
Partial; Australia, New Guinea; localities indicated; partial revision.

Asystasia australasica F.M. Bailey — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 135, fig. 21.
Complete; S. New Guinea + Torres Strait Islands; localities indicated; partial revision.

Brunoniella acaulis (R. Br.) Bremek. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 101, fig. 12.
Complete; Australia, New Guinea; localities indicated; partial revision.

Dicliptera ciliata Decne — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 180, fig. 32.
Partial; Australia, New Guinea; localities indicated; partial revision.

Dipteracanthus bracteatus (R. Br.) Nees — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 94, fig. 10.
Complete; Australia, New Guinea; localities indicated; partial revision.

Graptophyllum spinigerum F. Muell. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 160, fig. 26.
Complete; Australia, New Guinea; localities indicated; partial revision.

Hygrophila angustifolia R. Br. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 122, fig. 18.
Complete; Australia, New Guinea; localities indicated; partial revision.

H. triflora (Roxb.) Fosb. & Sachet — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 122, fig. 18.
Partial; Australia, New Guinea (intr.); localities indicated; partial revision.

*) As in the previous supplement, author names of taxa are added only if they are cited by the author of the paper concerned.

Hypoestes floribunda R. Br. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 204, fig 37.

Partial; Australia, New Guinea; localities indicated; partial revision.

Isoglossa Oersted — B. Hansen, Nord. J. Bot. 5 (1985) 5, fig. 3.

Partial (5 spp.); SE. Asia, Malesia (not Borneo and New Guinea); localities indicated; monograph.

Justicia sect. **Grossa** Hansen — B. Hansen, Nord. J. Bot.7 (1987) 507, fig. 2.

Complete (3 spp.); SE. Asia, Malay Peninsula; localities indicated; monograph.

Lepidagathis cf **royenii** Bremek. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 132, fig. 20.

Complete; Australia (Queensland), New Guinea; localities indicated; partial revision.

Leptostachya wallichii Nees (monotypic genus) — B. Hansen, Nord. J. Bot. 5 (1985) 470, fig. 2.

Complete; India, SE Asia, Sumatra, Java; localities indicated; monograph.

Pseuderanthemum variabile (R. Br.) Radlk. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 151, fig. 24.

Partial; Australia, New Guinea; localities indicated; partial revision.

Thunbergia — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 44, fig. 2.

Partial (3 cult. spp.); Australia, New Guinea; localities indicated; partial revision.

T. arnhemica F. Muell. — R.M. Barker, J. Adel. Bot. Gard. 9 (1986) 44, fig. 2.

Complete; Australia, New Guinea; localities indicated; partial revision.

Thysanostigma Imlay — B. Hansen, Nord. J. Bot. 8 (1988) 229, fig. 2.

Complete (2 spp.); Malay Peninsula; localities indicated; monograph.

ACERACEAE

Acer L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 276.

Complete; World, mainly N. hemisphere, incl. Malesia; delineated; fossil localities indicated; species density; lit. + herb.

Actinidia Lindl. — Liang Chou Fen, Guihaia 3 (1983) 241–245, maps 1–5.

Nearly complete, Malesia not on map (4 sect.); delineated; localities indicated; monograph.

AGAVACEAE see Liliaceae

ALANGIACEAE

Alangium — H.J. Hewson, Fl. Austr. 22 (1984) 17.

Partial; Australia; shaded; partial revision.

W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 35, fig. 11.

Complete; World (present distribution Paleotropics); delineated; fossil localities indicated; occasional.

AMARANTHACEAE

Ptilotus distans (R.Br.) Poir. — G. Benl, Austrobaileya 2 (1984) 19, map. 1.

Partial; N. Australia (incl. islands in Torres Strait); partial revision.

AMARYLLIDACEAE

Campynema lineare (as Liliaceae) — H.J. Hewson, Fl. Austr. 45 (1987) 450, map 440.

Complete; Tasmania; shaded; partial revision.

Campynemanthe Baill. (as Campynemataceae) — P. Goldblatt, Fl. Nouv. Caléd. et Dépend. 16 (1990) 130–133.

Complete; New Caledonia (3 end. spp.); localities indicated; partial revision.

P. Goldblatt, Bull. Nat. Hist. Nat. IV, 8, sect. B, Adansonia 2 (1986) 124, 126, figs. 1 & 2.

Complete; New Caledonia (3 spp.); localities indicated; partial revision.

Curculigo ensifolia (as Liliaceae) — R.J.F. Henderson, Fl. Austr. 45 (1987) 434, maps 206 & 207.

Partial (2 var.); Australia; localities indicated; partial revision.

Hypoxis (as Liliaceae) — R.J.F. Henderson, Fl. Austr. 45 (1987) 433–434, maps 190–205.

Partial (10 spp.); Australia + Tasmania; localities indicated; partial revision.

Molineria capitulata (as Liliaceae) — R.J.F. Henderson, Fl. Austr. 45 (1987) 434, map 208.

Partial; Australia; localities indicated; partial revision.

Proiphys (as Liliaceae) — J.R.H. Telford, Fl. Austr. 45 (1987) 449, maps 431–433.

Partial; Australia (3 spp.); delineated; shaded + localities indicated; partial revision.

ANACARDIACEAE

Pistacia L. - H. Meusel et.al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 274.

Complete; mainly N. hemisphere incl. Malesia (Malay Peninsula, Luzon); delineated; species density.

ANISOPHYLLEACEAE see Rhizophoraceae

ANNONACEAE

Annonaceae — P.J.A. Keßler, Annon. Newsl. 6 (1987) 16, fig. 1.

Complete; pantropical; delineated; occasional.

Anaxagorea St. Hill. — P.J.M. Maas & L.Y.Th. Westra, Bot. Jahrb. 105 (1984) 96, fig. 8.

Complete; Neotropics, Paleotropics (Sri Lanka to Malesia excl. New Guinea); shaded; monograph.

P.J.M. Maas & L.Y.Th.Westra, Bot. Jahrb. 105 (1984) 191–194, figs. 30–38.

Complete; New World (22 spp.); Old World (SE. Asia, Malesia, 7 spp.); localities indicated; monograph.

P.J.A. Keßler, Annon. Newsl. 6 (1987) 21, fig. 6.

Complete; Neotropics, Indo-Malesia (Sri Lanka to Moluccas); delineated; monograph.

Artabotrys — P.J.A. Keßler, Annon. Newsl. 6 (1987) 22, fig.7.

Complete; Paleotropics (E. to New Guinea); delineated; occasional.

Disepalum Hook. f — D.M. Johnson, Brittonia 41 (1989) 365, fig.4, 369, fig. 5.

Complete (9 spp.); SE. Asia, W. Malesia; localities indicated; monograph.

Orophea Blume — P.J.A. Keßler, Annon. Newsl. 6 (1987) 16–25.

Complete (all 27 spp.); S. & SE. Asia, Malesia (India to Moluccas); localities indicated; monograph.

Polyalthia discolor, P. ovalifolia, P. multinervis — S.H. Rogstad, J. Arnold Arbor. 70 (1989) 215, map 3.

Complete; Borneo, New Guinea; localities indicated; monograph.

P. glauca — S.H. Rogstad, J. Arnold Arbor. 70 (1989) 211, map 2.

Complete; Malesia; localities indicated; monograph.

P. hypoleuca — S.H. Rogstad, J. Arnold Arbor. 70 (1989) 229, map 5.

Complete; W. Malesia; localities indicated; monograph.

P. sumatrana — S.H. Rogstad, J. Arnold Arbor. 70 (1989) 221, map 4.

Complete; W. Malesia; localities indicated; monograph.

Uvaria — P.J.A. Keßler, Annon. Newsl. 6 (1987) 23, fig. 8.

Complete; Paleotropics (E. to Fiji); delineated; occasional.

Xylopia — P.J.A. Keßler, Annon. Newsl. 6 (1987) 18, fig. 3.

Complete; pantropical (in Malesia not in Java, Celebes, Lesser Sunda Islands); delineated; occasional.

APOCYNACEAE

Tabernaemontanoideae — L. Allorge, C. R. Soc. Biogéogr. 61 (1985) 138, carte 1.

Complete; World; shaded; species density; occasional.

Tabernaemontana antheonycta Leeuwenb. — A.J.M. Leeuwenberg, A revision of Tabernaemontana 1 (1991) 120, map 18.

Complete; Borneo; localities indicated; partial revision.

T. aurantiaca Gaud. — A.J.M. Leeuwenberg, l.c. 120, map 18.

Complete; E. Malesia, Melanesia (E. to Vanuatu), Palau; localities indicated; partial revision.

T. cerifera Panch. & Séb. — A.J.M. Leeuwenberg, l.c. 135, map 20.

Complete; New Caledonia; localities indicated; partial revision.

T. cordata Merr. — A.J.M. Leeuwenberg, l.c. 131, map 19.

Complete; Philippines (Mindanao); locality indicated; partial revision.

T. corymbosa Roxb. ex Wall. — A.J.M. Leeuwenberg, l.c. 143, map 21.

Complete; SE. Asia, W. Malesia (not Java), Flores; localities indicated; partial revision.

T. crispa Roxb. ex Wall. — A.J.M. Leeuwenberg, l.c. 149, map 22.

Complete; SE. Asia, Malesia (Sumatra to Luzon); localities indicated; partial revision.

T. dichotoma Roxb. ex Wall. — A.J.M. Leeuwenberg, l.c. 131, map 19.

Complete; SE. Asia incl. Sri Lanka, Malay Peninsula; localities indicated; partial revision.

T. macrocarpa Jack — A.J.M. Leeuwenberg, l.c. 161, map 25.

Complete; Malesia (Sumatra to Philippines); localities indicated; partial revision.

T. pandacaqui Lam. — A.J.M. Leeuwenberg, l.c. 179, map 26.

Complete; SE. Asia, Malesia, Australia, Pacific E. to Tahiti; localities indicated; partial revision.

T. pauciflora Blume — A.J.M. Leeuwenberg, l.c. 185, map 27.

Complete; SE. Asia, W. Malesia; localities indicated; partial revision.

T. peduncularis Wall. — A.J.M. Leeuwenberg, l.c. 190, map 28.

Complete; Indo-China, Malay Peninsula; localities indicated; partial revision.

T. remota Leeuwenb. — A.J.M. Leeuwenberg, l.c. 185, map 27.

Complete; Celebes, New Guinea (Rossel I.); localities indicated; partial revision.

T. solomonensis (Markgr.) Leeuwenb. — A.J.M. Leeuwenberg, l.c. 135, map 20.

Complete; Solomons; localities indicated; partial revision.

T. sphaerocarpa Blume — A.J.M. Leeuwenberg, l.c. 161, map 25.

Complete; Java, Lesser Sunda Islands, Celebes, Moluccas; localities indicated; partial revision.

T. thurstonii Horne ex Baker — A.J.M. Leeuwenberg, l.c. 135, map 20.

Complete; Fiji; localities indicated; partial revision.

Beaumontia Wall. — Rudjiman, Agr. Univ. Wageningen Pap. 86-5 (1986) 7–34, maps 1–9.

Complete S. & SE. Asia, W. Malesia (9 spp.); localities indicated; monograph.

Kibatalia G. Don — Rudjiman, Agr. Univ. Wageningen Pap. 86-5 (1986) 42–88, maps 10–24.

Complete (15 spp.); SE. Asia, Malesia; localities indicated; monograph.

Ochrosia minima (Markgr.) Fosb. & Boiteau — P.I. Forster, Austrobaileya 3 (1991) 559, map 1.

Complete; Queensland, S. New Guinea; localities indicated; partial revision.

Vallariopsis lancifolia (Hook. f.) Woodson - Rudjiman, Agr. Univ. Wageningen Pap. 86-5 (1986) 91, map 25.

Complete; Malaya, Sumatra (Siberut and Pagai); localities indicated; monograph.

Vallaris Burm. f. — Rudjiman, Meded. Landbouwhogeschool Wageningen 82-11 (1982) 6, map 1.

Complete (3 spp.); Pakistan to Hainan and W. Malesia; localities indicated; monograph.

Voacanga Thou. – A.J.M. Leeuwenberg, Agr. Univ. Wageningen Pap. 85-3 (1985) 16–65, maps 1–9.

Complete; Africa, Madagascar (7 spp.), Malesia (5 spp.); localities indicated; monograph.

APONOGETONACEAE

Aponogeton L. f. — H.W.E. van Bruggen, Bibl. Bot. 137 (1985) 15–25.

Complete (43 spp.); Old World, incl. Malesia; localities indicated; monograph.

H.W.E.van Bruggen, Aqua Planta Sonderheft 2 (1990) 10.

Complete; Old World, incl. Malesia; shaded; monograph.

AQUIFOLIACEAE

Aquifoliaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 274.

Complete (all genera separately); World; delineated; species density; lit. + herb.

Ilex — L. Pedley, Fl. Australia 22 (1984) 217.

Partial; Australia; shaded; localities indicated; partial revision.

ARACEAE

Lasieae subtr. **Lasiinae** — A. Hay, Ann. Missouri Bot. Gard. 79 (1992) 203 fig. 12.

Complete; World; shaded; monograph.

A. Hay, Ann. Missouri Bot. Gard. 79 (1992) 203, fig. 13.

Partial (distribution of *Lasia*, *Podolasia* and *Cyrtosperma* in Indo-Malesia); delineated; monograph.

Cryptocoryne — N. Jacobsen, Nord. J. Bot. 5 (1985) 48, fig. 6.

Partial; Borneo (14 spp.), 1 also Malaya, 1 from India to New Guinea; localities indicated; partial revision.

N. Jacobsen & J. Bogner, Aqua Planta 3 (1987) 102.

Partial (6 spp.); Malay Peninsula; localities indicated; monograph.

C. fusca de Wit — J. Bastmeijer & B. Bouwmeester, Het Aquarium 60 (10) (1990) 268.

Complete; W. Borneo; localities indicated; occasional.

Cyrtosperma 'Carrii group' — A. Hay, Blumea 33 (1988) 450, fig. 11.

Complete (2 spp.); New Guinea, Aru; localities indicated; partial revision.

C. 'Cuspidispathum group' — A. Hay, Blumea 33 (1988) 446, fig. 8.

Complete (2 spp.); New Guinea; localities indicated; partial revision.

C. 'Merkusii group' — A. Hay, Blumea 33 (1988) 433, 436, figs.2 & 3.

Partial (6 spp.); Malesia, Pacific; localities indicated; partial revision.

Lasia spinosa (L.) Thw. — A. Hay, Blumea 33 (1988) 461, fig. 16.

Complete; India to New Guinea; localities indicated; partial revision.

Podolasia stipitata N.E. Brown — A. Hay, Blumea 33 (1988) 464, fig. 17.

Complete; W. Malesia; localities indicated; partial revision.

ARALIACEAE

Araliaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 303.

Complete; World; delineated; species density; lit. + herb.

Delarbrea Vieill. — P.P. Lowry II, Allertonia 4 (1986) 178–198, figs. 2, 3, 6, 8, 11, 14.

Complete (6 spp.); E. Malesia, Queensland, Melanesia, New Caledonia; localities indicated; monograph.

Macropanax Miq. — C.B. Shang, Bull. Mus. Nat. Hist. Nat, IV, 5 sect. B, Adansonia 1 (1985) 36.

Complete; SE. Asia, Java; localities indicated by numbers; monograph.

ARAUCARIACEAE see Coniferales

ARISTOLOCHIACEAE

Aristolochia engleriana Schmidt — D. Hou, Fl. Males. I, 10 (1984) 107, fig. 18.

Complete; New Guinea; localities indicated; partial revision.

A. foveolata Merr. — D. Hou, Fl. Males. I, 10 (1984) 91, fig. 11.

Complete; W. Malesia incl. Philippines, Taiwan; localities indicated; partial revision.

ASCLEPIADACEAE

Cynanchum — P.I. Forster, Austrobaileya 3 (1991) 465, maps 1–3.

Partial; Australia (11 spp., some also in Malesia); localities indicated; partial revision.

C. brachystelmoides P.I. Forster — P.I. Forster & A. Thongpukdee, Austrobaileya 2 (1988) 457, map 1.

Complete; S. New Guinea, N. Australia; localities indicated; monograph.

Gymnanthera R. Br. — P.I. Forster, Austr. Syst. Bot . 4 (1991) 567, fig. 2.

Partial; Australia (2 spp.); localities indicated; partial revision.

Sarcolobus R. Br. — P.I. Forster, Austrobaileya 3 (1991) 359, maps 1–4.

Partial (11 spp.); Australia, Papuasia; localities indicated; partial revision.

ASTERACEAE see Compositae

BALANOPHORACEAE

Balanophora — H.J. Hewson, Fl. Austr. 22 (1984) 163.

Partial; Australia; shaded; partial revision.

B. fungosa J.R.& G. Forst. — J. Murata, J. Jap. Bot. 63 (1988) 207.

Complete; India to Ryukyu, Australia, and Pacific; localities indicated; partial revision.

BALSAMINACEAE

Hydrocera triflora (L.) W. & A. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 278.

Complete; S. India to Java; delineated; lit. + herb.

Impatiens L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 273.

Complete; World (not in S. America and Australia, incl. Malesia); delineated; species density; lit. + herb.

BATACEAE

Batis argillicola Royen — J.H. Ross, Fl. Austr. 8 (1982) 382, fig. 446.
Partial; Australia; shaded; partial revision.

BEGONIACEAE

Begonia — M.J.S. Sands, Kew Mag. 7 (1990) 58.
Partial; Sabah (6 end. spp.); localities indicated; partial revision.

BIGNONIACEAE

Dolichandrone spathacea — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 61.
Partial; Australia; localities indicated; occasional.

BOMBACACEAE

Bombacaceae — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 24, 25, fig 7.
Complete; pantropical; delineated; fossil range of *Bombacacidites* indicated; occasional.

Camptostemon schultzii — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 61.
Partial; Australia; localities indicated; occasional.

BURMANNIACEAE

Burmannia disticha, B. caelestis, B. championii — Li Heng, Acta Phytotax. Sin. 21 (1983) 122.
Complete; S. & E. Asia, Malesia, N. Australia; localities indicated; partial revision.

BURSERACEAE

Canarium (domesticated species) — D.E. Yen in P.A. Cox & S.A. Banack (eds.), Islands, Plants and Polynesians (1991) 85, fig. 5.
Partial; Malesia, Pacific; localities indicated; occasional.

BUXACEAE

Buxaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 273.
Complete (all genera separately), partial; World incl. W. Malesia; delineated; lit. + herb.

Buxus — M. Bessedik, Pollen et Spores 25 (1983) 474, fig. 3.
Complete; World, incl. Malesia; delineated + shaded (sections indicated); pollen study.

Pachysandra + Sarcococca — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 41, fig. 14.

Complete; World, incl. E. Asia & W. Malesia; delineated; fossil localities (*Erdtmanipollis*) in N. hemisphere indicated; delineated; localities; occasional.

Sarcococca Lindl. — J.R. Sealy, Bot. J. Linn. Soc. 92 (1986) 123, fig. 1.

Complete; S. & E. Asia, W. Malesia; distribution roughly indicated by country; monograph.

CALLITRICHACEAE

Callitriche L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 271.

Complete; World, mainly temperate, also Malesia (Java, New Guinea), New Zealand; delineated.

CAMPANULACEAE

Brighamia — M. Johnson, Kew Mag. 3 (1986) 70.

Complete; Hawaiian Islands; shaded; occasional.

Lobelia L.(and subgenera) — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 450.

Complete; World, incl. Malesia + Pacific; delineated; species density; lit. + herb.

Trematolobelia Zahlbr. — H. St. John, Pacif. Sci. 36 (1982) 504, 505, figs. 8, 9.

Complete (9 spp.); Hawaiian Islands; localities indicated; monograph.

Wahlenbergia Schrad. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 449.

Complete; World, mainly S. hemisphere, incl. Malesia & Pacific; delineated; species density; lit. + herb.

CAMPYNEMATACEAE see Amaryllidaceae

CAPPARACEAE

Cadaba capparoides DC. — H.J. Hewson, Fl. Austr. 8 (1982) 229, fig. 248.

Partial; Australia; shaded; partial revision.

Capparis L. — H.J. Hewson, Fl. Austr. 8 (1982) 219, 229, figs. 228–246.

Partial (17 spp.); Australia; shaded; partial revision.

Cleome L. — H.J. Hewson, Fl. Austr. 8 (1982) 229, 243, figs. 249–260.

Partial (10 spp.); Australia; shaded; partial revision.

Crateva religiosa G. Forster — H.J. Hewson, Fl. Austr. 8 (1982) 219, fig. 228.

Partial; Australia; shaded; partial revision.

CAPRIFOLIACEAE

Lonicera L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 430.

Complete; N. hemisphere, also W. Malesia (S. to Java); delineated; species density; lit. + herb.

L. japonica Thunb. — H. Hara, Ginkgoana 5 (1983) 87, fig. 25.

Partial; E. Asia incl. Japan, Ryukyu, Bonin Is.; delineated + localities partly indicated; partial revision.

Sambucus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 429.

Complete; World, mainly N. hemisphere, incl. Malesia; delineated; species density; lit. + herb.

Viburnum japonicum (Thunb.) Spreng.— H. Hara, Ginkgoana 5 (1983) 232, fig. 36.

Complete (2 var.); Japan, Ryukyu, Bonin (end. var.); localities indicated; partial revision.

CARDIOPTERIDACEAE

Cardiopteris — H.J. Hewson, Fl. Austr. 22 (1984) 217.

Partial; Australia; shaded; partial revision.

CARYOPHYLLACEAE

Scleranthus — J.G. West & P.J. Garnock Jones in B.A. Barlow (ed.), Flora & Fauna of Alpine Australasia (1986) 436, fig. 1.

Complete; Europe, Australasia; delineated; occasional.

S. sect. **Mniarum** — J.G. West & P.J. Garnock Jones in B.A. Barlow (ed.), Flora & Fauna of Alpine Australasia (1986) 440, fig. 3.

Complete; SE. Australia, E. New Guinea, New Zealand; delineated; occasional.

CELASTRACEAE

Hippocrateae — N. Hallé, Bull. Mus. Nat. Hist. Nat. IV, 5, sect. B, Adansonia 1 (1983) 14.

Complete; World, incl. Malesia & W. Pacific (Solomons only, actually widespread in Pacific); delineated; monograph.

Cassine, Celastrus, Euonymus, Lophopetalum, Maytenus, Perrottetia, Pleurostylia, Siphonodon — L.W. Jessup, Fl. Austr. 22 (1984) 163, 173, 191.

Partial; Australia; shaded; partial revision.

Euonymus L.— H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 274.

Complete; World, mainly N. hemisphere, incl. Malesia; delineated; species density; lit. + herb.

Hippocratea, Salacia — L.W. Jessup, Fl. Austr. 22 (1984) 191.

Partial; Australia; shaded; partial revision.

CENTROLEPIDACEAE

Centrolepidaceae/Restionaceae — J. Muller, Ann. Missouri Bot. Gard. 71 (1984) 434, fig. 9.

Partial; Australia; localities indicated; occasional.

CERATOPHYLLACEAE

Ceratophyllum L. — M. Wilmot-Dear, Kew Bull. 40 (1985) 249, map.1.

More or less complete; World incl. Malesia & Pacific; shaded + localities indicated; monograph.

CHENOPODIACEAE

Atriplex — G.A. Parr-Smith in Barker & Greenslade, Evol. of the Flora & Fauna of Arid Austr. (1982) 294–298, fig. 2 & 3.

Partial (several spp.); localities indicated, on grid map; species density; partial revision.

CHLORANTHACEAE

Chloranthaceae (& Austrobaileyaceae) — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 48, 49, fig. 16.

Complete; World; delineated + fossil localities of *Clavatipollenites* indicated; occasional.

Ascarina Forst. — B. Verdcourt, Fl. Males. I, 10 (1986) 140, fig. 6.

More or less complete; Malesia, Pacific (Madagascar not on map); localities indicated; partial revision + Pacific Plant Areas 2: map 64.

W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 48, 49. fig. 16.

Complete; E. Malesia, New Zealand, Pacific, fossil worldwide; delineated; fossil localities (*Clavatipollenites*) indicated; occasional.

A. subsessilis Verdc. — B. Verdcourt, Kew. Bull. 40 (1985) 215, map 1.

Complete; E. New Guinea; localities indicated; partial revision.

Hedyosmum orientale Merr. & Chun — B. Verdcourt, Fl. Males. I, 10 (1986) 144, fig. 8.

Complete; SE. Asia, Malesia (Sumatra, Borneo, Celebes); localities indicated; partial revision.

CHRYSOBALANACEAE

Chrysobalanaceae — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 44, fig. 7.

Complete; World, pantropical; delineated and shaded; monograph.

Atuna — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 48, fig. 14.

Complete; India to Samoa (not Australia and New Caledonia); delineated; monograph.

G.T. Prance, Fl. Males. I, 10 (1989) 666–670, fig. 17–20.

Partial (5 spp.); Malesia, W. Pacific; localities indicated; partial revision.

Chrysobalanus icaco L. — G.T. Prance, Fl. Males. I, 10 (1989) 644, fig. 1.

Partial; Vietnam, Singapore, Fiji (intr.); localities indicated; partial revision.

Hunga Pancher ex Prance — G.T. Prance, Fl. Nouv. Caléd. & Dépend. 12 (1983) 108–123, maps 23–30.

Partial; New Caledonia & Loyalty Islands (8 spp., all end.); localities indicated; partial revision.

G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 47, fig. 11.

Complete; E. New Guinea, New Caledonia; delineated; monograph.

G.T. Prance, Fl. Mal.es I, 10 (1989) 653, fig. 7.

Partial (3 spp.); E. New Guinea; localities indicated; partial revision.

Kostermanthus — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 47, fig. 13.

Complete; W. Malesia, Philippines, Celebes; delineated; monograph.

G.T. Prance, Fl. Males. I, 10 (1989) 677, fig. 24.

Complete (2 spp.); Sumatra, Malaya, Borneo, Philippines, Celebes; localities indicated; partial revision.

Licania — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 46, fig. 8.

Complete; Neotropics, W. Africa, Indo-Malesia (not Moluccas and New Guinea); delineated; monograph.

Maranthes — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 47, fig. 12.

Complete; Old World: Africa, Indo-Australia; New World: Panama; delineated; monograph.

M. corymbosa Blume — G.T. Prance, Fl. Males. I, 10 (1989) 674, fig. 22.

Complete; Malesia, W. Australia, W. Pacific; localities indicated; partial revision.

Parastemon — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 46, fig. 9.

Complete; Indo-Malesia (not Philippines, Java, Lesser Sunda Islands); delineated; monograph.

G.T. Prance, Fl. Males. I, 10 (1989) 650, fig. 5.

Complete (3 spp.); Malesia (not Philippines, Celebes, Java, Lesser Sunda Islands); localities indicated; partial revision.

Parinari — G.T. Prance & F. White, Phil. Trans. Roy. Soc. London, B, Biol. Sc. 320 (1988) 46, fig. 10.

Complete; pantropical, Malesia & Pacific; delineated; monograph.

G.T. Prance, Fl. Males. I, 10 (1989) 656–664, fig. 9–16.

Partial (13 spp.); Malesia, N. Australia, Solomons, SE. Asia; localities indicated; partial revision.

COCHLOSPERMACEAE

Cochlospermum Kunth — A.S. George, Fl. Austr. 8 (1982) 89, fig. 87–90.

Partial (3 spp.); Australia; shaded; partial revision.

COMBRETACEAE

Lumnitzera — D.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 63, 64.

Partial (3 spp.); Australia; localities indicated; occasional.

COMPOSITAE

Asteraceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 450.

Complete; World; percentage of total flora; delineated; lit. + herb.

Anthemideae Cass. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 474.

Complete; World, incl. Malesia & Pacific; delineated; genus density; lit. + herb.

Cardueae Cass. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 504.

Complete; World, mainly N. hemisphere, incl. Luzon; delineated; lit. + herb.

Eupatorieae Cass.— H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 451.

Complete; World; delineated; genus density; lit. + herb.

Heliantheae Cass. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 469.

Complete; World, incl. Malesia and Pacific; delineated; genus density; lit. + herb.

Lactuceae Cass.— H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 525.

Complete; World, incl. Malesia and Pacific; delineated; genus density; lit. + herb.

Senecioneae Cass. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 491.

Complete; World, incl. Malesia & Pacific; delineated; species density; lit. + herb.

Angiantinae — A.A. Anderberg, Opera Bot. 104 (1991) 178, fig. 79.

Complete; Australia, New Zealand; delineated; monograph.

Cassiniinae — A.A. Anderberg, Opera Bot. 104 (1991) 179, fig. 80.

Complete; World, incl. Malesia & W. Pacific; delineated; monograph.

Loricariinae — A.A. Anderberg, Opera Bot. 104 (1991) 176, fig. 77.

Complete; 3 disjunct areas, S. America, Tasmania & New Zealand, China; delineated; monograph.

Argyroxiphium DC. — D.G. Carr, Allertonia 4 (1985) 34–38, fig. 24–29.

Complete (5 spp.); Hawaiian Islands; localities indicated; monograph.

Artemisia L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 484.

Complete; World, incl. Hawaii & Luzon; delineated; species density; lit. + herb.

A. capillaris Thunb. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 489.

Complete; SE. & E. Asia, Luzon; delineated; lit. + herb.

Bidens L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 470.

Complete; World, incl. Malesia & Pacific; delineated; species density; lit. + herb.

B. tripartita L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 470.

Complete; Eurasia, Malesia, Australia, New Zealand; delineated + localities indicated; lit. + herb.

Carpesium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 467.

Complete; Eurasia, incl. Luzon; delineated; species density; lit. + herb.

Celmisia Cass. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 455.

Complete; Australia, New Zealand; delineated; species density; lit. + herb.

C. subg. **Lignosae** sect. **Serratae** — D.R. Given & M. Gray in B.A. Barlow (ed.), Flora & Fauna of Alpine Australasia (1986) 461, fig. 5.

Partial; New Zealand (South Island); localities indicated; occasional.

C. subg. **Pelliculatae** sect. **Petiolatae** — D.R. Given, New Zeal. J. Bot. 22 (1984) 143, fig. 2.

Complete (5 spp.); New Zealand; localities indicated; partial revision.

C. celvisia — D.R. Given & M. Gray in B.A. Barlow (ed), Flora & Fauna of Alpine Australasia (1986) 460, fig. 4.

Partial; New Zealand (South Island); species density; occasional.

C. traversii — D.R. Given, New Zeal. J. Bot. 22 (1984) 143, fig. 2.

Complete; New Zealand; shaded; partial revision.

M.S. McGlone, New Zeal. J. Bot. 23 (1985) 727, fig. 3.

Complete; New Zealand (South Island); shaded; after Given (1984).

C. traversii & C. petriei — D.R. Given & M. Gray in B.A. Barlow, Flora & Fauna of Alpine Australasia (1986) 461, fig. 5.

Complete; New Zealand; localities indicated; occasional.

Cotula L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 484.
Complete; World, incl. New Guinea & New Zealand; delineated; species density; lit. + herb.

Crossostephium chinense — Y. Jotani & H. Ohba, J. Jap. Bot. 61 (1986) 99, fig. 2.
Complete; Taiwan, Ryukyu Islands, Bonin Islands; delineated; occasional.

Darwiniothamnus — J.E. Lawesson & H. Adsersen, Opera Bot.92 (1987) 8, fig. 1.
Complete (3 spp.); Galapagos; localities indicated; monograph.

Dubautia Gaud. — G.D. Carr, Allertonia 4 (1985) 29–39, fig. 16–30.
Complete (21 spp.); Hawaiian Islands; localities indicated; monograph.

Embergeria Boulos — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 536.
Complete; S. Australia, New Zealand; delineated + localities indicated; lit. + herb.

Gnaphalium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 462.
Complete; World; delineated; species density; lit. + herb.

G. luteo-album L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 463.
Complete; Old World, incl. Malesia & Pacific; delineated; lit. + herb.

Helichrysum Mill. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 463.
Complete; Old World, incl. New Guinea & Pacific; delineated; species density; lit. + herb.

Lactuca L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 537.
Complete; World, incl. Malesia; delineated; species density; lit. + herb.

Picris L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 531.
Complete; Eurasia, Africa, (intr.? Australia & New Zealand); delineated; species density; lit. + herb.

Prenanthes L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 540.
Complete; N. hemisphere incl. Sumatra; delineated; species density; lit. + herb.

Robinsonia DC. — P. Pacheco et. al., Am. J. Bot. 72 (1985) 991, fig. 2.
More or less complete; 6 out of 7 spp., Juan Fernandez; localities indicated; occasional (lit. + field observations).

Sigesbeckia L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 469.
Complete; World; delineated + localities indicated; lit. + herb.

Solidago L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 451.
Complete; N. hemisphere incl. Luzon, S. America; delineated; species density; lit. + herb.

Sonchus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 536.
Complete; World incl. Malesia; delineated + localities indicated; species density; lit. + herb.

Tetramolopium Nees — T.K. Lowrey, Allertonia 4 (1986) 217–220, fig. 2–4.
Partial; Hawaiian Islands (11 spp.); localities indicated; partial revision.

Wilkesia — G.D. Carr, Allertonia 4 (1985) 29, fig. 16.
Complete (2 spp.); Hawaiian Islands (Kauai); localities indicated; monograph.

CONIFERALES

Agathis Salisb. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 43, fig. 7.
Complete; Malesia, E. Australia, W. Pacific; delineated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 434–442, fig. 74–86.
Partial (9 spp.); Malesia, Queensland; localities indicated; partial revision.

A. australis — M.S. McGlone, New Zeal. J. Bot. 23 (1985) 742, fig. 8A.
Complete; New Zealand (North Island); shaded; occasional.

A. lanceolata Lindley ex Warb. — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 283, fig. 6.
Complete; New Caledonia; localities indicated; partial revision.

A. montana Laub. — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 282, fig. 5.
Complete; New Caledonia; localities indicated; partial revision.

A. ovata — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 84, fig. 7.
Complete; New Caledonia; localities indicated; partial revision.

Araucaria Juss. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 423, fig. 65.
Complete; New Guinea, E. Australia, W. Pacific, S. America; delineated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 424, 427, fig. 66, 68.
Partial (2 spp.); New Guinea; localities indicated; partial revision.

Dacrycarpus (Endl.) Laub. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 376, fig. 24.
Complete; SE. Asia, Malesia, Pacific; delineated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 377–384, fig. 25–34.
Partial (7 spp.); SE. Asia, Malesia, Pacific; localities indicated; partial revision.

Dacrydium Soland. ex Forst. f. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 362, fig. 10.

Complete; SE. Asia, Malesia, Pacific, S. America; delineated; localities indicated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 364–371, fig. 11–21.

Partial (13 spp.); SE. Asia, Malesia, Fiji; localities indi cated; partial revision.

D. lycopodioides B. & G. — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 280, fig. 4.

Complete; New Caledonia; localities indicated; partial revision.

D. nausoriense — J. Ash, Austr. J. Bot. 34 (1986)199, fig. 1.

Complete; Fiji Islands (Viti Levu); shaded; occasional.

Falcatifolium Laub. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 373, fig. 23.

Partial (4 spp.); Malesia; localities indicated; partial revision.

Libocedrus Endl. — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 273, fig. 1.

Complete (3 spp.); New Caledonia; localities indicated; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 444, fig. 87.

Complete; E. Malesia, New Caledonia, New Zealand, S. America; delineated; species density; partial revision.

L. antrocaledonica — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 275, fig. 2.

Complete; New Caledonia; localities indicated; partial revision.

L. papuana F. Muell. (2 var.) — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 447, fig. 90.

Complete; New Guinea, Moluccas; localities indicated; partial revision.

L. yateensis Guil. — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 286, fig. 8.

Complete; New Caledonia; localities indicated; partial revision.

Nageia Gaertn. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 391, fig. 40.

Complete; World, incl. Malesia, Pacific; delineated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 393, 394, fig.42, 43.

Complete (4 spp.); SE. Asia, Malesia, Pacific; localities indicated; partial revision.

Neocallitropsis pancheri (Carr.) — T. Jaffré et al., Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 276, fig. 3.

Complete; New Caledonia; localities indicated; partial revision.

Phyllocladus — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 357, fig. 5.
Complete; Malesia, Tasmania, New Zealand; delineated; partial revision.

P. hypophyllus Hook.f. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 360, fig. 9.
Complete; Malesia (Borneo, Philippines, Moluccas, New Guinea); localities indicated; partial revision.

P. trichomanoides — M.S. McGlone, New Zeal. J. Bot. 20 (1985) 742, fig. 8 B.
Complete; New Zealand; shaded; occasional.

Pinus L. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 451, 452, fig. 94, 95.
Partial (2 spp.); SE. Asia, Malesia (Sumatra, Philippines); localities indicated; partial revision.

P. subsect. **Sylvestres** — D.I. Axelrod, Ann. Missouri Bot. Gard. 70 (1983) 638, fig. 21.
Complete; N. hemisphere, incl. Luzon; delineated; individual species indicated; occasional.

Podocarpus L'Herit. ex Pers. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 397, fig. 44.
Complete; World; delineated; species density; partial revision.

D.J. de Laubenfels, Fl. Males. I, 10 (1988) 400–418, fig. 45–64.
Partial (30 spp.); SE. Asia, Malesia; localities indicated; partial revision.

Prumnopitys Philippi — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 385, fig. 35.
Complete; Malesia, Pacific, S. America; delineated; species density; partial revision.

P. amara (Blume) Laub. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 387, fig. 37.
Complete; Malesia, Queensland; localities indicated; partial revision.

Sequoia — J. Tivy, Biogeography, ed. 2 (1982) 122, fig. 6, 4.
Complete; N. hemisphere; localities, also fossil, indicated; after Cain 1944 and Good 1964.

Taxus L. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 349, fig. 1.
Complete; World, mainly N. hemisphere, delineated; partial revision.

T. sumatrana (Miq.) Laub. — D.J. de Laubenfels, Fl. Males. I, 10 (1988) 351, fig. 4.
Partial; Malesia (Sumatra, Philippines, Celebes); localities indicated; partial revision.

CONNARACEAE

Cnestis palala — C.C.H. Jongkind in R.H.M.J. Lemmens, The Connaraceae, Agric. Univ. Wageningen Pap. 89-6 (1989) 216, fig. 99.
Complete; SE. Asia, W. Malesia; localities indicated; monograph.

CONVOLVULACEAE

Calystegia soldanella (L.) R. Br. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 359.

Complete; World, bipolar, incl. New Caledonia, New Zealand; delineated; species density.

CORYNOCARPACEAE

Corynocarpus — G.P. Guymer, Fl. Austr. 22 (1984) 217.

Partial; Australia; shaded; partial revision.

CRUCIFERAE

Cardamine L. — H.J. Hewson, Fl. Austr. 8 (1982) 321, fig. 363–370.

Partial (8 spp., some intr.); Australia; shaded; partial revision.

Cheesemannia radicata (Hook.) O. Schulz — H.J. Hewson, Fl. Austr. 8 (1982) 321, fig. 375.

Partial; Australia (Tasmania); shaded; partial revision.

Iti lacustris Garn.-Jones — P.J. Garnock-Jones, New Zeal. J. Bot. 25 (1987) 606, fig. 3.

Complete; New Zealand (South Island, monotypic genus); localities indicated; monograph.

Lepidium L. — H.J. Hewson, Fl. Austr. 8 (1982) 269, 277, fig. 286–328.

Partial (43 spp., several intr.); Australia; shaded; partial revision.

Rorippa Scop. — H.J. Hewson, Fl. Austr. 8 (1982) 333, fig. 377–384.

Partial (8 spp., some intr.); Australia; shaded; partial revision.

CTENOLOPHONACEAE

Ctenolophon — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 20, 21, fig. 6.

Complete; World (present-day distr.: W. Africa, SE. Asia, W. Malesia); delineated; fossil localities indicated; occasional.

C. parvifolius Oliv. — A.M.N. van Hooren & H.P. Nooteboom, Fl. Males. I, 10 (1988) 633, fig. 3.

Complete; Malesia (not: Java, Lesser Sunda Islands, Celebes, Moluccas); localities indicated; partial revision.

CUCURBITACEAE

Benincasa hispida (Thunb.) Cogn. — I.R. Telford, Fl. Austr. 8 (1982) 177, fig. 190.

Partial; Australia; localities indicated; partial revision.

Coccinia grandis (L.) Voigt — I.R. Telford, Fl. Austr. 8 (1982) 177, fig. 195.
Partial; Australia; localities indicated; partial revision.

Diplocyclos palmatus (L.) C. Jeffrey — I.R. Telford, Fl. Austr. 8 (1982) 191, fig. 196.
Partial; Australia; shaded; partial revision.

Gymnostemma Blume subg. **Gymnostemma** — Wu Cheng-Yih & Chen Shu-Kun, Acta Phytotax. Sin. 21 (1983) 356, fig. 1.
Complete; India to Japan and New Guinea; delineated; partial revision.

Luffa Miller — I.R. Telford, Fl. Austr. 8 (1982) 191, fig. 198–199.
Partial (2 spp.); Australia; shaded; species density; partial revision.

Momordica L. — I.R. Telford, Fl. Austr. 8 (1982) 177, fig. 187–189.
Partial (3 spp.); Australia; shaded; partial revision.

Muellerargia timorensis Cogn. — I.R. Telford, Fl. Austr. 8 (1982) 191, fig. 209.
Partial; Australia; localities indicated; partial revision.

Mukia Arn. — I.R. Telford, Fl. Austr. 8 (1982) 191, fig. 202–208.
Partial (7 spp.); Australia; shaded; partial revision.

Neoalsomitra Hutch. — I.R. Telford, Fl. Austr. 8 (1982) 177, fig. 183–186.
Partial (4 spp.); Australia; shaded; partial revision.

Sicyos australis Endl. — I.R. Telford, Fl. Austr. 8 (1982) 205, fig. 214.
Partial; Australia; shaded; partial revision.

Trichosanthes L. — I.R. Telford, Fl. Austr. 8 (1982) 205, fig. 215–220.
Partial (6 spp.); Australia; shaded; partial revision.

Zehneria Endl. — I.R. Telford, Fl. Austr. 8 (1982) 191, fig. 200, 201.
Partial (2 spp.); Australia; shaded; partial revision.

CUNONIACEAE

Acsmithia Hoogl. — R.D. Hoogland, Bull. Mus. Nat. Hist. Nat. IV, 9, sect. B, Adansonia 3 (1987) 394, fig. 1.
Partial (3 spp.); New Caledonia; shaded; partial revision.

CYCADACEAE

Cycadaceae — L. Butt, Austr. Pl. 13 (1984) 19.
Partial (3 genera); Australia; delineated; partial revision.

CYMODOCEACEAE see Potamogetonaceae

CYPERACEAE

Carpha — K.L. Wilson in B.A. Barlow, Flora & Fauna of Alpine Australasia (1986) 476, fig. 3.

Complete; New Guinea, SE. Australia, New Zealand, S. America; delineated; occasional.

Cyperus sect. **Leucocephali** — D.A. Simpson, Kew Bull. 45 (1990) 488, map 1.

Complete (7 spp.); pantropical, incl. Malesia (Luzon, *C. pulchellus*); localities indicated; partial revision.

Oreobolus — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 141, fig. 7.

Complete; circumpacific, incl. Malesia & Pacific; shaded; monograph.

O. ambiguus Kük. & Steen. — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 152, fig. 12.

Complete; Malesia (N. Borneo, Celebes, New Guinea); localities indicated; monograph.

O. impar Edgar — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 181, fig. 31.

Complete; New Zealand (South Island); localities indicated; monograph.

O. kuekentalii Steen. — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 154, fig. 13.

Complete; Malesia (N. Sumatra, Malaya, N. Borneo); localities indicated; monograph.

O. pectinatus Hook.f. — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 184, fig. 32.

More or less complete; New Zealand (in text mentioned for Auckland and Campbell Islands); localities indicated; monograph.

O. pumilio R.Br. — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 170, 171, fig. 23, 24.

Complete (25 spp.); Tasmania, SE. Australia, New Guinea; localities indicated; monograph.

O. strictus Berggreen — O. Seberg, Bot. J. Linn. Soc. 96 (1988) 187, fig. 33.

Complete; New Zealand; localities indicated; monograph.

Rhynchospora — D. Ueno & T. Koyama, Bot. Mag. Tokyo 100 (1987) 78, 79, fig. 3, 4.

Complete; World; shaded; occasional.

D. Ueno & T. Koyama, Bot. Mag. Tokyo 100 (1987) 79, 80, fig. 5.

Complete; World; species density; occasional.

DATISCACEAE

Tetrameles nudiflora R.Br. — T.D. Stanley, Fl. Austr. 8 (1982) 205, fig. 221.

Partial; Australia; locality indicated; partial revision.

DEGENERIACEAE

Degeneria — J.M. Miller, J. Arnold Arbor. 69 (1988) 278, map.1.
Complete; Fiji (2 spp.); localities indicated; partial revision.

DICHAPETALACEAE

Dichapetalum — H.J. Hewson, Fl. Austr. 22 (1984) 219.
Partial; Australia; shaded; partial revision.

DILLENIACEAE

Acrotrema — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 260, fig. 3.
Complete; S. India, Sri Lanka, Sumatra, Malaya; delineated; after Hoogland 1949.

Dillenia — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 260, fig. 3.
Complete; Madagascar to Fiji; delineated; after Hoogland 1949.

Hibbertia — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 260, fig. 3.
Complete; Madagascar to Fiji, incl. E. Malesia; delineated; after Hoogland 1949.

J.M. Veillon, Fl. Nouv. Caléd. et Dépend. 16 (1990) 16–81.
Partial (24 spp., 23 end.); New Caledonia; localities indicated; partial revision.

Tetracera billardieri Mart. — J.M. Veillon, Fl. Nouv. Caléd. et Dépend. 16 (1990) 9.
Complete; New Caledonia; localities indicated; partial revision.

DIPTEROCARPACEAE

Dipterocarpaceae — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 258, fig. 2.
Complete; World (fossil records in parentheses); delineated; after Ashton 1982.

Cotylelobium + Dryobalanops — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 258, fig. 2.
Complete; Sumatra, Malaya, Borneo (fossil in Deccan Peninsula); delineated; after Ashton 1982.

DROSERACEAE

Aldrovanda vesiculosa L. — H.I. Aston, Fl. Austr. 8 (1982) 77, fig. 68.
Partial; Australia; localities indicated; partial revision.

Drosera L. — N.G. Marchant, Fl. Austr. 8 (1982) 25, 35, 47, 59, 77, fig. 7–67.
Partial (54 spp.); Australia; shaded + localities indicated; partial revision.

EHRETIACEAE see Boraginaceae

ELAEOCARPACEAE

Aceratium — M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.1.

Complete; E. Malesia, Bismarcks, Solomons, Vanuatu; delineated; occasional.

Aristotelia — M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.2.

Complete; SE. Australia, New Zealand, S. America; delineated; occasional.

Dubouzetia — M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.1.

Complete; E. Malesia, N. Australia, New Caledonia; delineated; occasional.

Elaeocarpus — M.J.E. Coode, Kew Bull. 39 (1984) 525–571, maps 1–13.

Partial (several spp.); Australia; localities indicated; partial revision.

M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.2.

Complete; Old World, Madagascar to S. Japan, E. Australlia & Pacific; delineated; occasional.

M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 59, map 6.3.

Complete; Madagascar to S. Japan, E. Australia, New Zealand & Hawaii; delineated; species density in outlying areas; occasional.

Sericolea — M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.1.

Complete; New Guinea; delineated; occasional.

Sloanea — M.J.E. Coode in P. Baas et al. (eds.), Plant Diversity of Malesia (1990) 58, map 6.2.

Complete; Madagascar, SE. Asia, Malesia, E. Australia, New Caledonia, Central & S. America; delineated; occasional.

ELAEAGNACEAE

Elaeagnus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 296.

Complete; World, mainly N. hemisphere, incl. Malesia; delineated; species density; lit.+ herb.

ELATINACEAE

Bergia L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 285.

Complete; Africa, Asia, Australia; delineated; lit. + herb.

Elatine L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 285.
Complete; World, incl. Pacific & Malesia; delineated; lit. + herb.

EMPETRACEAE

Empetrum — J. Tivy, Biogeography, ed. 2 (1982) 120, fig. 6.3.
Complete; N. hemisphere, S. America; delineated; after Good 1964.

E. rubrum Vahl ex Willd. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 326.
Complete; S. Atlantic, S. America, Juan Fernandez; delineated; localities indicated; lit. + herb.

EPACRIDACEAE

Epacridaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 331.
Complete; Australia, Malesia, Pacific, S. America; delineated; lit. + herb.

ERICACEAE

Ericaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 331.
Complete; World; delineated; lit. + herb.

Paphia vitiensis Seem. — A.C. Smith, Fl. Vit. Nov. 2 (1981) 717, fig. 185.
Complete; Fiji; localities indicated; partial revision.

Rhododendron L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 327.
Complete; World (not in Africa and S. America), incl. Malesia; delineated; species density; lit. + herb.

R. subg. **Azaleastrum** Planch. sect. **Choniastrum** Franchet — W.R. & M.N. Philipson, Notes Roy. Bot. Gard. Edinb. 44 (1986) 3, fig. 1.
Complete; E.& SE. Asia, Malay Peninsula; localities indicated; partial revision.

Vaccinium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 329.
Complete; World; delineated, isolated locaties indicated; species density; lit. + herb.

EUPHORBIACEAE

Acalypha L. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 112, 113.
Partial (2 spp.); New Caledonia ¶ Loyalties; localities indicated; partial revision.

Aleurites moluccana (L.) Willd. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 74.
Partial; New Caledonia; localities indicated; partial revision.

Alphandia Baill. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 88.

Partial (2 spp.); New Caledonia; localities indicated; partial revision.

Austrobuxus Miq. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 199–221.

Partial (15 spp.); New Caledonia; localities indicated; partial revision.

Baloghia Endl. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 46–72.

Partial (13 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Bocquillonia Baill. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 117.

Complete (end. genus, 14 spp.); New Caledonia; localities indicated; partial revision.

Chamaesyce S.F. Gray (see also under **Euphorbia**) — D.L. Koutnik, Allertonia 4 (1987) 344–374, fig. 3–20.

Partial (14 spp.); Hawaiian Islands; localities indicated; partial revision.

C. clusiifolia, C. herbstii, C. rockii — W.L. Wagner, Occ. Pap. Bish. Mus. 28 (1988) 74.

Complete; Hawaiian Islands (Oahu); localities indicated; partial revision.

Claoxylon insulanum — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 170.

Complete; New Caledonia & Loyalties; localities indicated; partial revision.

Cleidion Blume — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 145–168.

Partial (13 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Cocconerion Baill. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 39–42.

Complete (end. genus, 2 spp.); New Caledonia; localities indicated; partial revision.

Codiaeum Juss. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 98.

Partial (2 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Croton L. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 98.

Partial (2 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Drypetes deplanchei (B.& G.) Megr. — G. McPherson, Fl. Nouv. Caléd. et Dépend. 17 (1991) 13.

Complete; New Caledonia & Loyalties; localities indicated; partial revision.

Euphorbia & Chamaesyce — D.C. Hassal in R.M. Barker & Greenslade, Evol. of Flora & Fauna of Arid Australia (1982) 324–326, fig. 2–9.

Partial (several spp.); Australia; localities indicated; partial revision.

Euphorbia L. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 13–21.

Partial (6 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Excoecaria agallocha L. — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 62.

Partial; Australia; localities indicated; occasional.

G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 33.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

Flueggea Willd. — G.L. Webster, Allertonia 3 (1984) 272, fig. 6.

Complete; World, incl. Malesia & Pacific; delinated; monograph.

F. flexuosa M.A., **F. virosa** (Roxb. ex Willd.) Voigt — G.L. Webster, Allertonia 3 (1984) 281–297, fig. 7–10.

Complete; Old World, incl. Malesia & Pacific; localities indicated; monograph.

Fontainea Heckel — L.W. Jessup & G.P. Guymer, Austrobaileya 2 (1985) 113, map 1.

Complete; E. Australia, New Guinea, New Caledonia, Vanuatu; delineated; partial revision.

F. pancheri (Baill.) Heckel — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 77.

Complete; New Caledonia & Lifu; localities indicated; partial revision.

Glochidion J.R. & G. Forst. — G.McPherson, Fl. Nouv. Caléd. et Dépend. 17 (1991) 22, 24.

Complete (2 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Longetia buxoides Baill. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 192.

Complete (monotypic genus); New Caledonia; localities indicated; partial revision.

Macaranga Petit-Thouars — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 175–185.

Partial (5 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Mallotus repandus (Willd.) M.A. — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 107.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

Myricanthe discolor Airy Shaw — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 74.

Complete (monotypic genus); New Caledonia; localities indicated; partial revision.

Neoguillauminia cleopatra (Baill.) Croizat — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 24.

Complete (monotypic genus); New Caledonia; localities indicated; partial revision.

Omalanthus A. Juss.(= **Homalanthus** A. Juss.) — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 27.

Partial (3 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Phyllanthus — G.L. Webster, Pacif. Sci. 40 (1986) 90, fig. 2.

Partial (4 spp.); Fiji; localities indicated; partial revision.

M. Schmid, Fl. Nouv. Caléd. et Dépend. 17 (1991) 47–318.

Partial (87 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

P. amicorum — G.L. Webster, Pacif. Sci. 40 (1986) 89, fig. 1.

Partial; Tonga; locality indicated; partial revision.

P. myrianthus — G.L. Webster, Pacif. Sci. 40 (1986) 89, fig. 1.

Complete; Vanuatu; localities indicated; partial revision.

Scagea McPherson — G. McPherson & C. Tirèl, Fl. Nouv. Caléd. et Dépend. 14, Euphorbiaceae 1 (1987) 92.

Complete (end. genus, 2 spp.); New Caledonia; localities indicated; partial revision.

FAGACEAE

Fagaceae — T.C. Whitmore, Ann. Missouri Bot. Gard. 69 (1982) 674, fig. 7.

Partial (several genera); SE. Asia, Malesia, Australia, New Zealand; delineated; after Whitmore 1981.

W.N. Ellis, Biogeografie (1988) 73, fig. 37.

Complete; World; delineated; occasional.

Castanopsis & Lithocarpus — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 584, fig. 12.

Complete; W. North America, S. & E. Asia, Malesia; delineated; occasional.

Lithocarpus — R.J. Johns, Common Forest Trees of Papua New Guinea 3 (1983) 99.

Complete; S. & E. Asia, Malesia, W. North America; shaded; partial revision.

Nothofagus — R.J. Johns, Common Forest Trees of Papua New Guinea 3 (1983) 108.

Complete; New Guinea, Australia, New Zealand, S. America; shaded; partial revision.

M.D. Wilcox & N.J. Ledgard, New Zeal. J. Ecol. 6 (1983) 20, fig. 1.

Partial (4 spp.); New Zealand; shaded; lit.

M.S. McGlone, New Zeal. J. Bot. 23 (1985) 734, fig. 5.

Partial; New Zealand; delineated + localities indicated; after Wardle 1984.

N. truncata — M.S. McGlone, New Zeal. J. Bot. 23 (1985) 742, fig. 8D.

Complete; New Zealand, shaded; occasional.

Trigonobalanus verticillata — K.C. Nixon & W.H. Crepet, Amer. J. Bot. 76 (1989) 837, fig. 6.

Complete; Malesia (Sumatra, Malaya, Borneo, Celebes); localities indicated; monograph.

Trisyngyne Baill. (**Nothofagus** subsect. **Bipartitae**) — M.G. Baumann-Bodenheim, Bot. Helv. 93 (1983) 297–307, fig. 23–28.

Partial (5 spp.); New Caledonia; localities indicated; partial revision.

FLACOURTIACEAE

Casearia Jacq. — L.W. Jessup, Fl. Austr. 8 (1982) 89, fig. 81–85.

Partial (5 spp.); Australia; shaded; partial revision.

Flacourtia Comm. ex L'Hérit. — L.W. Jessup, Fl. Austr. 8 (1982) 77, fig. 73, 74.

Partial (2 spp.); Australia; localities indicated; partial revision.

Homalium Jacq. — L.W. Jessup, Fl. Austr. 8 (1982) 89, fig. 78–80.

Partial (3 spp.); Australia; shaded; partial revision.

Ryparosa javanica (Blume) Kurz ex K. & V. — L.W. Jessup, Fl. Austr. 8 (1982) 77, fig. 70.

Partial; Australia; delineated; partial revision.

Scolopia braunii (Klotzsch) Sleum. — L.W. Jessup, Fl. Austr. 8 (1982) 77, fig. 75.

Complete; Australia; shaded; partial revision.

Xylosma G. Forst. — L.W. Jessup, Fl. Austr. 8 (1982) 77, fig. 71, 72.

Partial (2 spp.); Australia; shaded; partial revision.

FRANKENIACEAE

Frankenia L. — A. Barnsley, Fl. Austr. 8 (1982) 135, 145, 157, fig. 136–169.

Partial (47 spp.); Australia; shaded; partial revision.

GENTIANACEAE

Canscora pentanthera C.B. Clarke, **C. andrographioides** C.B.Clarke — A. Ubolcholaket, Thai For. Bull. 14 (1983) 98, 102, fig. 2, 5.

Complete; SE. Asia, incl. Thailand & Malay Peninsula; localities rougly indicated; partial revision.

Centaurium Hill — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 349.

Complete; World, incl. New Caledonia; delineated; species density; lit. + herb.

Crawfurdia trinervis (Blume) Dietr. — A. Ubolcholaket, Thai For. Bull. 14 (1983) 120, fig. 16.

More or less complete; Malay Peninsula, Java; localities rougly indicated; partial revision.

Exacum L. — J. Klackenberg, Opera Bot. 84 (1985) 27, 28; fig. 11, 12.

Complete; Old World, incl. Malesia; delineated; monograph.

E. tetragonum Roxb. — A. Ubolcholaket, Thai For Bull. 14 (1983) 106, fig. 7.

Complete; SE. Asia, incl.Malay Peninsula; localities rougly indicated; partial revision.

E. tetragonum Roxb. — J. Klackenberg, Opera Bot. 84 (1985) 40, fig. 18.

Complete; India & SE. Asia, Malesia, N. Australia; delineated; monograph.

Gentiana L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 350.

Complete; World, incl. Malesia, New Zealand; delineated; species density; lit. + herb.

T.N. Ho & S.W. Liu, Bull. Brit. Mus. Nat. Hist. 20 (1990) 185, fig. 5.

Complete; World; species density; monograph.

G. sect. **Chondrophyllae** Bunge — T.N. Ho & S.W. Liu, Bull. Brit. Mus. Nat. Hist. 20 (1990) 186, fig. 6.

Complete; mainly N. hemisphere, also Malesia, E. Australia; shaded; monograph.

G. sect. **Monopodiae** (H. Sm.) T.N. Ho — T.N. Ho & S.W. Liu, Bull. Brit. Mus. Nat. Hist. 20 (1990) 190, fig. 12.

Complete; E. Asia, W. Malesia; delineated; monograph.

Microrphium pubesens C.B.Clarke — A. Ubolchaket, Thai For. Bull. 14 (1983) 110, fig. 10.

Complete; S. Thailand, Langkawi; localities indicated; partial revision.

Swertia L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 357.

Complete; World (not in Australia, S. America), incl. Malesia; delineated; isolated localities indicated; species density; lit. + herb.

GERANIACEAE

Geraniaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 259.

Complete; World; delineated; isolated localities indicated; lit. + herb.

Geranium solanderi Carol. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 262.

Complete; S. Australia, New Zealand; delineated; lit. + herb.

R.O. Gardner, New Zeal. J. Bot. 22 (1984) 131, fig. 4.

Partial; New Zealand; localities indicated; partial revision.

Pelargonium L'Hérit. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 259.

Complete; Africa, Australia, New Zealand; delineated; lit. + herb.

J.J.A. van der Walt & P.J. Voster, Bothalia 14 (1983) 518–521, fig. 1–4.

Complete; Africa, Australia, Asia minor; shaded; species density (only S. Africa); partial revision.

GESNERIACEAE

Boea Comm. ex Lamk. — B.L. Burtt, Notes Roy. Bot. Gard. Edinb. 41 (1984) 410, fig. 2.

Complete; SE. Asia, Philippines, E. Malesia, Solomons, Queensland; delineated; partial revision.

B. herbacea, B. minutiflora — B.L. Burtt, Notes Roy. Bot. Gard. Edinb. 41 (1984) 410, fig. 2.

Complete; Thailand, Malay Peninsula; localities indicated; partial revision.

Paraboea (C.B.Clarke) Ridley — B.L. Burtt, Notes Roy. Bot. Gard. Edinb. 41 (1984) 410, fig. 2.

Complete; SE. Asia, W. Malesia, Philippines, Celebes (not Java); delineated; partial revision.

P. bettiana — B.L. Burtt, Notes Roy. Bot. Gard. Edinb. 41 (1984) 410, fig. 2.

Complete; Malay Peninsula; locality indicated; partial revision.

Trisepalum C.B.Clarke — B.L. Burtt, Notes Roy. Bot. Gard. Edinb. 41 (1984) 410, fig. 2.

Complete; SE. Asia, N. Malesia; delineated; partial revision.

GOODENIACEAE

Scaevola L. — J.H. Müller, Fl. Nouv. Caléd. et Dépend. 16 (1990) 95–115.

Partial (8 spp., 7 end.); New Caledonia; localities indicated; partial revision.

GRAMINAE

Agrostis — E. Edgar & M.B. Forde, New Zeal. J. Bot. 29 (1991) 144, fig. 1; 146, fig. 3; 148, fig. 4; 150, fig. 5; 151, fig. 6; 155, fig. 7.

Partial (14 spp., 10 native); New Zealand and adjacent islands; partial revision.

Anemanthele lessoniana (Steud.) Veldk. — S.W.L. Jacobs et al., New Zeal. J. Bot. 27 (1989) 581, fig. 6.

Complete (monotypic genus); New Zealand; delineated; partial revision.

Aristida L. sect. **Aristida** — B.K. Simon, Austr. Syst. Bot. 5 (1992) 139, map 1.

Complete; World (incl. S. New Guinea); shaded; partial revision.

A. sect. **Streptachne** — B.K. Simon, Austr. Syst. Bot. 5 (1992) 140, map 5.

Complete; Tropics, incl. S. New Guinea; shaded; partial revision.

Chionochloa Zotov — H.E. Connor, New Zeal. J. Bot. 29 (1991) 229–256, fig. 1–21.

Partial (21 spp.); New Zealand; delineated; species density; partial revision.

Danthonia A.P. DC. — M. Gies, Untersuchungen zur Taxonomie und Chorologie der Gramineengattung Danthonia in Neu Guinea, Thesis, Frankfurt (1991) 65–104, fig. 57–75.

Partial (7 spp.); New Guinea; localities indicated; partial revision.

Dinochloa Buse — D. Ohrnberger & J. Goerrings, The Bamboos of the World, ed. 1 (1984) Dinochloa, p. 7–11, fig. 1–17.

Complete (genus and individual spp.); SE. Asia, Malesia (not New Guinea); delineated; shaded; occasional.

Greslania Balansa — D. Ohrnberger & J. Goerrings, The Bamboos of the World, ed. 1 (1984) Greslania, p. 5–7, map 1–5.

Complete (4 spp.); New Caledonia; delineated; occasional.

Heteropholis C.E. Hubb. — R. de Koning, M.S.M. Sosef & J.F. Veldkamp, Gard. Bull. Sing. 36 (1983) 144, map 1.

Complete (4 spp.); Africa, Madagascar, Sri Lanka, SE. Asia, Malesia, Micronesia; delineated + localities indicated; monograph.

Ichnanthus pallens (Swartz) Munro ex Benth. — M. Stieber, Syst. Bot. 12 (1987) fig. 1, 2, fig. 188, 189.

Complete; pantropical, incl. Malesia; localities indicated; monograph.

Mnesithea laevis (Retz.) Knuth — J.F. Veldkamp, R. de Koning & M.S.M. Sosef, Blumea 31 (1986) 285, map 1.

Complete (3 var.); Asia, Malesia (not Sumatra, Borneo, New Guinea), Micronesia; localities indicated; monograph.

Nasella trichotoma (Nees) Hackel ex Arech. — S.W.L. Jacobs et al., New Zeal. J. Bot. 27 (1985) 580, fig. 5.

Partial; New Zealand (introd.); delineated; partial revision.

Piptatherum miliaceum (L.) Cosson — S.W.L. Jacobs et al., New Zeal. J. Bot. 27 (1985) 581, fig. 6.

Partial; New Zealand (introd.); delineated; partial revision.

Poa L. — E. Edgar, New Zeal. J. Bot. 24 (1986) 478–500, fig. 12–34.

Partial (45 spp., 34 end., 10 naturalized); New Zealand; delineated; partial revision.

Racemobambos Holtt. — D. Ohrnberger & J. Goerrings, The Bamboos of the World, ed. 1 (1984) Racemobambos, p. 7–11, fig. 1–16.

Complete (genus + indiv. spp.); Malesia (not Sumatra, Java, Philippines); delineated + shaded; occasional.

Schizostachyum Nees — D. Ohrnberger & J. Goerrings, The Bamboos of the World, ed. 1 (1984) Schizostachyum, p.13–25, fig. 1–45.

Complete (genus + indiv. spp.); Madagascar, SE. Asia, Malesia, Pacific; delineated; shaded; localities indicated; occasional.

Spartina — T.R. Partridge, New Zeal. J. Bot. 25 (1987) 569, fig. 1.

Partial (3 spp., intr.); New Zealand; localities indicated; partial revision.

Stipa — S.W.L. Jacobs et al., New Zeal. J. Bot. 27 (1989) 573, 575, fig. 2, 3.

Partial (11 spp.); New Zealand; delineated + localities indicated; partial revision.

Thaumastochloa C.E. Hubb. — R. de Koning, M.S.M. Sosef & J.F. Veldkamp, Gard. Bull. Sing. 36 (1983) 144–159, map 1–5.

Complete (7 spp.); Australia, New Guinea, Aru Islands; delineated + localities indicated; monograph.

GUTTIFERAE

Hypericum L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 283.

Complete; World; delineated; lit. + herb.

H. geminiflorum Hemsl. subsp. **geminiflorum** — N.K.B. Robson, Bull. Brit. Mus. Bot. Ser. 12 (1985) 294, map 30.

Complete; Taiwan, Luzon; localities indicated; monograph.

H. henryi H. Lev. & Vaniot subsp. **hancockii** Robson — N.K.B. Robson, Bull. Brit. Mus. Bot. Ser. 12 (1985) 260, map 21.

Complete; SE. Asia, Sumatra; localities indicated; monograph.

H. leschenaultii Choisy — N.K.B. Robson, Bull. Brit. Mus. Bot. Ser. 12 (1985) 242, map 16.

Complete; Sumatra, Java, Lesser Sunda Islands, Celebes; localities indicated; monograph.

HAEMODORACEAE

Haemodoraceae — M.G. Simpson, Ann. Missouri Bot. Gard. 77 (1990) 723.

Complete; World (mainly S. hemisphere, incl. *Haemodorum*: Australia, New Guinea), shaded monograph.

Haemodorum — T.D. Macfarlane, Fl. Austr. 45 (1987) 430–431, map 145–164.

Partial; Australia (20 spp., *H. coccineum* R. Br. also in New Guinea); shaded + localities indicated; species density; monograph.

HERNANDIACEAE

Gyrocarpus americanus — J. Jérémie, Fl. Nouv. Caléd. et Dépend. 15 (1988) 14.

Partial; New Caledonia; localities indicated; partial revision.

Hernandia nymphaeifolia, H. cordigera — J. Jérémie, Fl. Nouv. Caléd. et Dépend. 15 (1988) 6, 11.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

HYDATELLACEAE

Hydatella — D.A. Cooke, Fl. Austr. 45 (1987) 421, maps 1–4.

Partial (4 spp.); Australia; localities indicated; partial revision.

HYDROCHARITACEAE

Blyxa Thou. — C.D.K. Cook & R. Luond, Aquat. Bot. 15 (1983) 47–51, maps 1–5.

Complete (9 spp.); Old World, incl. Malesia; localities indicated; monograph.

Enhalus acoroides (L.f.) Royle — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 25, fig. 16.

Complete (Old World, rough, Philippines, detailed); localities indicated; partial revision.

Halophila beccarii Aschers. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 27, fig. 18.

Complete (SE. Asia, rough, Philippines, Luzon); localities indicated; partial revision.

H. minor (Zoll.) den Hartog — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 29, fig. 20.

Complete (Old World, rough, Philippines, detailed); localities indicated; partial revision.

H. ovalis–H. minor complex — C. McMillan, Aquat. Bot.16 (1983) 340, fig. 1.

Partial; Pacific; localities rougly indicated; occasional.

H. ovalis (R.Br.) Hook.f. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 32, fig. 22.

Complete (Old World, rough, Philippines, detailed); localities indicated; partial revision.

H. spinulosa (R.Br.) Aschers. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 35, fig. 24.

Complete (Malesia, Australia, rough, Philippines, detailed); localities indicated; partial revision.

Thalassia hemprichii (Ehrenb.) Aschers. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 37, fig. 26.

Complete (Old World, rough, Philippines, detailed); localities indicated; partial revision.

HYPERICACEAE see Guttiferae

HYPOXIDACEAE see Amaryllidaceae

ICACINACEAE

Apodytes, Citronella, Gomphandra, Pennantia, Ryticaryum — G.P. Guymer, Fl. Austr. 22 (1984) 217.

Partial; Australia; shaded; partial revision.

ILLICIACEAE

Illiciaceae — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 581, fig. 6.

Complete; N. America, SE. Asia, W. Malesia; delineated; occasional.

JUGLANDACEAE

Juglandaceae — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 584, fig. 11.

Complete (7 genera); America, Asia, Malesia; delineated + shaded; occasional.

Engelhardieae — S.R. Manchester, Monogr. Syst. Bot. Missouri Bot. Gard. 21 (1987) 60, fig. 29.

Complete; mainly N. hemisphere, incl. Malesia; fossil localities indicated; shaded; monograph.

JUNCACEAE

Marsipposperma — K.L. Wilson in B.A. Barlow, Flora & Fauna of Alpine Australasia (1986) 473, fig. 1.

Complete; S. America, New Zealand; delineated; occasional.

Rostkovia — K.L. Wilson in B.A. Barlow, Flora & Fauna of Alpine Australasia (1986) 474, fig. 2.

Complete; New Zealand, S. America, S. Atlantic; delineated; occasional.

LABIATAE

Ajuga bracteosa Benth. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 370.

Complete; Africa, Asia, E. Malesia; delineated; lit. + herb.

Clinopodium L. ser. **Umbrosa** Boriss. (*Clinopodium* is a synonym of *Satureja* L.) — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 382.

Complete; S. & E. Asia, Malesia; delineated; lit. + herb.

Mentha L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 384.

Complete; World, incl. Malesia and New Zealand; delineated; lit. + herb.

Scutellaria L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 371.

Complete; World, incl. Malesia; delineated; lit. + herb.

Teucrium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 369.

Complete; World, incl. Malesia; delineated; species density; lit. + herb.

LAURACEAE

Beilschmiedia Nees — A.E. Wright, New Zeal. J. Bot. 22 1984) 114–122, fig. 3, 6, 9.

Partial (3 spp.); New Zealand; localities indicated; partial revision.

LECYTHIDACEAE

Barringtonia J.R. & G. Forst. — R.J. Henderson, Fl. Austr. 8 (1982) 25, fig. 2–5.

Partial (4 spp.); Australia; shaded; partial revision.

Planchonia careya (F. Muell.) Knuth. — R.J. Henderson, Fl. Austr. 8 (1982) 25, fig. 1.

Partial; Australia; shaded; partial revision.

LEGUMINOSAE

Acacia Mill. (sections) — B.R. Maslin & S.D. Hopper in R.M. Barker & Greenslade, Evol. of Flora & Fauna of Arid Austr. (1982) 309, 310, fig. 5, 6.

Partial; Australia; shaded; partial revision.

A. subg. **Aculeiferum** Vassal — I. Nielsen, Opera Bot. 81 (1985) 9–25, fig. 1–11.

Partial (13 spp.); Malesia, SE. Asia; partly delineated; localities indicated; partial revision.

A. simplex (Sparrm.) Pedl., **A. spirorbis** Labill. — I. Nielsen, Fl. Nouv. Caléd. et Dépend. 12 (1983) 32, 36, maps.3, 4.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

Adenanthera (11 spp., *A. pavonina* excluded) — I. Nielsen & Ph. Guinet, Nord. J. Bot. 12 (1991) 91, fig. 4.

Complete (each sp. separately indicated); SE Asia, Malesia, Queensland; delineated; monograph.

Albizia Durazz. — I. Nielsen, Opera Bot. 81 (1985) 31–46. fig. 12–27.

Partial; Malesia and environs (18 spp.); some delineated; localities indicated; partial revision.

A. guillanii Guillaum. — I. Nielsen, Fl. Nouv. Caléd. et Dépend.12 (1983) 44, map 5.

Complete; New Caledonia; localities indicated; partial revision.

Archidendron F. Muell. — I. Nielsen, T. Baretta-Kuipers & Ph. Guinet, Opera Bot. 76 (1984) 21, fig. 6.

Complete; India to Taiwan throughout Malesia to E. Australia; delineated; species density; monograph.

I. Nielsen, T. Baretta-Kuipers & Ph. Guinet, Opera Bot. 76 (1984) 27, fig. 11.

Complete (distribution of infrageneric categories); Asia, Malesia, Australia; delineated; monograph.

I. Nielsen, T. Baretta-Kuipers & Ph. Guinet, Opera Bot. 76 (1984) 41–102, fig. 14–71.

Partial (several species); Asia, Malesia, Australia; delineated; localities indicated; monograph.

Archidendropsis Nielsen — I. Nielsen, Ph. Guinet, T. Baretta-Kuipers, Bull. Mus. Nat. Hist. Nat. Paris. IV, 5, sect. B, Adansonia 4 (1983) 337, pl. 7.

Complete; E. Australia, New Guinea, Bismarcks, Solomons, New Caledonia, Loyalties; delineated + localities indicated (area of New Caledonian spp. delineated); monograph.

I. Nielsen, Fl. Nouv. Caléd. et Dépend. 12 (1983) 72–102, maps 12–22.

Partial (8 spp., all end.); New Caledonia & Loyalties; localities indicated; partial revision.

Cajanus DC. — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 2, map 1.

Complete; World (original area Old World incl. Malesia); shaded; monograph.

C. cajan (L.) Millsp. — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 67, 72, 77, 78, 89, maps 5–9.

Complete; World, incl. Malesia and Pacific; delineated; and localities indicated; monograph.

C. crassus (Prain ex King) Maesen — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 107, 108, maps 13, 14.

Complete; SE. Asia, W. Malesia, Java, Philippines; localities indicated; monograph.

C. goensis Dalz. — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 121, 122, maps 17, 18.

Complete; SE. Asia, W. Malesia (Java); localities indicated; monograph.

C. platycarpus (Benth.) Maesen — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 163, map 27.

Complete; India, E. Java; localities indicated; monograph.

C. reticulatus (Dryand.) F. Muell. (3 var.) — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 169, map 29.

Complete; Australia, New Guinea; localities indicated; monograph.

C. scarabaeoides (L.) Thouars — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 185–187, maps 31–34.

Complete; Old World, incl. Malesia, Pacific (Micronesia); localities indicated; monograph.

C. volubilis (Blanco) Blanco — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 212, map 37.

Complete; Philippines, Celebes; localities indicated; monograph.

Carmichaelia — A.W. Purdie, New Zeal. J. Bot. 22 (1984) 10, fig. 1; 11, fig.2; 12, fig. 3.

Partial (31 species, distr. of flavonoid components); New Zealand; delineated; occasional.

Cynometra ramiflora — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 62.

Partial; Australia; localities indicated; occasional.

Desmodium microphyllum (Thunb.) DC. — H. Ohashi, J. Jap. Bot. 59 (1984) 43, fig. 1.

Complete; India to Japan and E. Australia, Malesia; delineated; partial revision.

Entada phaseoloides (L.) Merr.— I. Nielsen, Fl. Nouv. Caléd. et Dépend. 12 (1983) 12, map 1.

Partial; New Caledonia; localities indicated; partial revision.

Fordia Hemsley — J.R.M. Buysen, Blumea 33 (1988) 246, fig. 2.

Complete; W. Malesia (not Java), S. China; delineated; species density; monograph.

Mucuna — C.M. Wilmot-Dear, Kew Bull. 45 (1989) 9–29, maps 1–5.

Partial (11 spp.); Pacific (Palau to Hawaii, New Caledonia & Tahiti); localities indicated; partial revision.

Paraserianthes Nielsen — I. Nielsen, Ph. Guinet & T. Baretta-Kuipers, Bull. Mus. Nat. Hist. Nat. Paris. IV, 5, sect. B, Adansonia 4 (1983) 353, pl. 9.

Complete; Sumatra, Java, Lesser Sunda Islands, Moluccas, New Guinea, Bismarcks & Solomons, Australia (SW. & NE.); delineated + localities indicated; monograph.

P. falcataria (L.) Nielsen — I. Nielsen, Ph. Guinet & T. Baretta-Kuipers, Bull. Mus. Nat. Hist. Nat. Paris. IV, 5, sect. B, Adansonia 4 (1983) 359, pl. 10.

Complete (3 subspp.); E. Malesia (incl. Solomons); localities indicated; monograph.

Parkia — H.C.F. Hopkins, Bull. Int. Group Study Mimos. 17 (1989) 68–71, fig. 2–6.

Complete; Pantropical, incl. Malesia & Pacific; delineated; species density; monograph.

Pueraria DC — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 17–99, maps 2–14.

Complete; Indo-Malesia & Pacific, intr. in America, Africa, Australia, Hawaii; shaded; monograph.

L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 17–99, maps 2–14.

Complete (all 16 spp.); Indo-Malesia & Pacific; localities indicated; monograph.

P. lobata (Willd.) Ohwi — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 48–50, maps 6, 7.

Complete (all 16 spp.); SE. Asia, Malesia (not Sumatra, Borneo, Java, Celebes), Pacific (E. to Samoa); localities indicated; monograph.

P. phaseoloides (Roxb.) Benth. — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 74, 77, 82, maps 9, 11.

Complete (3 var.); SE. Asia, Malesia, Pacific; localities indicated; monograph.

P. pulcherrima (Koord.) Merr. — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 91, map 12.

Complete; E. Malesia (Celebes, Philippines to Solomons); localities indicated; monograph.

Schleinitzia insularum (Guillem.) Burk. — I. Nielsen, Fl. Nouv. Caléd. et Dépend. 12 (1983) 14, map 2.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

Serianthes Benth. — I. Nielsen, Fl. Nouv. Caléd. et Dépend. 12 (1983) 50–65, maps 6–11.

Partial (6 spp., all end.); New Caledonia & Loyalties; localities indicated; partial revision.

Spatholobus Hassk. — J.W.A. Ridder-Numan & H. Wiriadinata, Reinwardtia 10 (1985) 155–193, map 5.

Partial (15 out of 29 spp.); SE. Asia, Malesia; localities indicated; monograph.

Strongylodon Vogel — S.F. Huang, Agric. Univ. Wageningen Pap. 90-8 (1991) 17, map 1.

Complete (4 sections); Madagascar to Hawaii; localities indicated; monograph.

S.F. Huang, Agric. Univ. Wageningen Pap. 90-8 (1991) 28–61, maps 1–13.

Complete (all 12 spp.); Indo-Pacific; localities indicated; monograph.

Teyleria Backer — L.J.G. van der Maesen, Agr. Univ. Wageningen Pap. 85-4 (1985) 120, map 16.

Complete (3 spp.); SE. Asia, Malesia,S. Sumatra, Java, Flores, Palawan; localities indicated; monograph.

Wallaceodendron Koord. — I. Nielsen, Ph. Guinet & T. Baretta-Kuipers, Bull. Mus. Nat. Hist. Nat. Paris. IV, 5, sect. B, Adansonia 4 (1983) 349, pl. 8.

Complete; Philippines, Celebes; localities indicated; monograph.

LEMNACEAE

Lemnaceae — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 235, fig. 6.2.
Complete; World, incl. Malesia & Pacific; localities indicated; monograph.

E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 236, fig. 6.3.
Complete; World; shaded; species density; monograph.

Lemna aequinoctialis — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 319, fig. 6.13.
Complete; World, incl. Malesia & Pacific; localities indicated; monograph.

L. disperma — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 272, fig. 6.7.
Complete; Australia, New Zealand; localities indicated; monograph.

L. minor — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 282, fig. 6.8.
Complete; World (not S. America and Malesia), incl. New Zealand; localities indicated; monograph.

L. tenera — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 302, fig. 6.12.
Complete; Burma, Malaya; localities indicated; monograph.

L. trisulca — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 300, fig. 6.11.
Complete; World (not S. America), incl. Malesia; localities indicated; monograph.

Spirodela polyrrhiza — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 254, fig. 6.5.
Complete; World, incl. Malesia; localities indicated; monograph.

S. punctata — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 261, fig. 6.6.
Complete; World, incl. Malesia & Pacific; localities indicated; monograph.

E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 407, fig. 6.35.
Complete; World (year of first record); localities indicated; monograph.

Wolffia augusta — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 370, fig. 6.24.
Complete; SE. Asia, Malaya, Australia; localities indicated; monograph.

W. australiana — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 364, fig. 6.23.
Complete; Australia, New Zealand; localities indicated; monograph.

W. globosa — E. Landolt, Veröff. Geobot. Inst. Rübel 71 (1986) 380, fig. 6.26.
Complete; N. America, Africa, S.& E. Asia incl. Malesia, Hawaii; localities indicated; monograph.

LENTIBULARIACEAE

Utricularia australis R. Br. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 417.

Incomplete; Old World, incl. New Guinea, New Zealand (actually also E. Asia, Java, Sumatra, Philippines); delineated + localities indicated; lit. + herb.

LILIACEAE

Ripogonaceae — J.G. Conran & H.T. Clifford, Nord. J. Bot. 5 (1985) 218, fig. 3.

Complete (as separate family); New Guinea, Australia, New Zealand; shaded; species density; occasional.

Arthropodium — N.H. Brittan, Fl. Austr. 45 (1987) 446, 447, maps 390–393.

Partial (4 spp.); Australia & Tasmania; shaded; localities indicated; partial revision.

Astelia — J.B. Williams, Fl. Austr. 45 (1987) 432, maps 176–179.

Partial (3 spp.); Australia & Tasmania; shaded; partial revision.

A. papuana Skottsb. — R.J. Morley & J.R. Flenley in T.C. Whitmore (ed.), Biogeog. Evol. Mal. Arch. (1987) 52, fig. 5.2.

Partial; New Guinea; localities indicated; herb.

Bulbine — E.M. Watson, Fl. Austr. 45 (1987) 437–438, maps 252–256.

Partial (5 spp.); Australia & Tasmania; shaded; partial revision.

Caesia — R.J.F. Henderson, Fl. Austr. 45 (1987) 441, maps 304, 313.

Partial (8 spp.); Australia & Tasmania; localities indicated; partial revision.

Crinum — H.J. Hewson, Fl. Austr. 45 (1987) 449, fig. 424–428.

Partial (5 spp.); Australia; localities indicated; partial revision.

Dianella — R.J.F. Henderson, Fl. Austr. 45 (1987) 434–437, maps 210, 244.

Partial (15 spp.); Australia & Tasmania; localities indicated; partial revision.

Eustrephus latifolius R. Br. ex Ker-Gawl. — J.G. Conran, Muelleria 6 (1987) 364, fig. 1a.

Complete; E. Australia, New Guinea, Lord Howe, New Caledonia, Loyalties; localities indicated; monograph.

Geitonoplesium cymosum (R. Br.) Cunn. ex R. Br. — J.G. Conran, Muelleria 6 (1987) 364, fig. 1b.

Complete; E. Australia, New Guinea, Lesser Sunda Islands, Bismarcks, Solomons, Lord Howe, Norfolk, New Caledonia, Fiji; localities indicated; monograph.

Herpolirion - H.J. Hewson, Fl. Austr. 45 (1987) 438, map 257.

Partial; Australia & Tasmania; shaded; partial revision.

Heterosmilax Kunth — T. Koyama, Brittonia 36 (1984) 185, fig. 1.
Complete; SE. Asia, W. Malesia; delineated; monograph.

H. borneensis A.DC. — T. Koyama, Brittonia 36 (1984) 185.
Complete; SE. Asia, W. Malesia; localities indicated; monograph.

Pleomele — H. St.John, Pacif. Sci. 39 (1985) 189, fig. 8.
Partial (9 end. spp.); Hawaiian Islands; localities indicated; partial revision.

Schelhammera — H.T. Clifford & J.G. Conran, Fl. Austr. 45 (1987) 452, maps 471, 472.
Partial (2 spp.); Australia; shaded; partial revision.

Thysanotus — N.H. Brittan, Fl. Austr. 45 (1987) 443–446, maps 338–387.
Partial (49 spp.); Australia; shaded; localities indicated; partial revision.

LINACEAE

Hugonia L. (**Durandea** Planch.) — A.M.N. van Hooren & H.P. Nooteboom, Blumea 29 (1984) 553, fig. 1.
Complete; Old World, W. Africa to Fiji, incl. Malesia; delineated; partial revision.

A.M.N. van Hooren & H.P. Nooteboom, Fl. Males. I, 10 (1988) 612, fig. 2.
Partial (2 spp.); Malesia, Pacific, E. Australia; localities indicated; partial revision.

Indorouchera Hall.f. — A.M.N. van Hooren & H.P. Nooteboom, Fl. Males. I, 10 (1988) 618, 619, fig. 9, 10.
Partial (2 spp.); SE. Asia, W. Malesia; localities indicated; partial revision.

Philbornea magnifolia (Stapf) Hall.f. — A.M.N. van Hooren & H.P. Nooteboom, Fl. Males. I, 10 (1988) 615, fig. 5.
Complete; Malesia (Sumatra, Borneo, Palawan); localities indicated; partial revision.

LOBELIACEAE see Campanulaceae

LOGANIACEAE

Strychnos — L.T.C. Khanh, Feddes Rep. 97 (1986) 718, map 1.
Complete; World; delineated; after Pacif. Plant Areas 2 (1966).

LORANTHACEAE

Loranthaceae — B.A. Barlow in H.C. Weber & W. Forstreuter (eds.), Parasitic Flow. Pl. (1987) 809, fig. 3.
Complete; World, relictual genera indicated; delineated; occasional.

B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 278, fig. 22.3.

Complete; World, relictual genera (Australia, S. America) shown in black; delineated; monograph.

Amyema, Amylotheca, Cecarria, Dactyliophora, Decaisnina, Dendrophthoe — B.A. Barlow, Fl. Austr. 22 (1984) 83, 97, 135.

Partial; Australia; localities indicated; partial revision.

Amyema — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig. 22.5.

Partial; Malesia, Australia; delineated; monograph.

A. acuta (Tieghem) Danser, **A. apoensis** (Elmer) Barlow, **A. arthrocaulis** Barlow, **A. beccarii** (Tieghem) Danser — B.A. Barlow, Blumea 36 (1992) 307, fig. 1.

Complete; Malesia; localities indicated, monograph.

A. artensis (Montr.) Danser — B.A. Barlow, Blumea 36 (1992) 310, fig. 2.

Complete; New Guinea, New Caledonia, Vanuatu, Carolines, Samoa; localities indicated; monograph.

A. benguetensis (Merr.) Danser, **A. brassii** Barlow, **A. canaliculata** Barlow, **A. cauliflora** (Merr.) Barlow, **A. celebica** (Tieghem) Danser — B.A. Barlow, Blumea 36 (1992) 315, fig. 3.

Complete; Malesia; localities indicated; monograph.

A. caudiciflora (Laut.) Danser — B.A. Barlow, Blumea 36 (1992) 336, fig. 8.

Complete; Malesia; localities indicated; monograph.

A. cercidioides (Krause) Danser, **A. conspicua** (F.M. Bailey) Danser subsp. **fulvicalyx** Barlow + subsp. **conspicua** + subsp. **obscurinervis** Barlow — B.A. Barlow, Blumea 36 (1992) 322, fig. 4.

Complete; Australia, New Guinea; localities indicated; monograph.

A. corniculata Danser, **A. cuernosensis** (Elmer) Barlow, **A. curranii** (Merr.) Danser, **A. dilatipes** Barlow, **A. dolichopoda** Barlow — B.A. Barlow, Blumea 36 (1992) 324, fig. 5.

Complete; Australia, Malesia; localities indicated; monograph.

A. eburna (Barlow) Barlow, **A. edanoi** (Merr.) Barlow, **A. enneantha** Barlow, **A. finisterrae** (Warb.) Danser — B.A. Barlow, Blumea 36 (1992) 327, fig. 6.

Complete; Australia, Malesia; localities indicated; monograph.

A. fasciculata (Bl.) Danser, **A. friesiana** (Schumann) Danser — B.A. Barlow, Blumea 36 (1992) 331, fig. 7.

Complete; Malesia; localities indicated; monograph.

A. gravis Danser, **A. haenkeana** (Schultes) Danser, **A. halconensis** (Merr.) Danser, **A. hastifolia** (Ridley) Danser, **A. hexameres** (Danser) Danser, **A. hexantha** (Merr.) Barlow — B.A. Barlow, Blumea 36 (1992) 336, fig. 8.

Complete; Malesia; localities indicated; monograph.

A. incarnatiflora (Elmer) Danser, **A. irrubescens** Barlow, **A. kebarensis** Barlow, **A. longipes** (Danser) Barlow, **A. luzonensis** (Schultes) Danser — B.A. Barlow, Blumea 36 (1992) 342, fig. 9.

Complete; Malesia; localities indicated; monograph.

A. mackayensis (Blakely) Danser, **A. novaebritanniae** (Schumann) Danser — B.A. Barlow, Blumea 36 (1992) 346, fig. 10.

Complete; Australia, New Guinea; localities indicated; monograph.

A. miraculosa (Miq.) Tieghem subsp **boormanii** (Blakely) Barlow + subsp **latifolia** Barlow + subsp **miraculosa**, **A. pachypus** (Burk.) Danser, **A. polillensis** (Robinson) Danser, **A. polytrias** Danser, **A. pyriformis** Barlow, **A. queenslandica** (Blakely) Danser — — B.A. Barlow, Blumea 36 (1992) 351, fig. 11.

Complete; Australia, Malesia; localities indicated; monograph.

A. plicatula (Krause) Danser — B.A. Barlow, Blumea 36 (1992) 354, fig. 12.

Complete; Australia, New Guinea; localities indicated; monograph.

A. rhytidoderma Barlow, **A. rigidiflora** (Krause) Danser, **A. scheffleroides** Barlow, **A. seemeniana** (Schum.) Danser, **A. seriata** (Merr.) Danser — B.A. Barlow, Blumea 36 (1992) 360, fig. 13.

Complete; Malesia; localities indicated; monograph.

A. scandens (Tieghem) Danser — B.A. Barlow, Blumea 36 (1992) 363, fig. 14.

Complete; New Guinea, New Caledonia; localities indicated; monograph.

A. squarrosa (Krause) Danser, **A. strongylophylla** (Laut.) Danser, **A. tetraflora** (Barlow) Barlow, **A. timorana** Danser, **A. triantha** (Korth.) Tieghem, **A. urdanetensis** (Elmer) Danser, **A. verticillata** (Merr.) Danser — B.A. Barlow, Blumea 36 (1992) 367, fig. 15.

Complete; Malesia; localities indicated; monograph.

A. tetrapetala (Danser) Barlow, **A. tristis** (Zoll.) Tieghem, **A. umbellata** Danser, **A. vernicosa** Barlow, **A. villiflora** (Domin) Barlow, **A. wenzelii** (Merr.) Danser, **A. wichmannii** (Krause) Danser — B.A. Barlow, Blumea 36 (1992) 376, fig. 16.

Complete; Australia, Malesia; localities indicated; monograph.

Amylotheca — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 289, fig. 22.6.

Partial; New Guinea, New Caledonia, Australia (incomplete); delineated; monograph.

Dactyliophora — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig. 22.5.

Complete; New Guinea, Solomons, N. Queensland; delineated; monograph.

Decaisnina — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 284, fig. 22.6.

More or less complete; Malesia, N. Australia, distribution in Pacific incomplete; delineated; monograph.

Dendrophthoe — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 285, fig. 22.7.

More or less complete; E.& N. Australia, Malesia, distribution in Asia incomplete; delineated; monograph.

Distrianthes — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig. 22.5.

Complete; New Guinea; delineated; monograph.

Ginalloa — B.A. Barlow in H.C. Weber & W. Forstreuter (eds.), Parasitic Flow. Pl. (1987) 814, fig. 6.

Complete; Indo-Malesia; delineated; occasional.

B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 289, fig. 22.8.

Complete; E. Asia, Malesia; delineated; monograph.

Korthalsella Tieghem — B.A. Barlow, Brunonia 6 (1983) 42, 43, fig. 23–25.

Partial (8 spp.); Australia incl. Lord Howe and Norfolk Islands; localities indicated; monograph.

B.A. Barlow in H.C. Weber & W. Forstreuter (eds.), Parasitic Flow. Pl. (1987) 815, fig. 7.

Complete; Old World, incl. Malesia, Pacific; delineated; occasional.

Korthalsella, Notothixos, Viscum — B.A. Barlow, Fl. Austr. 22 (1984) 135, 145.

Partial; Australia; shaded; partial revision.

Lepeostegeres — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 284, fig. 22.6.

Complete; Malesia; delineated ; monograph.

Macrosolen — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 284, fig. 22.6.

More or less complete; SE. Asia; Malesia, distribution in Asia incomplete; delineated; monograph.

Notothixos Oliv. — B.A. Barlow, Brunonia 6 (1983) 4, fig. 1.

Complete; Ceylon through Malesia (not Sumatra) to St. Cruz and E. Australia; delineated; monograph.

B.A. Barlow, Brunonia 6 (1983) 10–21, fig. 6–11.

Complete (8 spp.); Ceylon, Malesia, E. Australia, Solomons, St. Cruz; localities indicated; monograph.

B.A. Barlow in H.C. Weber & W. Forstreuter (eds.), Parasitic Flow. Pl. (1987) 814, fig. 6.

Complete; Ceylon, Malesia, E. Australia, Solomons; delineated; occasional.

B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 289, fig. 22.8.

Partial; Malesia, Australia, incomplete for Asia and Australia; delineated; monograph.

Papuanthes — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig.22.5.

Complete; New Guinea; delineated; monograph.

Scurrula — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 285, fig. 22.7.

Partial; Malesia, eastern limit Celebes, Lesser Sunda Islands; delineated; monograph.

B.A. Barlow, Blumea 36 (1991) 69–81, fig. 2–8.

Complete (8 spp.); SE. Asia, Malesia, W. to Moluccas; delineated (outside Malesia), localities indicated (in Malesia); monograph.

Sogerianthe — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig. 22.5.

Complete; New Guinea, Solomons; delineated; monograph.

Taxillus chinensis (DC.) Danser — B.A. Barlow, Blumea 36 (1991) 83, fig. 9.

Complete; SE. Asia, W. Malesia;delineated + localities indicated; monograph.

Tetradyas — B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 282, fig. 22.5.

Complete; New Guinea; delineated; monograph.

Viscum — B.A. Barlow in H.C. Weber & W. Forstreuter (eds.), Parasitic Flow. Pl. (1987) 816, fig. 8.

Complete; Old World, incl. Malesia; delineated; species density; occasional.

B.A. Barlow in P. Baas, C. Kalkman & R. Geesink (eds.), Plant Diversity of Malesia (1990) 289, fig. 22.8.

Incomplete; Asia, Malesia, Australia (incomplete for Asia and Australia); delineated; monograph.

LYTHRACEAE

Lythrum hyssopifolia — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 297.

Complete; World (bipolar), incl. New Zealand; delineated; lit. + herb.

MAGNOLIACEAE

Magnoliaceae — T.C. Whitmore, Ann. Missouri Bot. Gard. 69 (1982) 670, fig. 2.

Partial (several genera); SE. Asia, Malesia; delineated; after Whitmore 1981.

Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 582, fig. 9.

Complete (*Magnolia* + *Liriodendron* separately indicated); America, S.& E. Asia, Malesia; delineated; occasional.

Law Yuh — Wu, Acta Phytotax. Sin. 22 (1984) facing p. 9, fig. 7.

Complete (several genera); S.& E. Asia, Malesia, America; delineated; partial revision.

Magnolia — J. Tivy, Biography, ed. 2 (1982) 120, fig. 6.3.

Complete; SE. Asia, Malesia, America; shaded; after Good 1964.

W.N. Ellis, Biographie (1988) 65, fig. 29.

Complete; World; delineated + fossil localities indicated; occasional.

Michelia montana Bl. — H.P. Nooteboom, Blumea 31 (1985) 117, fig. 11.

Complete; W. Malesia; localities indicated; partial revision.

H.P. Nooteboom, Fl. Males. I, 10 (1988) 604, fig. 14.

Complete; Malesia; localities indicated; partial revision.

MALPIGHIACEAE

Malpighiaceae — S. Vogel, Mem. N.Y. Bot. Gard. 55 (1990) 133, fig. 2.

Complete; World, incl. Malesia & W. Pacific; delineated; occasional.

MALVACEAE

Hibiscadelphus Rock — R.W. Hobdy, Occ. Pap. Bish. Mus. 25, 11 (1984) 6, fig. 2.

Complete (6 spp.); Hawaiian Islands; localities of the species rougly indicated by letters; partial revision.

Hibiscus pacificus Nakai ex Jotani & Ohba — Y. Jotani & H. Ohba, J. Jap. Bot. 61 (1986) 99, fig. 2.

Complete; Bonin Is.; localities indicated; partial revision.

MELASTOMATACEAE

Brittenia subacaulis Cogn. ex Boerl. — C. Hansen, Willdenowia 15 (1985) 172, fig. 1.

Complete (monotypic genus); Borneo; localities indicated; monograph.

Campimia Ridley — C. Hansen, Willdenowia 17 (1985) 150, fig. 2.

Complete (monotypic genus); Malay Peninsula; locality indicated; monograph.

Memecylon L. — K. Bremer, Opera Bot. 69 (1983) 46, 47, fig. 1–27.

Partial (27 spp., 8 end.); Borneo; localities indicated; partial revision.

Neodriessenia Nayar — C. Hansen, Bot. Jahrb. 106 (1985) 9, fig. 2.

Complete (6 spp.); Borneo; localities indicated; monograph.

Sonerila Roxb. — R. Lundin, Nord. J. Bot . 3 (1983) 636.

Partial (22 spp., all end.); Sri Lanka; localities indicated; partial revision.

MELIACEAE

Aglaia elaeagnoidea — D.J. Mabberley, Fl. Nouv. Caléd. et Dépend. 15 (1988) 78.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

Anthocarapa nitidula — D.J. Mabberley, Fl. Nouv. Caléd. et Dépend. 15 (1988) 74.

Complete; New Caledonia; localities indicated; partial revision.

Dysoxylum — D.J. Mabberley, Fl. Nouv. Caléd. et Dépend. 15 (1988) 74.

Partial (9 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Xylocarpus — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 62, 64.

Partial (2 spp.); Australia; localities indicated; occasional.

X. rumphii, X. granatum — D.J. Mabberley, Fl. Nouv. Caléd. et Dépend. 15 (1988) 85.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

MESEMBRYANTHEMACEAE see Aizoaceae

MONIMIACEAE

Monimiaceae — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 260, fig. 3.

Partial; Old World, incl. Malesia & Pacific (New World range not indicated); delineated; occasional.

Mollinedieae — W.R. Philipson, Fl. Males. I, 10 (1986) 284, fig. 10.

Complete (6 monotypic genera); *Parakibara*, Halmahera; *Kairoa, Lauterbachia, Faika*, New Guinea; *Tetrasynandra, Austromatthaea*, Queensland; delineated; partial revision.

Faika Philipson — W.R. Philipson, Fl. Males. I, 10 (1986) 284, fig. 10.

More or less complete; W. New Guinea (distribution actually extending to the Cyclops); delineated; partial revision.

Lauterbachia — W.R. Philipson, Blumea 30 (1985) 418, fig. 1.

Complete; W. New Guinea; locality indicated; partial revision.

Matthaea Bl. — W.R. Philipson, Fl. Males. I, 10 (1986) 321, fig. 25.

Complete (all 6 spp.); Malesia (5 Philippines, *M. sancta* W. to Sumatra); delineated; partial revision.

Parakibara Philipson — W.R. Philipson, Blumea 30 (1985) 418, fig. 25.

Complete; E. Malesia: Halmahera; locality indicated; partial revision.

Steganthera Perk. — W.R. Philipson, Blumea 29 (1984) 482, fig. 1.

Partial; New Guinea; delineated + localities indicated; partial revision.

MORACACEAE

Ficus racemosa L.— E.J.H. Corner, Allertonia 4 (1985) 155, fig. 8.

More or less complete (3 var.); Asia, Malesia, Australia; delineated; occasional.

MUSACEAE

Heliconia — W.J. Kress, Allertonia 6 (1990) 22–39, fig. 4–16.

Partial (6 spp.); Old World species, E. Malesia, Pacific ; localities indicated; partial revision.

Musa — E.M. Ross, Fl. Austr. 45 (1987) 421, 422, maps 15–17.

Partial (3 spp.); Australia; localities indicated; partial revision.

MYRISTICACEAE

Gymnacranthera (A. DC.) Warb. — R.T.A. Schouten, Blumea 31 (1986) 460, fig. 1.

Complete; S. India, Malesia; delineated; monograph.

G. farquhariana (Hook.f. & Th.) Warb. — R.T.A. Schouten, Blumea 31 (1986) 478, fig. 7.

Complete (4 var.); Malesia to Bismarcks; delineated; monograph.

G. forbesii, G. ocellata — R.T.A. Schouten, Blumea 31 (1986) 473, fig. 6.

Complete; W. Malesia; delineated; monograph.

Horsfieldia Willd. — W.J.J.O. de Wilde, Gard. Bull. Sing. 37 (1985) 137, fig. 2.

Complete (3 sections); Sri Lanka, SE. Asia, Malesia E. to the Solomons, N. Australia; delineated; monograph.

Knema globularia (Lamk.) Warb. — W.J.J.O. de Wilde, Gard. Bull. Sing. 37 (1985) 121, fig. 2.

Complete; SE. Asia, W. Malesia; localities indicated; monograph.

MYRSINACEAE

Aegiceras corniculatum — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 64.

Partial; Australia; localities indicated; occasional.

Ardisia subg. **Bladhia** (Hornst.) Mez — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 209, fig. 28.

Complete (2 sections); SE. & E. Asia, W. Malesia; delineated; partial revision.

Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 216–292, fig. 30–78.

Partial (various spp.), E. Asia, Malesia; localities indicated; partial revision.

A. subg. **Chinensia** (Nakai) Yang — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 207, fig. 27.

Complete; China, Ryukyu, Malaya; delineated; partial revision.

A. bambusetorum K. & G. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 250, fig. 50.

Complete; S. Thailand, Malaya; localities indicated; partial revision.

A. cameronensis Yang — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 260, fig. 57.

Complete; Malaya; localities indicated; partial revision.

A. chinensis Benth. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 292, fig. 78.

Complete; China, Ryukyu, Malaya; localities indicated; partial revision.

A. foliosa Furt. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 254, fig. 53.

Complete; Malaya; localities indicated; partial revision.

A. maingayi (Cl.) K. & G. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 257, fig. 55.

Complete; Malaya, Hainan, Borneo; localities indicated; partial revision.

A. metallica N.E. Br. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 263, fig. 59.

Complete; W. Malaya; localities indicated; partial revision.

Ardisia theaefolia K. & G. — Y.P. Yang & J.D. Dwyer, Taiwania 34 (1989) 238, fig. 42.

Complete; Malaya, localities indicated; partial revision.

MYRTACEAE

Acmena DC. — B.P.M. Hyland, Austr. J. Bot. Suppl. ser. 9 (1983) 155, fig. 51.

Partial (7 spp.); Australia; localities indicated; partial revision.

Acmenosperma E. Kausel — B.P.M. Hyland, Austr. J. Bot. Suppl. ser. 9 (1983) 154, 160, fig. 50, 56.

Partial (2 spp.); Australia; localities indicated; partial revision.

Arillastrum — P.Y. Ladiges & C.J. Humphries, Bot. J. Linn. Soc. 87 (1983) 124, fig. 7.

Complete; New Caledonia; shaded; occasional.

Asteromyrtus Schauer — L.A. Craven, Austr. Syst. Bot. 1, 4 (1988) 378, 382, fig. 1–4.

Complete (7 spp.); N. Australia, New Guinea, Moluccas; localities indicated; monograph.

Eucalyptus — P.Y. Ladiges & C.J. Humphries, Bot. J. Linn. Soc. 77 (1983) 124–126, fig. 8–11.

Complete (+ subgenera); Australia, E. Malesia; shaded; occasional.

Eugenia reinwardtiana (Bl.) DC. — B.P.M. Hyland, Austr. J. Bot. Suppl. ser. 9 (1983) 159, fig. 55.

Partial; Australia; localities indicated; partial revision.

Leptospermum — J. Thompson, Telopea 3 (1989) 303, map 1.

Complete; Australia, Malesia; species density; monograph.

J. Thompson, Telopea 3 (1989) 330–426, maps 2–9.

Complete (79 spp.); Australia, Malesia; delineated; shaded; localities indicated; monograph.

L. parviflorum Val. — R.J. Johns, Klinkii 2 (1983) 46, fig. 2.

Partial; New Guinea; localities indicated; partial revision.

Lindsayomyrtus brachyandrus (C.T. White) Hyland & Steen. — R.J. Johns, Klinkii 2 (1983) 32, fig. 2.

Partial; New Guinea; localities indicated; partial revision.

Metrosideros bartletii Dawson — J.W. Dawson, New Zeal. J. Bot. 23 (1985) 610, fig. 3.

Complete; New Zealand (North Island); localities indicated; partial revision.

Osbornia octodonta — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 65.

Partial; Australia; localities indicated; partial revision.

Syzygium J. Gaertn. — B.P.M. Hyland, Austr. J. Bot. Suppl. ser. 9 (1983) 153–160, fig. 49–56.

Partial (52 spp.); Australia; localities indicated; partial revision.

NAJADACEAE

Najas — L. Triest, A revision of the genus Najas, Bruxelles (1988) fig. 3, facing p. 28.

Partial (11 spp.); species from ricefields; Old World incl. Malesia; species density; monograph.

N. browniana — L. Triest, A revision of the genus Najas, Bruxelles (1988) 117, fig. 20.

Complete; Australia, New Caledonia; localities indicated; monograph.

N. celebica — L. Triest, A revision of the genus Najas, Bruxelles (1988) 122, fig. 22.

Complete; Celebes; localities indicated; monograph.

N. foveolata — L. Triest, A revision of the genus Najas, Bruxelles (1988) 105, fig. 17.

Complete; Malesia (not Sumatra, Malay Peninsula, Borneo), N. Australia; localities indicated; monograph.

N. graminea — L. Triest, A revision of the genus Najas, Bruxelles (1988) 140, fig. 26.

Complete; Old World, incl. Malesia, New Caledonia; localities indicated; monograph.

N. halophila — L. Triest, A revision of the genus Najas, Bruxelles (1988) 115, fig. 19.

Complete; Java, New Guinea, Queensland; localities indicated; monograph.

N. kingii — L. Triest, A revision of the genus Najas, Bruxelles (1988) 89, fig. 13.

Complete; SE. Asia, W. Malesia (Sumatra, Malay Peninsula); localities indicated; monograph.

N. malesiana — L. Triest, A revision of the genus Najas, Bruxelles (1988) 134, fig. 25.

Complete; India to Queensland (not Philippines, New Guinea); localities indicated; monograph.

N. marina — L. Triest, A revision of the genus Najas, Bruxelles (1988) 61–71, fig. 8–10.

Complete (5 spp.); Old World, incl. Malesia; localities indicated; monograph.

Najas pseudograminea — L. Triest, A revision of the genus Najas, Bruxelles (1988) 124, fig. 23.

Complete; SE. Asia, N. Australia, Solomons; localities indicated; monograph.

NEPENTHACEAE

Nepenthaceae — W.N. Ellis, Biogeografie (1988) 115.

Complete; Old World, Madagascar to New Caledonia; delineated; occasional.

Nepenthes — B.E. Juniper, R.J. Robins & D.M. Joel, The Carnivorous plants (1989) 33, fig. 3.7.

More or less complete; Old World, Madagascar to SE. Asia, Malesia, Queensland (New Caledonia not indicated); delineated; occasional.

W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. & Evol. 162 (1989) 37, fig. 12.

Complete; Old World (fossil in Europe, present-day distribution Madagascar to SE. Asia and New Caledonia); delineated + fossil localities indicated; occasional.

M. Jebb, Science in New Guinea 17 (1991) 21, fig. 7.

Partial; New Guinea; localities indicated; partial revision.

M. Jebb, Science in New Guinea 17 (1991) 23–47, fig. 9–28.

Partial (11 spp.); New Guinea; localities indicated; partial revision.

N. bicalcarata Hook. f. — G. Zizka, Palmengarten 55 (1991) 41, fig. 7.

Complete; Borneo; localities indicated; occasional.

N. mirabilis (Lour.) Druce — T.D. Stanley, Fl. Austr. 8 (1982) 25, fig. 6.

Partial; Australia; shaded; partial revision.

NYMPHAEACEAE

Nelumbo — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 582, fig. 8.

Complete; America, Asia, Malesia, Australia; delineated; occasional.

NYSSACEAE

Nyssa — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 581, fig. 4.

Complete; N. America, SE. Asia, W. Malesia; delineated; occasional.

OCHNACEAE

Sauvagesia serrata (Korth.) Sastre — C. Sastre, C.R. Séances Soc. Biogéogr. 48 (1972) 47.

Complete; W. Malesia; localities indicated; monograph.

OLACACEAE

Olacaceae p.p. (*Anacolosa, Cathedra, Ptychopetalum*) — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. & Evol. 162 (1989) 26, 27, fig. 8.

Complete; World; fossil range of *Anacolosidites*; delineated; occasional.

Anacolosa fructescens (Bl.) Bl. — H. Sleumer, Fl. Males. I, 10 (1984) 26, fig. 14.

Complete; Thailand, W. Malesia, incl. Philippines, Celebes; delineated; partial revision.

Harmandia Pierre — H. Sleumer, Fl. Males. I, 10 (1984) 10, fig. 3.

Complete (monotypic); Indochina, Malesia, Sumatra, Borneo; localities indicated; practically a monograph.

Ochanostachys Mast. — H. Sleumer, Fl. Males. I, 10 (1984) 14, fig. 6.

Complete (monotypic); Sumatra, Malaya, Borneo; delineated; practically a monograph.

Olax, Ximenia — A.S. George, Fl. Austr. 22 (1984) 17, 35.

Partial; Australia; shaded; partial revision.

Scorodocarpus Becc. — H. Sleumer, Fl. Males. I, 10 (1984) 14, fig. 6.

Complete (monotypic); Sumatra, Malaya, Borneo; delineated; practically a monograph.

Strombosia ceylanica Gardn. — H. Sleumer, Fl. Males. I, 10 (1984) 23, fig. 11.

Complete; S. India, Ceylon, W. Malesia; delineated; partial revision.

S. javanica Bl. — H. Sleumer, Fl. Males. I, 10 (1984) 21, fig. 10.

Complete; Sumatra, Malaya, W. Java, Borneo; delineated; partial revision.

OLEACEAE

Fraxinus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 346.

Complete; World, N. hemisphere, also Luzon & Java; delineated; species density; lit. + herb.

Jasminum — P.S. Green, Allertonia 3 (1984) 410–417, fig. 4–17.

Partial (9 spp.); Australia; localities indicated; partial revision.

Myxopyrum L. — R. Kiew, Blumea 29 (1984) 507, 510, fig. 2, 3.

Complete (4 spp.); India to New Guinea & Bismarcks; localities indicated; monograph.

Nestegis — N.T. Moar, New Zeal. J. Bot. 22 (1984) 170, fig. 1.

Partial; New Zealand (South Island); localities indicated; occasional.

ONAGRACEAE

Fuchsia (sections) — P.E. Berry, Ann. Missouri Bot. Gard. 69 (1982) 5, 6, fig. 1, 2.

Partial; America; delineated; shaded; partial revision.

(Fuchsia, sections)
P.E. Berry, J.J. Skvarla, A.D. Partridge & M.K. Macphail, Austr. Syst. Bot. 3 (1990) 742, fig. 5.

Partial; Australia, New Zealand; fossil localities indicated; palynological study.

Ludwigia — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 298.

Complete; World; delineated; isolated localities indicated; lit. + herb.

ONCOTHECACEAE

Oncotheca balansae, O. humboldtiana — P.H. Morat & J.M. Veillon, Fl. Nouv. Caléd. et Dépend. 15 (1988) 94, 98.

Complete; New Caledonia; localities indicated; partial revision.

OPILIACEAE

Cansjera Juss. — P. Hiepko, Fl. Males. I, 10 (1984) 49, fig. 15.

Complete (3 spp.); India to N. Australia and Bismarcks; localities indicated; partial revision.

Champereia manillana (Bl.) Merr. — P. Hiepko, Fl. Males. I, 10 (1984) 38, fig. 3.

Partial; Australia; shaded; partial revision (after Hiepko 1979).

Lepionurus sylvestris Bl. — P. Hiepko, Fl. Males. I, 10 (1984) 44, fig. 11.

Complete (monotypic genus); SE. Asia, W. Malesia; localities indicated; partial revision (after Hiepko 1979).

Melientha Pierre — P. Hiepko, Fl. Males. I, 10 (1984) 40, fig. 6.

Complete (monotypic, 2 subsp.); SE. Asia, Malesia, Borneo, Philippines; localities indicated; partial revision (after Hiepko 1979).

Opilia — P. Hiepko, Fl. Austr. 22 (1984) 35.

Partial; Australia; shaded; partial revision.

O. amentacea Roxb. — P. Hiepko, Fl. Males. I, 10 (1984) 47, fig. 13.

Partial; India to N. Australia & Solomons (also Africa); localities indicated; partial revision.

Urobotrya Stapf sect. **Lepionuroides** Hiepko — P. Hiepko, Fl. Males. I, 10 (1984) 42, fig.8.

Complete (5 spp.); SE. Asia, Borneo, Flores; localities indicated; partial revision.

ORCHIDACEAE

Aerides — H.L. Valmayor, Orch. Philip. 1 (1984) 65.

More or less complete; SE. Asia & Malesia (Philippines more detailed); localities more or less indicated; partial revision.

Bromheadia pulchra Schltr. — P.S. Lavarack, M.A. Clements, N.H.S. Howcroft, the Orchadian 7 (1984) 294, fig. 3.

Complete; New Guinea, Australia (Cape York); localities indicated; partial revision.

Bulbophyllum — H.L. Valmayor, Orch. Philip. 1 (1984) 67.

Partial; SE. Asia, Malesia, Pacific (Philippines more detailed); localities indicated; partial revision.

B. digodence J.J.S — N.H.S. Howcroft, The Orchadian 10 (1990) 49, fig. 2.

Complete; New Caledonia; localities indicated; monograph.

Calanthe triplicata Group — S. Kobayashi, J. Phytogeogr. & Taxon. 36 (1988) 23, fig. 2.

Complete; S. Africa through Madagascar, India to N. Australia & West Pacific; delineated; occasional.

Coelogyne beccarii Reichb.f. — N.H.S. Howcroft, The Orchadian 7 (1983) 157, fig. 3.

Complete; New Guinea; localities indicated; partial revision.

Dendrobium — H.L. Valmayor, Orch. Philip. 1 (1984) 69.

Partial; SE. Asia, Malesia, Pacific (Philippines more detailed); localities indicated; partial revision.

D. sect. **Ceratobium** — P.J. Cribb, Kew Bull. 37 (1983) 582, 587, maps 1, 2.

Partial (11 spp.); Pacific; delineated or localities indicated; partial revision.

D. sect. **Oxyglossum** — T.M. Reeve & P.J.B. Woods, Notes Roy. Bot. Gard. Edinb. 46 (1989) 283, map 1.

Complete; E. Malesia to Ponape, Fiji & New Caledonia; delineated; monograph.

T.M. Reeve & P.J.B. Woods, Notes Roy. Bot. Gard. Edinb. 46 (1989) 283–290, maps 2–19.

Complete (28 spp.); E. Malesia, Pacific; localities indicated; monograph.

D. sect. **Spatulata** Lindl. — P.J. Cribb, Kew Bull. 41 (1986) 616, map 1.

Complete; Malesia, Australia, Pacific; species density; monograph.

D. sect. **Spatulata** — P.J. Cribb, Kew Bull. 41 (1986) 629–660, maps 2–7.

Partial (35 out of 46 species); Malesia, Australia, Pacific; delineated + localities indicated; monograph.

D. mirbelianum Gaud. — N.H.S. Howcroft, The Orchadian 7 (1983) 175, fig. 1.

Complete; Moluccas, New Guinea, New Ireland, Queensland; localities indicated; partial revision.

Eria — H.L. Valmayor, Orch. Philip. 1 (1984) 71.

More or less complete; SE. Asia, Malesia, Pacific; localities more or less indicated; partial revision.

Goodyera boninensis and related species — S. Kobayashi, J. Phytogeogr. & Taxon. 36 (1988) 25, fig. 4.

Complete; India to S. Japan; delineated; occasional.

Malaxis biloba group — S. Kobayashi, J. Phytogeogr. & Taxon. 36 (1988) 23, fig. 2.

Complete; S. Africa through Madagascar, India to N. Australia & West Pacific; delineated; occasional.

Microtis R. Br. — R. Bates, J. Adel. Bot. Gard. 7 (1984) 57–82, maps 1–11.

Complete; Australia, New Zealand, New Caledonia, Malesia, E. Asia; localities in Australia + New Zealand indicated; monograph.

Paphiopedilum — P. Cribb, The genus Paphiopedilum (1987) 26, map 1.

Complete; India to Solomons; delineated; species density; occasional.

P. Cribb, The genus Paphiopedilum (1987) 103, map 6.

Partial (8 spp.); Sumatra, Malay Peninsula, Borneo; delineated + localities indicated; monograph.

P. Cribb, The genus Paphiopedilum (1987) 132, map 8.

Partial (6 spp.); W. Malesia & Celebes; delineated + localities indicated; monograph.

P. Cribb, The genus Paphiopedilum (1987) 164, map 12.

Partial (5 spp.); W. Malesia & Celebes, Flores; delineated + localities indicated; monograph.

P. Cribb, The genus Paphiopedilum (1987) 170, map 13.

Partial (6 spp.); Moluccas, New Guinea, Solomons; delineated + localities indicated; monograph.

P. Cribb, The genus Paphiopedilum (1987) 195, map 15.

Partial (9 spp.); Philippines; delineated + localities indicated; monograph.

P. subg. **Brachypetalum** — Chen Sing-Chi & Tsi Zhan-Huo, Acta Phytotax. Sin. 22 (1984) 121, fig. 2.

Complete; SE. Asia, N. Malaya; delineated; partial revision.

P. Cribb, The genus Paphiopedilum (1987) 76, map 3.

Complete; SE. Asia, N. Malay Peninsula; shaded; monograph.

P. sect. **Barbata** — P. Cribb, The genus Paphiopedilum (1987) 123, map 7.

Complete; SE. Asia, Malesia incl. Solomons; delineated; monograph.

P. sect. **Cochlopetalum** — P. Cribb, The genus Paphiopedilum (1987) 123, map 7.

Complete; Sumatra, Java; shaded; monograph.

P. sect. **Coryopedilum** — P. Cribb, The genus Paphiopedilum (1987) 98, map 5.

Complete; Malesia, Borneo, Philippines, New Guinea; delineated; monograph.

P. sect. Pardalopetalum — P. Cribb, The genus Paphiopedilum (1987) 98, map 5.
Complete; SE. Asia, W. Malesia, Luzon, Celebes; delineated; monograph.

P. calosum — P. Cribb, The genus Paphiopedilum (1987) 185, map 14.
Complete; SE. Asia, Malay Peninsula; delineated; monograph.

P. niveum — P. Cribb, The genus Paphiopedilum (1987) 87, map 4.
Complete; N. Malay Peninsula, incl. Langkawi; delineated; monograph.

Phalaenopsis — H.L. Valmayor, Orch. Philip. 1 (1984) 73.
More or less complete; SE. Asia, Malesia (Philippines more detailed); localities more or less indicated; partial revision.

Renanthera — H.L. Valmayor, Orch. Philip. 1 (1984) 65.
More or less complete; SE. Asia, Malesia (Philippines more detailed); localities indicated; partial revision.

Trichoglottis — H.L. Valmayor, Orch. Philip. 1 (1984) 65.
More or less complete; SE. Asia, Malesia (Philippines more detailed); localities indicated; partial revision.

Vanda — H.L. Valmayor, Orch. Philip. 1 (1984) 65.
More or less complete; SE. Asia, Malesia (Philippines more detailed); localities more or less indicated; partial revision.

OXALIDACEAE

Oxalis corniculata L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 259.
Complete; World; delineated and shaded, isolated localities indicated; lit. + herb.

PALMAE

Palmae — J. Tivy, Biogeography ed. 2 (1982) 120, fig. 6.3.
Complete; Northern and Southern limits; delineated; after Good 1964.

N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 59, map 1.
Complete; World; shaded; monograph.

Arecoideae Uhl & Dransfield — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 311, map 26.
Complete; World, incl. Malesia & Pacific; shaded; monograph.

Calamoideae — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 234, map 11.
Complete; World, incl. Malesia & Pacific; shaded; monograph.

Ceryoxyloideae Drude — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 290, map 22.
Complete; America, Juan Fernandez, Madagascar, Queensland; shaded; monograph.

Nypoideae Griffith — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 286, map 21.

Complete; present range India to Pacific (Micronesia, Solomons) & Queensland; fossil localities indicated (America, Africa, Europe, Tasmania); monograph.

Borasseae Martius — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 217, map 8.

Complete; Old World (Africa, S. & SE. Asia, Malesia, Queensland); shaded; monograph.

Caryoteae Drude — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 312, map 27.

Complete; India to Ryukyu, Queensland & Solomons; shaded; monograph.

Ceroxyleae Satake — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 292, map 24.

Complete; S. America incl. Juan Fernandez, Madagascar, Queensland; shaded; monograph.

Coccoeae Martius — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 480, map 45.

Complete; pantropical; shaded; fossil localities indicated in Europe, India, New Zealand; monograph.

Corypheae Martius — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 167, map 2.

Complete; World, incl. Malesia & Pacific; shaded; monograph.

Phoeniceae Drude — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 215, map 7.

Complete; Old World: Africa, S. & E. Asia, W. Malesia (Sumatra, Malay Peninsula); fossil in Europe and N. America; shaded; monograph.

Archontophoenicinae Dransfield & Uhl — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 367, map 37.

Complete; Australia (E. Australia), New Zealand, New Caledonia, Lord Howe, Norfolk; shaded; monograph.

Arecinae Uhl & Dransfield — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 403, map 41.

Complete; India to N. Australia & Solomons; shaded; monograph.

Butiinae Saakov — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 485, map 47.

Complete; S. America, Madagascar to Pacific; shaded; monograph.

Calaminae Meisner — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 248, map 15.

Complete; W. Africa, SE. Asia (India to Taiwan), W. Malesia, E. Australia, Pacific E. to Fiji; shaded; monograph.

Coryphinae Dransfield & Uhl — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 207, map 5.

Complete; Arabia, S. & E. Asia, Malesia, Queensland; shaded; monograph.

Cyrtoschidinae Dransfield & Uhl — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 308, map 38.

Complete; Malesia (Sumatra, Malay Peninsula, Borneo, New Guinea, Solomons); shaded; monograph.

Eugeissoninae Becc. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 241, map 13.

Complete; Malesia (Malay Peninsula & Borneo); shaded; monograph.

Iguanurinae Hook. f. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 417, map 42.

Complete; Mascarenes, India through Malesia to Pacific (incl. Bonin, Lord Howe, Fiji); shaded; monograph.

Lataniinae Meisner — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 218, map 9.

Complete; Old World: Africa, S. & SE. Asia, Malesia, N. Australia; shaded; monograph.

Linospadicinae Hook. f. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 380, map 39.

Complete; Moluccas, New Guinea, E. Australia, Lord Howe; shaded; monograph.

Livistoninae Saakov — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 191, map 4.

Complete; World, incl. Malesia & Pacific; shaded; shaded; monograph.

Metroxylinae Bl. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 244, map 14.

Complete; SE. Asia, Malesia, Pacific (E. to Samoa); shaded; monograph.

Oncospermatinae Hook. f. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 461, map 43.

Complete; Mascarenes, Seychelles, Sri Lanka, W. Malesia incl. Philippines & Moluccas; delineated; monograph.

Oraniinae Dransfield & Uhl — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 334, map 30.

Complete; Madagascar, Malesia (not Lesser Sunda Islands); shaded; monograph.

Pigafettinae Dransfield — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 271, map 17.

Complete; E. Malesia (Celebes, Moluccas, New Guinea); shaded; monograph.

Plectocomiinae Dransfield & Uhl — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 265, map 16.

Complete; SE. Asia, W. Malesia incl. Philippines; shaded; monograph.

Ptychospermatinae Hook. f. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 387, map 40.

Complete; Malesia (Palawan, Moluccas, New Guinea); N. Australia, Pacific (E. to Fiji); shaded; monograph.

Trinacinae Becc. — N.W. Uhl & J. Dransfield, Genera Palmarum (1987) 170, map 3.

Complete; World, incl. Malay Peninsula; shaded; monograph.

Actinokentia Dammer — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 325, fig. 4.

Complete (2 spp.); New Caledonia; localities indicated; partial revision.

Alloschmidia glabrata (Becc.) H.E. Moore (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 355, fig. 21.

Complete; New Caledonia; localities indicated; partial revision.

Basselinia Vieill. sect. **Basselinia** — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 374, 375, 378, fig. 27, 28, 29.

Complete (3 spp.); New Caledonia; localities indicated; partial revision.

Basselinia Vieill. sect. **Talona** H.E. Moore — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 364, fig. 24.

Complete (7 spp.); New Caledonia; localities indicated; partial revision.

Brongniartikentia Becc. — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 343, fig. 16.

Complete (2 spp.); New Caledonia; localities indicated; partial revision.

Burretiokentia Pichi Sermolli — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 397, fig. 39.

Complete; New Caledonia; localities indicated; partial revision.

Campecarpus fulcitus (Brongn.) Wendl. (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 386, fig. 33.

Complete; New Caledonia; localities indicated; partial revision.

Chambeyronia Vieill. — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 333, fig. 10.

Complete (2 spp.); New Caledonia; localities indicated; partial revision.

Clinosperma bracteale (Brongn.) Becc. (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 339, fig. 14.

Complete; New Caledonia; localities indicated; partial revision.

Cyphokentia macrostachya Brongn. (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 355, fig. 21.

Complete; New Caledonia; localities indicated; partial revision.

Cyphophoenix Wendl. ex Hook.f. — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 386, fig. 33.

Complete (2 spp.); New Caledonia, Lifu; localities indicated; partial revision.

Cyphosperma balansae (Brongn.) Wendl. — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 393, fig. 37.

Complete; New Caledonia; localities indicated; partial revision.

Heterospathe Scheff. — E.S. Fernando, Kew Bull. 35 (1990) 220, map 1.

Complete; Philippines, Moluccas, New Guinea, Solomons, Carolines; delineated; partial revision.

Kentiopsis oliviformis (B. & G.) Brongn. (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 330, fig. 8.

Complete; New Caledonia; localities indicated; partial revision.

Lavoixia macrocarpa H.E. Moore (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 339, fig. 14.

Complete; New Caledonia; localities indicated; partial revision.

Livistona — T.C. Whitmore, Ann. Missouri Bot. Gard. 69 (1982) 673, fig. 6.

Partial; S. & E. Asia, Malesia, N. Australia; delineated; after Dransfield 1981.

Mackeea magnifica H.E. Moore (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 330, fig. 8.

Complete; New Caledonia; localities indicated; partial revision.

Moratia cerifera H.E. Moore (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 355, fig. 21.

Complete; New Caledonia; localities indicated; partial revision.

Nipa [sic] — J. Tivy, Biogeography ed. 2 (1982) 122, fig. 6–4.

Complete; Indo-Malesia (living), scattered, N. hemisphere (fossil); delineated living + fossil localities indicated; after Cain 1944 and Good 1964.

Nypa — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 17, fig. 4.

Complete; World; delineated + fossil localities indicated; after Walter 1962.

N. fruticans — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 65.

Partial; Australia; localities indicated; occasional.

W.N. Ellis, Biogeography (1988) 71, fig. 34.

Complete; World; delineated + fossil localities indicated; occasional.

Pritchardiopsis jeanneneyi Becc. (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 325, fig. 4.

Complete; New Caledonia; localities indicated; partial revision.

Veillonia alba H.E. Moore (monotypic genus) — H.E. Moore & N.W. Uhl, Allertonia 5 (1984) 3935, fig. 37.

Complete; New Caledonia; localities indicated; partial revision.

PASSIFLORACEAE

Adenia heterophylla (Bl.) Koord.— D.R. Satterthwait, Fl. Austr. 8 (1982) 157, fig. 170, 171.

Partial (2 spp.); Australia; shaded; partial revision.

Passiflora L. — D.R. Satterthwait, Fl. Austr. 8 (1982) 157, 177, fig. 172–182.

Partial (10 spp.); Australia; shaded; partial revision.

PANDANACEAE

Pandanaceae — D.M. Jarzen, Gard. Bull. Sing. 36 (1983) 167, map 1.

Complete; Old World tropics (stippled) (also Eurasia, America); fossil localities indicated; species density; lit.

PHILYDRACEAE

Helmholtzia, Philydrium lanuginosum — L.G. Adams, Fl. Austr. 45 (1987) 423, 424, maps 41, 45, 46.

Partial (2 spp.); Australia; shaded; partial revision.

PIPERACEAE

Piper methysticum, P. wichmannii — V. Lebot in P.A. Cox & S.A. Banack (eds.), Islands, Plants and Polynesians (1991) 173, fig. 2.

Complete; E. Malesia, Pacific; delineated + localities indicated; occasional.

PLANTAGINACEAE

Plantago major L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 421.

Complete; World (in Malesia & Pacific introduced); delineated; species density; lit. + herb.

PLUMBAGINACEAE

Aegialitis — P.B. Tomlinson, The Botany of Mangroves (1986) 54, fig. 3.4.

Complete (2 spp.); SE. Asia, E. Malesia, N. Australia; delineated; after Van Steenis 1949.

(Aegialitis)

R.J. Morley & J.R. Flenley in T.C. Whitmore (ed.), Biogeogr. Evol. Mal. Arch. (1987) 51, fig. 5.1.

Partial; SE. Asia, Malesia, Australia; fossil occurrence delineated; modern localities indicated; after Van Steenis 1949.

A. annulata — A.G. Wells in H.J. Teas, Biol. & Evol. of Mangroves (1983) 65.

Partial; Australia; localities indicated; occasional.

Limonium tetragonum (Thunb.) Bull. — J.R. Edmondson, Fl. Nouv. Caléd. et Dépend. 12 (1983) 134, map 31.

Partial; New Caledonia & Loyalties; localities indicated; partial revision.

POLYGALACEAE

Balgoya pacifica Morat & Meijden (monotypic genus) — Ph. Morat & R. van der Meijden, Bull. Mus. Nat. Hist. Nat. Paris IV, 13, Sect. B, Adansonia (1991) 6, fig. 2.

Complete; New Caledonia; localities indicated; monograph.

Polygala subg. **Chamaebuxus** (DC.) Duchartre — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 266.

Complete; N. America, Africa, Asia, W. Malesia; delineated; lit. + herb.

PONTEDERIACEAE

Monochoria Presl. — H.J. Aston, Fl. Austr. 45 (1987) 424, maps 47–50.

Partial (4 spp.); Australia; localities indicated + shaded; partial revision.

C.D.K. Cook in Kit Tan (ed.), Davis & Hedge Festschr. (1989) 178–182, maps 1–7.

Complete (all 8 spp.); Old World, incl. Malesia ; delineated; monograph.

POTAMOGETONACEAE

Cymodocea rotundata Ehrenb. & Hempr. ex Aschers. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 10, fig. 4.

Complete (Old World, rough); partial (Philippines, detailed); localities indicated; partial revision.

C. serrulata (R. Br.) Aschers. & Magnus — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 12, fig. 6.

Complet; (Old World, rough); partial (Philippines, detailed); localities indicated; partial revision.

Halodule pinifolia (Miki) den Hartog — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 15, fig. 8.

Complete (SE Asia–Pacific, rough); partial (Philippines, detailed); localities indicated; partial revision.

H. uninervis (Forsk.) Aschers. — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 17, fig. 10.

Complete (Old World, rough); partial (Philippines, detailed); localities indicated; partial revision.

Potamogeton wrightii Morong — G. Wiegleb, Pl. Syst. & Evol. 170 (1990) 66, fig. 13.

Complete; Asia, Malesia E. to New Guinea; localities indicated; partial revision.

Syringodium isoetifolium (Aschers.) Dandy — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 20, fig. 12.

Complete (Old World, rough); partial (Philippines, detailed); localities indicated; partial revision.

Thalassodendron ciliatum (Forsk.) den Hartog — E.G. Menez, R.C. Phillips & H.P. Calumpong, Smithson. Contr. Mar. Sc. 21 (1983) 22, fig. 14.

Complete (Old World, rough); partial (Philippines, Palawan, detailed); localities indicated; partial revision.

PRIMULACEAE

Anagallis L. (several subg. + spp. separately indicated) — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 334, 335.

Complete; World; delineated + localities indicated; species density; lit. + herb.

Lysimachia L. (subgenera separately indicated) — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 332.

Complete; delineated; isolated localities indicated; species density; lit. + herb.

L. subsect. **Eunummularia** Hand. — Mazz.- H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 333.

Complete; Europa, S. & E. Asia, Malesia; delineated; lit. + herb.

Samolus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 344.

Complete; World, incl. Pacific; delineated; lit. + herb.

PROTEACEAE

Proteaceae — T.C. Whitmore, Ann. Missouri Bot. Gard. 69 (1982) 673, fig. 5.

Partial (several genera); SE. Asia, Malesia; delineated; after Whitmore 1981.

Embotrhriinae — P.H. Weston & M.D. Crisp in J.A. Armstrong (ed.), Waratahs, Austr. Nat. Bot. Gard. Occ. Publ. 9 (1987) 30, fig. 5.
Complete (9 genera); New Guinea, Aru Is., Australia, S. America; shaded; occasional.

Alloxylon Weston & Crisp — Telopea 4 (1991) 503, fig. 2; 506, fig. 3.
Complete (4 spp.); E. Australia, New Guinea, Aru Is.; localities indicated; monograph.

Helicia Lour. — D.B. Foreman, Brunonia 6 (1983) 62, maps 1–6.
Partial; Australia (8 spp., 1 also S. New Guinea); shaded; partial revision.

P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 261, fig. 4.
Complete; S. India, Sri Lanka to S. Japan and Australia; delineated; after Sleumer 1955.

Heliciopsis — P.S. Ashton & C.V.S. Gunatilleke, J. Biogeogr. 14 (1987) 261, fig. 4.
Complete; SE. Asia, W. Malesia; delineated; after Sleumer 1955.

RAFFLESIACEAE

Rafflesia — W. Meijer, Nat. Geogr. 168 (1985) 138.
Complete; W. Malesia incl. Philippines; shaded; occasional.

K. Mat Salleh & A. Latiff, Blumea 34 (1989) 115, fig. 3.
Partial; Sabah; localities indicated; partial revision.

R. arnoldii — W. Meijer, Nat. Geogr. 168 (1985) 138.
Complete; Sumatra, Borneo; shaded; occasional.

RANUNCULACEAE

Ceratocephalus pungens Garnock-Jones — P.J. Garnock-Jones, New Zeal. J. Bot. 22 (1984) 136, fig. 2.
Complete; New Zealand (South Island); localities indicated; partial revision.

Myosurus minimus L. subsp. **novae-zelandiae** (Oliv.) Garnock-Jones — P.J. Garnock-Jones, New Zeal. J. Bot. 24 (1986) 353, fig. 3.
Complete; New Zealand; localities indicated; partial revision.

Ranunculus — H.D. Wilson & P.J. Garnock-Jones, New Zeal. J. Bot. 21 (1983) 345, fig. 3.
Partial (3 end. spp.); Stewart I.; localities indicated; partial revision.

RESTIONACEAE

Restionaceae–Centrolepidaceae — J. Muller, Ann. Missouri Bot. Gard. 71 (1984) 434, fig. 9.
Complete; S. hemisphere incl. Malesia; shaded + fossil localities indicated (also N. hemisphere); occasional.

Restionaceae — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. & Evol. 162 (1989) 44, 45, 46, fig. 15.

Complete; World; delineated; fossil localities of *Milfordia* indicated; occasional.

RHAMNACEAE

Rhamnaceae — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 278.

Complete; World; delineated; lit. + herb.

Rhamnus L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 278.

Complete; World (not in S. America and Australia), incl. Malesia; delineated; species density; lit. + herb.

RHIZOPHORACEAE

Anisophyllea — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1292, fig. 17 (as *Anisophylleaceae*).

Complete; pantropical; delineated; occasional.

Bruguiera — A.G. Wells in H.J. Teas, Biol. & Ecol. in Mangroves (1983) 66, 67.

Partial (5 spp.); Australia; localities indicated; occasional.

A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1290, fig. 11.

Complete; Old World, E. Africa to N. Australia and New Caledonia; delineated; occasional.

Bruguiera (5 spp.), **Carallia** (1 sp.), **Ceriops** (2 spp.), **Rhizophora** (4 spp.) — A. McKusker, Fl. Austr. 22 (1984) 17.

Partial; Australia; shaded; partial revision.

Carallia — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1284, fig. 1.

Complete; Old World, Madagascar to India, Queensland and Solomons; delineated; occasional.

Ceriops — A.G. Wells in H.J. Teas, Biol. & Ecol. in Mangroves (1983) 68.

Partial (2 spp.); Australia; localities indicated; occasional.

A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1290, fig. 11.

Complete; Old World, E. Africa to N. Australia and Solomons; delineated; occasional.

Combretocarpus — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1292, fig. 17.

Complete; Malay Peninsula, Borneo, delineated; occasional.

Crossostylis — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1284, fig. 1.

Complete; Solomons, New Caledonia, Vanuatu, Fiji; delineated; occasional.

Gynotroches — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1284, fig. 1.

Complete; Malesia, Queensland, Solomons; delineated; occasional.

Kandelia — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1290, fig. 11.

Complete; SE. Asia, W. Malesia; delineated; occasional.

Pellacalyx — A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1284, fig.1.

Complete; SE. Asia, W. Malesia; delineated; occasional.

Rhizophora — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 69, 70.

Partial (4 spp.); Australia; localities indicated; occasional.

P.B. Tomlinson, The Botany of Mangroves (1986) 332, 335, fig. B64, B65.

Complete (9 spp.); Indo-Malesia & Pacific & Neotropics; delineated; occasional.

A.M. Juncosa & P.B. Tomlinson, Ann. Missouri Bot. Gard. 75 (1988) 1290, fig. 11.

Complete; pantropical; delineated; occasional.

J.C. Ellison, Bot J. Linn. Soc. 105 (1991) 273, fig. 1

Complete (7 spp.); World; delineated; occasional.

R. samoensis — C.D. Woodroffe, Pacif. Sc. 41 (1987) 167, fig. 1.

Complete; Pacific, New Caledonia to Samoa; shaded; occasional.

ROSACEAE

Acaena — C. Kalkman, Bot. J. Linn. Soc. 98 (1988) 52, fig. 5.

Complete; mainly S. hemisphere, incl. New Guinea, New Zealand, Hawaii, Juan Fernandez; delineated; after Moore 1972.

A. juvenea, A. emittens — B.H. MacMillan, New Zeal. J. Bot. 27 (1989) 112, fig. 3.

Complete; New Zealand; localities indicated; partial revision.

A. profundeincisa (Bitt.) MacMillan — B.H. MacMillan, New Zeal. J. Bot. 21 (1983) 350, fig. 2.

Complete; New Zealand; localities indicated; partial revision.

A. rorida MacMillan, **A. tesca** MacMillan — B.H. MacMillan, New Zeal. J.Bot. 29 (1991) 134, fig. 3.

Complete; New Zealand; localities indicated; partial revision.

Geum — C. Kalkman, Bot. J. Linn. Soc. 98 (1988) 53, fig. 6.

Complete; World, mostly extratropical, incl. New Zealand; delineated; after Meusel 1965.

Rubus L. — N. Naruhashi & T. Sato, Tukar-menukar 2 (1983) 19–33, fig.1–15.

Partial; Malesia, Sabah; localities indicated (+ alt. range); localities; partial revision.

R. moluccanus L. var. **angulosus** Kalkm. — C. Kalkman, Blumea 29 (1984) 364, fig. 4.

Complete; SE. Asia, W. Malesia; localities indicated; partial revision.

R. moluccanus L. var. **obtusangulus** Miq. — C. Kalkman, Blumea 29 (1984) 362, fig. 3.

Complete; SE. Asia, Malesia E. to Bismarcks (not Celebes, Lesser Sunda Islands and Moluccas); localities indicated; partial revision.

RUBIACEAE

Gardenieae (tribes and subtribes, not *Coffeeae*) — E. Robbrecht & C. Puff, Bot. Jahrb. 108 (1986) 110, fig. 22.

Complete; pantropical; delineated; generic revision.

Hydnophytinae — C.R. Huxley & M.H.P. Jebb, Blumea 36 (1991) 4, fig. 2

Complete (5 genera); Andamans to Fiji; delineated; monograph.

M. Jebb in C.R. Huxley & D.F. Cutler (eds), Ant-plant interactions (1991) 375, fig. 24.1.

Complete (5 genera); Indo-Australia from SE. Asia to Queensland and Fiji; delineated; occasional.

Knoxieae — C. Puff & E. Robbrecht, Bot. Jahrb. 110 (1986) 542, fig. 20.

Complete; Africa, SE. Asia, Malesia, Queensland; delineated; partial revision.

Aidia Lour. — D.D. Tirvengadum & C. Sastre, Bull. Mus. Nat. Hist. Nat. IV, 8, sect. B, Adansonia 3 (1986) 270, pl. 4.

Partial (3 spp.); SE Asia, Malesia (Sumatra, Malaya, Philippines); delineated; partial revision.

A. cochinchinensis Lour. — K.M. Wong, Mal. Nat. J. 38 (1984) 15, fig. 4.

Complete; SE. Asia to Solomons; localities indicated; partial revision.

Anthorrhiza Huxley & Jebb — C.R. Huxley & M.H.P. Jebb, Blumea 36 (1991) 22, fig. 1.

Complete (8 spp.); E. New Guinea and islands; delineated + localities indicated; monograph.

Argostemma — B. Bremer, Ann. Missouri Bot. Gard. 76 (1989) 15–46, maps 1–28.

Partial (28 spp.); Borneo; localities indicated; partial revision.

Caelospermum Bl. — J.T. Johansson, Blumea 33 (1988) 273–275, fig. 4–8.

Complete (7 spp.); SE. Asia, Malesia, E. Australia, Pacific (Carolines, Solomons, New Caledonia); localities indicated; monograph.

Coprosma pumila complex — A.E. Orchard, Brunonia 9 (1987) 124, fig. 2, 127, fig. 4.

More or less complete (6 spp.); Australia, New Zealand (except on the Subantartic islands); localities indicated; partial revision.

Cyclophyllum — D.E. Bridson, Kew Bull. 42 (1987) 614, map 1.

Complete; E. Malesia, Australia, Pacific, E. to Polynesia; delineated; partial revision.

Dolicholobium A. Gray — M.E. Jansen & C.E. Ridsdale, Blumea 29 (1983) 292, 295, 304, fig. 3, 4, 6.

Partial (7 spp.); Solomons; shaded or localities indicated; monograph.

Galium L. — D.J. McGillivray, Telopea 2 (1983) 361–375, fig. 1–7.

Partial; Australia & New Zealand; localities indicated; partial revision.

Guettardella Benth. — M.E. Jansen, Blumea 29 (1984) 567, fig. 1.

Complete (20 spp.); SE. Asia, Malesia (not Sumatra & Java), Queensland, Pacific (E. to Fiji & Samoa); delineated + localities indicated of every species; monograph.

Lerchea L. — B. Axelius, Blumea 32 (1987) 99, 104, 108, fig. 5, 9, 12.

Complete (8 spp.); Sumatra, Java; localities indicated; monograph.

Motleya borneensis J.T. Johansson — J.T. Johansson, Blumea 32 (1987) 154, fig. 4.

Complete (monotypic genus); Borneo; localities indicated; monograph.

Myrmephytum Becc. — C.R. Huxley & M.H.P. Jebb, Blumea 36 (1991) 44, fig. 1.

Complete (8 spp.); E. Malesia (locality of *M. selebicum* in SE. Celebes not indicated; *M naumannii* in text, *M. naumannianum* in map); delineated + localities indicated; monograph.

Neonauclea Merr. — C.E. Ridsdale, Blumea 34 (1989) 198–252, fig. 6–32.

Partial (54 spp.); Malesia; localities indicated; partial revision.

Neonauclea (with myrmedomes) — C.E. Ridsdale, Blumea 34 (1989) 179, fig. 2.

Complete; Malesia (Sumatra, Borneo, Celebes, Philippines); delineated; monograph.

Paederia L. — Ch. Puff (ed.), Op. Bot. Belg. 3 (1991) 198, fig. 1.

Complete; World, mainly tropics (not Australia & Pacific); delineated; monograph.

P. foetida L. — Ch. Puff (ed.), Op. Bot. Belg. 3 (1991) 216, fig. 2.

Complete; S. & SE. Asia, Malesia (not New Guinea), Bonin Is.; localities indicated; monograph.

Ch. Puff. (ed.), Op. Bot. Belg. 3 (1991) 216, fig. 2

Complete; S. & SE. Asia, Malesia (not New Guinea), Bonin Is.; localities indicated; monograph.

P. verticillata Bl. — Ch. Puff.(ed.), Op. Bot. Belg. 3 (1991) 284, fig. 29.

Complete; Malesia (Sumatra to Philippines and Celebes); localities indicated; monograph.

Prismatomeris Thw. — J.T. Johansson, Opera Bot. 94 (1987) 15, fig. 12.

Complete; SE. Asia, W. Malesia; delineated; monograph.

J.T. Johansson, Opera Bot. 94 (1987) 16–19, fig. 13–17.

Complete (all 15 spp.); SE. Asia incl. Sri Lanka, W. Malesia incl. the Philippines; localities indicated; monograph.

Pyrostria — D.E. Bridson, Kew Bull. 42 (1987) 614, map 1.

Complete; Africa, SE. Asia, Malesia; delineated; partial revision.

Rennellia Korth. — J.T. Johansson, Opera Bot. 34 (1987) 10–15, fig. 2–6.

Complete (4 spp.); SE. Asia, W. Malesia; localities indicated; monograph.

Scyphiphora hydrophyllacea — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 70.

Partial; Australia; localities indicated; occasional.

Steenisia Bakh.f. — B. Bremer, Nordic J. Bot. 4 (1984) 338, fig. 4.

Complete; Borneo, 1 sp., Natuna Islands; localities of all species indicated; monograph.

Timonius corneri Wong — K.M. Wong, Bot. J. Linn. Soc. 98 (1988) 502, map 1.

Complete; S. Thailand, Malay Peninsula; localities indicated; partial revision.

Xanthophytum Bl. — B. Axelius, Blumea 34 (1990) 437–473, fig. 5–32.

Complete (30 spp.); SE. Asia, Malesia, Pacific (Fiji, Vanuatu); localities indicated; monograph.

RUPPIACEAE see Najadaceae

RUTACEAE

Boronia (& Boronella) — P.H. Weston, R.C. Carolin & J.A. Armstrong, Austr. J. Bot. 32 (1984) 188, fig. 1.

Complete; Australia & New Caledonia; delineated + shaded; occasional.

Comptonella Baker f. — T.G. Hartley, Bull. Mus. Nat. Hist. Nat. IV, 5, sect. B, Adansonia 4 (1982) 399–411, maps 1–6.

Complete (8 spp.); New Caledonia; localities indicated; monograph.

Medicosma Hook. f.(incl. **Coombea** Royen) — T.G. Hartley, Austr. J. Bot. 33 (1985) 36, fig. 1.

Partial (7 spp.); Australia, New Guinea; localities indicated; monograph.

T.G. Hartley, Austr. J. Bot. 33 (1985) 45, fig. 4.

Partial (8 spp.); New Caledonia; localities indicated; monograph.

T.G. Hartley, Austr. J. Bot. 33 (1985) 55, fig. 8.

Partial (7 spp.); New Caledonia; localities indicated; monograph.

SABIACEAE

Meliosma — C.F.van Beusekom & Th.P.M. van de Water, Fl. Males. I, 10 (1989) 703, 708, fig. 11, 13.

Partial (6 spp.); SE. Asia, Malesia; delineated or localities indicated; monograph.

M. simplicifolia — C.F.van Beusekom & Th.P.M. van de Water, Fl. Males. I, 10 (1989) 699, fig. 9.

Partial (6 subspp.); SE. & E. Asia, W. Malesia; delineated; monograph.

Sabia Coleb. — C.F.van Beusekom & Th.P.M. van de Water, Fl. Males. I, 10 (1989) 683, fig. 1.

Partial; SE. Asia E. to Solomons; delineated; monograph.

SANTALACEAE

Santalaceae — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 100.

Partial; New Caledonia & Loyalties; shaded; partial revision.

Amphorogyne — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 145, 146, 149.

Partial (3 spp.); New Caledonia; localities indicated; partial revision.

Anthobolus, Dendrotrophe, Dendromyza, Exocarpos, Santalum — H.J. Hewson & A.S. George, Fl. Austr. 22 (1984) 35, 47, 69.

Partial; Australia; shaded; partial revision.

Daenikera corallina — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 152.

Complete; New Caledonia; localities indicated; partial revision.

Elaphanthera baumannii — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 140.

Complete; New Caledonia; localities indicated; partial revision.

Exocarpos — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 120–134.

Partial (5 spp.); New Caledonia; localities indicated; partial revision.

Santalum austrocaledonicum — N. Hallé, Fl. Nouv. Caléd. et Dépend. 15 (1988) 108, 113.

Complete (3 var.); New Caledonia & Loyalties; localities indicated; partial revision.

SAPINDACEAE

Nephelieae — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 6, fig. 1.

Complete; Old World, incl. Malesia & Pacific (E. to Samoa, New Zealand & Hawaii); delineated; pollen study.

Alectryon pollen type — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 162, fig. 19.

Complete; E. Malesia, Australia, Pacific incl. Hawaii & New Zealand; delineated; pollen study.

Cubilia pollen type — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 162, fig. 19.

Complete; Malesia (Philippines, Borneo, Celebes, Moluccas); delineated; pollen study.

Litchi pollen type — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 162, fig. 19.

Complete; Africa, Madagascar, Mascarenes, India to S. China, Philippines, N. Queensland; delineated; pollen study.

Nephelieae (psilate) pollen — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 103, fig. 10.

Complete; Madagascar, Ceylon, Malaya, Borneo, Philippines, Celebes; shaded; pollen study.

Pometia pollen type — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 162, fig. 19.

Complete; Ceylon to Taiwan, Malesia, Pacific E. to Samoa; delineated; pollen study.

Cupanieidites (fossil localities), recent (para)syncolporate **Sapindaceae** (delineated) — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 160, fig. 18.

Complete; World; pollen study.

Cupaniopsis Radlk. — F. Adema, Thesis, Leiden (1991) 62, fig. 23.

Complete; E. Malesia, Australia; Pacific; delineated isolated; monograph.

Dictyoneura Bl. — J. van Dijk, Blumea 31 (1986) 445, fig. 2.

Complete; Malesia, Borneo eastward to Bismarcks; localities indicated; monograph.

Dimocarpus longan — R.W.J.M. van der Ham, Nephelieae Pollen, Thesis, Leiden (1990) 59, fig. 7.

Complete; SE. Asia, W. Malesia; distribution of pollen ornamentation; localities indicated; pollen study.

Dodonaea Miller — J.G. West in R.M. Barker & Greenslade, Evol. of the Flora & Fauna of Arid Austral. (1982) 332, fig. 2, 3.

Partial (several spp.); Australia; shaded; species density; partial revision.

(Dodonaea)
J.G. West, Brunonia 7 (1984) 187–192, fig. 90–101.
Partial (61 spp.); Australia ; localities indicated; partial revision.

Guioa acutifolia Radlk. — P.C. van Welzen, Leiden Bot. Ser. 10 (1990) 175, fig. 63.
Complete; E. Malesia, Queensland; localities indicated; monograph.

G. bijuga (Hiern) Radlk. — P.C. van Welzen, Leiden Bot. Ser. 10 (1990) 185, fig. 69.
Complete; W. Malesia (Sumatra, Malaya, Borneo, Palawan); localities indicated; monograph.

G. diplopetala (Hassk.) Radlk. — P.C. van Welzen, Leiden Bot. Ser. 10 (1990) 201, fig. 77.
Complete; SE. Asia, W. Malesia (Celebes); localities indicated; monograph.

G. pleuropteris (Bl.) Radlk. — P.C. van Welzen, Leiden Bot. Ser. 10 (1990) 261, fig. 110.
Complete; SE. Asia, W. Malesia (not Java), Philippines; localities indicated; monograph.

G. pubescens (Zoll. & Mor.) Radlk. — P.C. van Welzen, Leiden Bot. Ser. 10 (1990) 275, fig. 116.
Complete; W. Malesia, Palawan; localities indicated; monograph.

Jagera dasyantha (Radlk.) Reyn. — S.T. Reynolds, Austrobaileya 3 (1991) 499, map 5.
Complete; Queensland, New Guinea (incl. New Britain); localities indicated; monograph.

Lepidopetalum — P.C. van Welzen, P. Piskaut, F.J. Windradi, Blumea 36 (1992) 451, fig. 7.
Complete (6 spp.); Malesia, Solomons; delineated; monograph.

Nephelium L. — P.W. Leenhouts, Blumea 31 (1986) 377, fig. 2.
Complete; SE. Asia, W. Malesia (not E. of Celebes, Java & Philippines); delineated; species density; monograph.

Paranephelium macrophyllum King, **P. joannis** Davids — M. Davids, Blumea 29 (1984) 434, fig. 2.
Complete; Malay Peninsula, Borneo; localities indicated; monograph.

P. spirei Lecomte, **P. xestophyllum** Miq. — M. Davids, Blumea 92 (1984) 432, fig. 1.
Complete; SE. Asia, W. Malesia; localities indicated; monograph.

Sarcopteryx Radlk. — P.C. van Welzen, Blumea 36 (1991) 93, map 1, 98, map 2.
Partial (5 of 7 spp.); New Guinea (the genus also in Australia); localities indicated; partial revision.

SAURURACEAE

Saururaceae — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 584, fig. 10.
Complete (3 genera); N. America, E. Asia, W. Malesia; delineated; occasional.

SAXIFRAGACEAE

Quintinia serrata — M.S. McGlone, New Zeal. J. Bot. 23 (1985) 742, fig. 8C.
Complete; New Zealand; shaded; occasional.

SCHISANDRACEAE

Schisandraceae (Kadsura & Schisandra) — Z. Wu, Ann. Missouri Bot. Gard. 70 (1983) 581, fig. 6.
Complete; N. America, S. & E. Asia, W. Malesia; delineated; occasional.

SCROPHULARIACEAE

Euphrasia L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 407.
Complete; World, incl. Malesia, New Zealand, Juan Fernandez; delineated; isolated localities indicated; species density; lit. + herb.

W.R. Barker in B.A. Barlow, Flora & Fauna of Alpine Australasia (1986) 490, fig. 1.
Complete (various sections indicated separately); World; delineated; occasional.

W.N. Ellis, Biogeografie (1988) 29, fig. 9.
Complete; World; shaded; occasional.

Hebe pauciramosa, H. macrantha — A.D. Macdonald, Cant. Bot. Soc. J. 16 (1982) 74, 76.
Complete; New Zealand (South Island); localities indicated; partial revision.

Limnophila R. Br. — B.C. Wannan & J.T. Waterhouse, Austr. J. Bot. 33 (1985) 376, 378.
Partial ; Australia (5 spp., of which 4 also in Malesia or Pacific); localities indicated; partial revision.

Limosella L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 402.
Complete; World, incl. New Zealand; delineated + localities indicated; lit. + herb.

Lindernia dubia (L.) Penn. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 402.
Complete; America, Galapagos; delineated + localities indicated; lit. + herb.

L. procumbens (Krock.) Borb. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 402.
Complete; Eurasia, Java; delineated + localities indicated; lit. + herb.

Mazus — W.R. Barker in R.M. Barker & Greenslade, Evol. of the Flora & Fauna of Arid Austr. (1982) 342, fig. 2.

Complete; S. & E. Asia, Malesia, Australia, New Zealand; shaded; species density; after Grant 1925.

Mimulus L. (sections seperately indicated) — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 402.

Complete; World, incl. Malesia (Sumatra), New Zealand; delineated; lit. + herb.

W.R. Barker in R.M. Barker & Greenslade Evol. of the Flora & Fauna of Arid Austr. (1982) 342, fig. 2.

Complete; World, incl. New Zealand; shaded; species density; after Grant 1925.

D.Y. Hong, Ann. Missouri Bot. Gard. 70 (1983) 707, fig. 9.

Complete; World, incl. New Zealand; shaded; occasional.

Veronica — D.Y. Hong, Ann. Missouri Bot. Gard. 70 (1983) 707, fig. 9.

Complete; mainly N. hemisphere, also Malesia, Australia; delineated; occasional.

SMILACACEAE see Liliaceae

SOLANACEAE

Lycium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 385.

Complete; World (Bonin,Tonga, SE. Polynesia not indicated); delineated; lit. + herb.

Solanum L. subg. **Leptostemonum** (Dunal) Bitter — M.D. Whalen, Gentes Herb. 12 (1984) 193, fig. 6.

Complete; World; delineated; species group density; monograph.

Solanum sect. **Archaesolanum** — D.E. Symon in R.M. Barker & Greenslade, Evol. of the Flora & Fauna of Arid Austr. (1982) 335, fig. 1.

Partial; Australia; shaded; species density; partial revision.

S. sect. **Graciliflorum, Melongena, Oliganthes** — D.E. Symon in R.M. Barker & Greenslade, Evol. of the Flora & Fauna of Arid Austral. (1982) 337, fig. 2–4.

Partial; Australia; shaded; species density; partial revision.

SONNERATIACEAE

Sonneratia — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 71, 72.

Partial (2 spp.); Australia; localities indicated; occasional.

N.C. Duke & B.R. Jackes, Blumea 32 (1987) 278, fig. 1, 2.

Partial (4 spp., 3 hybr.); Australia, New Guinea, SW. Pacific; localities indicated; occasional.

SPHENOSTEMONACEAE

Sphenostemon — L. Pedley, Fl. Austr. 22 (1984) 217.
Partial; Australia; shaded; partial revision (as *Aquifoliaceae*).

STACKHOUSIACEAE

Stackhousia — W.R. Barker, Fl. Austr. 22 (1984) 191, 203.
Partial; Australia; shaded; partial revision.

STERCULIACEAE

Acropogon — Ph. Morat, Bull. Mus. Nat. Hist. Nat. Paris IV, 10, sect. B, Adansonia 2 (1988) 94–100, maps 1–3.
Partial (3 spp.); New Caledonia & Loyalties; localities indicated; partial revision.

Brachychiton — G.P. Guymer, Austr. Syst. Bot. 1 (1988) 214, fig. 7.
Complete; Australia, New Guinea; shaded; monograph.

B. carruthersii F. Muell. — G.P. Guymer, Austr. Syst. Bot. 1 (1988) 223, fig. 8.
Complete; New Guinea; localities indicated; monograph.

B. velutinosus Kost. — G.P. Guymer, Austr. Syst. Bot. 1 (1988) 260, fig. 27.
Complete; N. Queensland, E. New Guinea; localities indicated; monograph.

Heritiera littoralis — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 71.
Partial; Australia; localities indicated; occasional.

Leptonychia — J.F. Veldkamp & R.C.H. Flipphi, Blumea 32 (1987) 449, fig. 1.
Complete (3 spp.); S. India, Indo-China, Malesia (not Java, Celebes, Lesser Sunda Islands); localities indicated; monograph.

STYLIDIACEAE

Donatia novae-zelandiae — N. Gibson & J.B. Kirkpatrick, New Zeal. J. Bot. 23 (1985) 564, fig. 9.
Complete; New Zealand, Tasmania; localities indicated on a 10 km/sq. grid; occasional.

Phyllachne colensoi — N. Gibson & J.B. Kirkpatrick, New Zeal. J. Bot. 23 (1985) 564, fig. 9.
Complete; New Zealand, Tasmania; localities indicated on a 10 km/sq. grid; occasional.

SYMPLOCACEAE

Symplocos — W. Krutzsch in F. Ehrendorfer (ed.), Pl. Syst. Evol. 162 (1989) 30, 31, fig. 9.
Complete; World; delineated + fossil localities indicated; occasional.

THEACEAE

Camellia — Chang Hung Ta, Edit. Staff, J. Sun Yatsen Univ. (1981) 7, fig. 2.
Complete; SE. & S. Asia, incl. W. Malesia; species density; monograph.

THYMELAEACEAE

Linostomatinae — J.R. Clarkson, Austrobaileya 2 (1986) 204, fig. 1.
Complete; *Lophostoma* (S. America), *Linostoma, Enkleia* (SE. Asia, Malesia), *Jedda* (Queensland); delineated; partial revision.

Daphne sect. **Daphnanthes** C.A. Mey. subsect. **Daphnanthoides** (Gilg) Keissl. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 295.
Complete; Himalaya to Japan, Luzon; delineated; lit. + herb.

Kelleria Endl. — M.J. Heads, Austr. Syst. Bot. 3 (1990) 638, fig. 24.
Complete; SE. Australia, New Zealand, New Guinea, Borneo; delineated; monograph.

M.J. Heads, Austr. Syst. Bot. 3 (1990) 639–643, fig. 25–32.
Complete (11 spp.); Australia, New Zealand, Malesia; localities indicated; partial revision.

Pimelea (37 spp.) — S. Threlfall, Brunonia 5 (1982) 124, 134, 146, 155, 165, 176, 188, 194, fig. 5–12.
Partial; E. Australia; localities indicated; partial revision.

TRIMENIACEAE

Trimenia Seem. — W.R. Philipson, Fl. Males. I, 10 (1986) 329, fig. 2.
Complete (5 spp.); E. Malesia, incl. Celebes, New South Wales, Pacific, E. to Marquesas; localities indicated; monograph.

TRIURIDACEAE

Triuridaceae — J.P.M. van den Meerendonk, Fl. Males. I, 10 (1984) 110, fig. 1.
Complete; pantropical, incl. Malesia & Pacific (E. to Fiji); delineated; partial revision.

Sciaphila Bl. — J.P.M. van den Meerendonk, Fl. Males. I, 10 (1984) 110, fig. 1.
Partial; S. & E. Asia, Malesia, N. Queensland, Pacific (Carolines, Melanesia, Fiji); delineated; partial revision.

TYPHACEAE

Typha — B.G. Briggs, Fl. Austr. 45 (1987) 421, maps 10–12.
Partial (3 spp.); Australia; localities indicated; partial revision.

UMBELLIFERAE

Apium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 310.
Complete; World; delineated; species density; lit. + herb.

Eryngium L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 305.
Complete; World (Ceram sp. not indicated); delineated; species density; lit. + herb.

Hydrocotyle L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 303.
Complete; World; delineated; species density; lit. + herb.

Oenanthe L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 314.
Complete; World, incl. Malesia; delineated; species density; lit. + herb.

Pimpinella L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 312.
Complete; Old World, incl. Luzon & Java; delineated; species density; lit. + herb.

Sanicula L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 303.
Complete; Old World, incl. Malesia; delineated; isolated localities indicated; species density; lit. + herb.

H. St.John, Bot. Jahrb. 108 (1986) 60, 61, fig. 3.
Partial; Hawaiian Islands; localities of each of the 5 endemic species indicated by letters; partial revision.

URTICACEAE

Cecropiaceae (as separate family; 3 genera America; 2 genera Africa; *Poikilospermum*, SE. Asia, Malesia) — K. Bonsen & B.J.H. ter Welle, Bull. Mus. Nat. Hist. Nat. IV, 5, sect. B, Adansonia 2 (1983) 152.
Complete; delineated and shaded; distribution of *Poikilospermum* after Chew Wee Lek 1963.

Australina pusilla (Poir.) Gaud. — I. Friis & C.M. Wilmot-Dear, Nordic J. Bot. 8 (1988) 55, fig. 20.
Complete (12 spp.); SE. Australia & Tasmania, New Zealand; localities indicated; monograph.

Debregeasia Gaud. — C.M. Wilmot-Dear, Kew Bull. 43 (1988) 679–687, maps 1–3.
Complete (4 spp.); E. Africa, SE Asia, Malesia; localities indicated; monograph.

Droguetia iners (Forssk.) Schweinf. subsp. **urticoides** (Wight) Friis & Wilmot-Dear — I. Friis & C.M. Wilmot-Dear, Nordic J. Bot. 8 (1988) 55, fig. 20.
Complete; S. India, Java; localities indicated; partial revision.

VALERIANACEAE

Valeriana L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 3 (1992) 433.
Complete; World, also Sumatra & Java; delineated; species density; lit. + herb.

VERBENACEAE

Avicennia — A.G. Wells in H.J. Teas, Biol. & Ecol. of Mangroves (1983) 60, 61.

Partial (2 spp.); Australia; localities indicated; occasional.

N.C. Duke, Austr. Syst. Bot. 4 (1991) 300, 301, fig. 1–3.

Partial (5 spp.); Australia, E. Malesia, Pacific; localities indicated; partial revision.

N.C. Duke, Austr. Syst. Bot. 4 (1991) 304, fig. 4.

Complete (all 8 spp.); World; delineated + shaded; partial revision.

Faradaya F. Muell. — A.A. Munir, J. Adel. Bot. Gard. 10 (1987) 167, map 1.

Complete; Borneo to Queensland, Samoa & Tonga; delineated; partial revision.

Gmelina L. — A.A. Munir, J. Adel. Bot. Gard. 10 (1987) 167, map 1.

Partial (5 spp.); Australia; delineated; localities of each species indicated; partial revision.

Premna L. — A.A. Munir, J. Adel. Bot. Gard. 10 (1987) 167, map 1.

Partial (7 spp.); Australia; delineated; localities of each species indicated; partial revision.

Verbena officinalis L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 368.

Complete; World (in Malesia & Pacific introduced only); delineated; lit. + herb.

Viticipremna Lam. incl. **V. vitilevuensis** Munir — A.A. Munir, J. Adel. Bot. Gard. 10 (1987) 167, map 1.

Complete (all 5 spp.); Philippines, E. Malesia incl. Bismarks, Queensland, Fiji; delineated + localities indicated; monograph.

VIOLACEAE

Hybanthus Jacq. — A.S. George, Fl. Austr. 8 (1982) 111, fig. 107–120.

Partial (10 spp.); Australia; shaded; partial revision.

Hymenanthera dentata R. Br. — A.S. George, Fl. Austr. 8 (1982) 127, fig. 121.

Partial; Australia; shaded; partial revision.

Rinorea — W.H.A. Hekking, Fl. Neotr. 46 (1988) 31, fig. 10.

Complete; pantropical, incl. Malesia & Pacific, E. to Samoa; shaded; localities indicated; partial revision.

Viola L. — H. Meusel et al., Vergl. Chorol. Zentraleur. Fl. 2 (1978) 289.

Complete (sections separately indicated); World; delineated; species density; lit. + herb.

L.G. Adams, Fl. Austr. 8 (1982) 101, 111, fig. 92–106.

Partial (8 spp.); Australia; shaded; partial revision.

(Viola)
H. St.John, Bot. Jahrb. 111 (1989) 202, fig. 18; 203. fig. 19.
Partial; Hawaiian Islands; shaded; partial revision.

VISCACEAE see Loranthaceae

VITACEAE

Ampelocissus acetosa (F.Muell.) Planch. — B.R. Jackes, Austrobaileya 2 (1984) 83, map 1.
Partial; N. Australia, S. New Guinea; localities indicated; partial revision.

Cissus aristata Bl. — B.R. Jackes, Austrobaileya 2 (1984) 504, map 6.
Partial; Torres Strait, Australia (the species widespread in Asia & Malesia); localities indicated; partial revision.

WINTERACEAE

Winteraceae (several genera) — T.C. Whitmore, Ann. Missouri Bot. Gard. 69 (1982) 672, fig. 4.
Partial; Malesia, Australia, Pacific; delineated; after Whitmore 1981.

XANTHORRHOEACEAE see Liliaceae

XYRIDACEAE

Xyris bancana Miq. — B. Hansen, Fl. Males. I, 9 (1982) 572, fig. 4.
Complete; SE. Asia, Malesia (not Sumatra, Java, Philippines, Celebes); localities indicated; partial revision.

ZANNICHELLIACEAE see Potamogetonaceae

ZINGIBERACEAE

Alpinia (6 spp.), **Amomum** (2 spp), **Costus** (1 spp), **Curcuma** (2 spp), **Etlingera** (1 spp), **Globba** (1 spp.), **Hedychium** (2 spp.), **Hornstedtia** (1 spp.), **Kaempferia** (1 sp.), **Tapeinochilus** (1 sp.), **Zingiber** (2 spp.) — R.M. Smith, Fl. Austr. 45 (1987) 422, 423, maps 18–37.
Partial; Australia; localities indicated + shaded; partial revision.

Alpinia (2 subg., 11 sect., 12 subsect.) — R.M. Smith, Edinb. J. Bot. 47 (1990) 66–74, maps 1–18.
Complete; Indo-Australia (from India to Ryukyu, E. Australia & Samoa); delineated; monograph.

ZOSTERACEAE see Potamogetonaceae