

FLORAE MALESIANAE PRECURSORES XLII
NOTES ON SAPINDACEAE II. TRISTIOPSIS

P. W. LEENHOUTS

***Tristiopsis ferruginea* Leenh., nov. sp.**

Ramuli 8 mm crassi, novelli dense ferrugineo-tomentelli, vetustiores glabrescentes. Petiolus 8 cm longus; petioluli 1—2 mm longi; foliola alterna vel rarius (sub)opposita, ovato-oblonga vel lanceolata, 5—10 cm longa, $1\frac{3}{4}$ —3 cm lata, subtus in nervorum axillis breviter barbata, supra in costa tomentella, ceterum glabra, basi decurrentia, apice subcaudato-acuminata, acumine acuto, costa supra prominula, nervis primariis utrinque c. 15, subcurvatis, in superiore dimidio foliolorum tantum arcuato-anastomosantibus, subtus prominulis, intermediis nervis primariis \pm parallelis. Thyrsi usque ad 17 cm longi, ut ramuli novelli tomentelli; pedunculi 2—3 $\frac{1}{2}$ cm longi; pedicelli 1—2 mm longi. Sepala late ovata vel late obovata, 2 $\frac{1}{2}$ mm longa, 2 $\frac{1}{4}$ mm lata, ut videtur haud petaloidea, intus subdense adpresse pubescentia, integra. Petala per 1 $\frac{1}{4}$ mm unguiculata, lamina transverse semi-elliptica, 1 mm longa, 1 $\frac{1}{2}$ mm lata, ad marginem unguis lanata, ceterum dense ciliata, intus (unguie lamina densius) lanata; squama lamina dimidio brevior, biloba, lanata. Discus 5-lobatus, lobis annuli forma, profunde excavatis, parte centrum floris spectante dense pilosa excepta glabris. Stamina 8; filamenta 2 mm longa, in parte $\frac{2}{3}$ inferiore subdense lanata; antherae $\frac{3}{4}$ mm longae, glabrae. Pistillodium pyriforme, 1 $\frac{1}{2}$ \times 1 mm, dense adpresse pilosum.

BORNEO. Sarawak: Baram Distr., G. Api, Melinau Gorge, c. 200 m, fl. 15-7-1961, *Anderson 4626* (L). — E. Borneo: E. Kutei, Sg. Menubar, Sangkulirang, 15 m, st. 13-6-1951, *Kostermans FRI. bb. 34689* (L) vern. *laras*; Mt Medadam N. of Sangkulirang, 400 m, fr. 9-8-1957, *Kostermans 13428* (L); Berau, Mt Njapa on Kelai River, 100 m, fr. 17-10-1963, *Kostermans 21330* (L). — Sabah: Mt Kinabalu, Koung, 500 m, fl. 8-5-1933, *Carr SF. 27276* (SING) *Typus*.

T. ferruginea differs from *T. acutangula* Radlk. — the only other species of the genus — mainly by the orange-brown to ferruginous instead of fulvous indument, the subcaudate instead of emarginate to acute leaflets, the not — instead of partly — petaloid sepals, and the long-clawed instead of cuneate petals which are inside woolly instead of glabrous.