

NOTULAE AD FLORAM AGARICINAM NEERLANDICAM — VI-VII
RICKENELLA AND OMPHALINA

THOMAS W. KUYPER

Rijksherbarium, Leiden

VI. *Rickenella* Raithelhuber

Rickenella swartzii (Fr.) Kuyp., comb. nov. — *Agaricus swartzii* Fr., Obs. mycol. 1: 90. 1815 (basionym).

This species has been known thus far as *Rickenella setipes* (Fr.: Fr.) Raithelhuber. However, the original diagnosis of *Agaricus setipes* (Fries, 1818: 162) clearly refers to another species, as *A. setipes* was characterized as follows: pileus 5–8 mm broad, campanulato-convex; stipe very long, 85–100 mm, setaceous; lamellae greyish; strongly resembling *Agaricus corticola* Pers.: Fr. As far as I can ascertain this protologue refers to a species of *Mycena*. The misapplication of the name *Agaricus setipes* might have arisen when Fries (1873: pl. 75 f. 4) published an illustration of an unnamed variety of *A. setipes*, which undoubtedly represents *Rickenella swartzii*. This unnamed variety was later (Fries, 1874: 164) called *A. setipes* var. *acrocyanus*.

The use of the epithet *swartzii* is in accordance with Kühner (1938: 608), Smith (1947: 123) and Dennis & al. (1960: 122).

VII. *Omphalina* Quél.

Furthermore a new combination in *Omphalina* seems desirable:

Omphalina viridis (Hornem.) Kuyp., comb. & stat. nov. — *Agaricus umbelliferus* var. *viridis* Hornem., Fl. dan. 10 (28): 10. 1819 (basionym).

REFERENCES

- DENNIS, R. W. G., ORTON, P. D. & HORA, F. B. (1960). New check list of British agarics and boleti. In *Trans. Br. mycol. Soc.* 43 (Suppl.): 1–224.
FRIES, E. M. (1818). *Observationes mycologicae Pars 2.* Hafniae.
— (1867–1884). *Icones selectae Hymenomycetum nondum delineatorum.* Holmiae.
— (1874). *Hymenomycetes europaei.* Upsaliae.
KÜHNER, R. (1938). Le genre *Mycena*. In *Encycl. mycol.* 10.
SMITH, A. H. (1947). North American species of *Mycena*. In *Univ. Mich. Stud. scient. Ser.* 17.