



A taxonomic review of *Penicillium* species producing conidiophores with solitary phialides, classified in section *Torulomyces*

C.M. Visagie^{1,2,3}, J. Houbraken², J. Dijksterhuis², K.A. Seifert³, K. Jacobs¹, R.A. Samson²

Key words

beta-tubulin
calmodulin
Eupenicillium
internal transcribed spacer rDNA region
low temperature Scanning Electron
Microscopy (cryo-SEM)
Monocillium
RNA polymerase II second
largest subunit
Trichocomaceae
Wollemi pine

Abstract The genus *Torulomyces* was characterised by species that typically have conidiophores consisting of solitary phialides that produce long chains of conidia connected by disjunctors. Based on the phylogenetic position of *P. lagena* (generic ex-neotype), the genus and its seven species were transferred to *Penicillium* and classified in sect. *Torulomyces* along with *P. cryptum* and *P. lassenii*. The aim of this study was to review the species currently classified in sect. *Torulomyces* using morphology and phylogenies of the ITS, *BenA*, *CaM* and *RPB2* regions. Based on our results, we accept 16 species in sect. *Torulomyces*, including 12 new species described as *P. aeris*, *P. austriaca*, *P. cantabricum*, *P. catalanicum*, *P. oregonense*, *P. marthae-christenseniae*, *P. riverlandense*, *P. tubakianum*, *P. variatense*, *P. williamettense*, *P. wisconsinense* and *P. wollemiicola*. In addition, we reclassify *P. laeve* and *P. ovatum* in sect. *Exilicaulis* and correct the typification of *P. lagena*. We provide descriptions and notes on the identification of the species.

Article info Received: 17 June 2015; Accepted: 26 November 2015; Published: 8 February 2016.

INTRODUCTION

Delitsch (1943) introduced the hyphomycete genus *Torulomyces* with *T. lagena* as generic type. The genus included species producing conidiophores consisting of solitary, swollen phialides borne on short, unbranched stipes (Delitsch 1943, Barron 1967, Gams 1971, Stolk & Samson 1983, Ando et al. 1998, Seifert et al. 2011). Conidia are linked together by connectives, i.e. short intercalary cylinders of cell wall material, to form very long chains (Gams 1971, Stolk & Samson 1983, Ando et al. 1998). Colonies of *T. lagena* and related species are brownish and their slow growth on agar media means that they are overgrown easily by other fungi and are infrequently isolated.

Stolk & Samson (1983) linked *T. lagena* with the sexual morph *Eupenicillium limoneum*, suggesting an association with *Penicillium* and introduced *P. lagena* as a new combination. Later, they proposed a sectional classification for *Penicillium* asexual morphs, including sect. *Torulomyces* for *P. lagena* (Stolk & Samson 1985). Pitt & Hocking (1985) argued that *T. lagena* did not fit the generic concept of *Penicillium*, at that time defined by a conidiophore with branches forming a penicillus. The transfer also was not accepted in the list of Current Names in Use for the *Trichocomaceae* (Pitt & Samson 1993, Pitt et al. 2000).

The taxonomy of such morphologically undifferentiated hyphomycetes is confusing, and prior to the availability of DNA-based phylogenies it was difficult to evaluate the phylogenetic significance of characters such as colony colours, phialide

shape and the nature of connectives between individual conidia in chains. Apart from *T. lagena*, 10 similar fungi were classified as monophialidic species of *Paecilomyces* (Onions & Barron 1967), but following the conclusions of Gams (1971), the monograph of Samson (1974) excluded these from *Paecilomyces*. They were distributed by various authors into genera such as:

- i. *Monocillium* S.B. Saksena (1955), sexual morphs in *Niesslia* (*Niessliaceae*, *Hypocreales*), including species with thick-walled, centrally swollen phialides, conidial chains lacking connectives, or often slimy rather than in chains, sometimes confused with *Torulomyces* (Hashmi et al. 1972);
- ii. *Phialosimplex* Sigler et al. (2010), sexual morphs unknown, including species with thin-walled, scarcely swollen mono- or polyphialides, and conidial chains with connectives, proposed as a synonym of *Aspergillus* (*Aspergillaceae*, *Eurotiales*) by Houbraken & Samson (2011) and Samson et al. (2014), but perhaps phylogenetically distinct (Tanney, pers. comm.);
- iii. *Sagenomella* W. Gams (1978), sexual morphs *Sagenoma* (*Trichocomaceae*, *Eurotiales*), including species with thin-walled, unswollen mono- or polyphialides, and conidial chains with connectives;
- iv. *Taifanglania* Z.Q. Liang et al. (2009), sexual morphs unknown but *Chaetomiaceae*, *Sordariales*, including species with thin-walled, flask-shaped monophialides and conidial chains without connectives, now considered a synonym of *Acrophialophora* Edward (Zhang et al. 2015).

Thus, it is clear that the combination of solitary phialides with chains of ameroconidia is plesiomorphic, and its notable this combination of characters appears in several unrelated monophyletic groups, nested among species with more complex conidiophores. In particular, *Phialosimplex* appears to represent a reduced form of the vesiculate *Aspergillus* conidiophores,

¹ Department of Microbiology, University of Stellenbosch, Private Bag X1, Stellenbosch 7600, South Africa; corresponding author e-mail: kj@sun.ac.za.

² CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands.

³ Biodiversity (Mycology), Eastern Cereal and Oilseed Research Centre, 960 Carling Ave., Ottawa, Ontario, K1A 0C6, Canada; corresponding author e-mail: cobusvisagie9@gmail.com.

Table 1 Strains used for the phylogenetic analysis.

Species name	Collection accession nr.	Substrate, Locality	GenBank accession nr.			
			ITS	BenA	CaM	RPB2
<i>P. aëris</i>	CBS 135897 [†] = DTO 207D4	Unknown origin	KF303654	KF303614	KF303627	KF303681
<i>P. austriicola</i>	CBS 135899 = CV 1811 = DTO 186D8	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	KF303672	JX091578	JX141599	KF303704
	CBS 135900 [†] = CV 1842 = DTO 183E6 = DAOM 241066	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091466	JX091579	JX141600	KF303705
	CBS 135901 = CV 1902 = DTO 183F5	Bract from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091467	JX091586	JX141602	KF303703
	CBS 135902 = CV 1943 = DTO 183F8 = DAOMC 241067	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091468	JX091587	JX141603	KF303701
	CBS 135903 = CV 1954 = DTO 183F9	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091469	JX091588	JX141604	KF303699
	CBS 135904 = CV 1808 = DTO 183D8 = DAOMC 241065	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091465	JX091584	JX141598	KF303702
<i>P. cantabricum</i>	CBS 136253 = CV 1968 = DTO 186G3	Mite from <i>Protea repens</i> infructescens, Struisbaai, South Africa	KF303671	JX091589	JX141605	KF303700
	CBS 120415 [†] = DTO 7619 = FMR 9121	Soil, Cantabria, Reinosa, Spain	KF303655	KF303615	KF303646	KF303682
<i>P. catalanicum</i>	CBS 110532 [†] = DTO 78H5	Soil, Catalonia, Montseny Natural Park, Spain	KF303650	KF303609	KF303644	KF303683
<i>P. corylophilum</i>	CBS 330.79 = IJFM 5147	Air sample, Barcelona, Spain	GU944557	GU944519	GU944607	JN406569
<i>P. cryptum</i>	CBS 271.89 [†] = DTO 122C9 = ATCC 60138 = IMI 296794 = NRRL 13460	Soil from <i>Quercus-Betula</i> forest, New York, USA	KF303647	KF303608	KF303628	JN121478
<i>P. dimorphosporum</i>	CBS 456.70 [†] = NRRL 5207 = ATCC 22783 = ATCC 52501 = FRR 1120 = IMI 149680	Mangrove swamp soil, Tooraddin, Australia	AF081804	KJ834448	KP016783	JN121517
	CBS 136665 [†] = DTO 270G8 = KY 12727 = NBRC 109724	Soil, unknown, Thailand	KF667369	KF667365	KF667367	KF667371
<i>P. legena</i>	CBS 129212 = DTO 199C2	Unknown, Halsey National forest, Nebraska, USA	KF303662	KF303616	KF303631	KF303680
	CBS 129389 = DTO 202C5	Unknown, Wisconsin, USA	KF303663	KF303617	KF303632	KF303679
	CBS 129620 = DTO 206D2	Unknown, Minnesota, USA	KF303664	KF303618	KF303633	KF303678
	CBS 185.65 [†] = DTO 7718 = MUCL 8221 = JCM10149 = OAC10034	Bog soil under <i>Thuja plicata</i> , Guelph, Canada	KF303665	KF303619	KF303634	JN121450
<i>P. lassenii</i>	CBS 277.70 [†] = DTO 95D6 = NRRL 5272 = ATCC 22054 = FRR 858 = IMI 148395	Soil under conifers, California, USA	KF303648	KF303607	KF303629	JN121481
<i>P. marthae-christenseniae</i>	CBS 129213 [†] = DTO 201B5	Unknown, Wisconsin, USA	KF303651	KF303613	KF303645	KF303711
	CBS 129775 [†] = DTO 208A5	Unknown, Williamette National Forest, near Blue River, Oregon, USA	KF303668	KF303623	KF303640	KF303710
<i>P. oregonensis</i>	CBS 136664 [†] = DTO 270G7 = KY 12726	Soil under <i>Pinus caribaea</i> leaf litter, Kuala Lumpur, Malaysia	KF667370	KF667366	KF667368	KF667372
<i>P. porphyreum</i>	CBS 132154 = DTO 210E2 = WSF 3133	Unknown, Wisconsin, USA	KF303661	KF303620	KF303635	KF303676
	CBS 382.64 [†] = DTO 78G7 = KY 12723	Soil in community of <i>Pinus strobus</i> , Wisconsin, USA	KF303666	KF303621	KF303636	KF303677
<i>P. restrictum</i>	CBS 367.48 [†] = ATCC 11257 = FRR 1748 = IMI 040228 = NRRL 1748	Soil, Honduras	AF033457	KJ834486	KP016803	JN121506
<i>P. riverlandense</i>	CBS 135883 = CV 1360 = DTO 183A4 = DAOMC 241061	Bract from <i>Protea repens</i> infructescens, Malmesbury, South Africa	JX091458	JX091576	JX141594	KF303686
	CBS 135884 = CV 1448 = DTO 183A9 = DAOMC 241063	Bract from <i>Protea repens</i> infructescens, Malmesbury, South Africa	JX091459	JX091582	JX141595	KF303684
	CBS 135885 = CV 1450 = DTO 183B1 = DAOMC 241064	Bract from <i>Protea repens</i> infructescens, Malmesbury, South Africa	JX091460	JX091583	JX141596	KF303695
	CBS 135886 = CV 1583 = DTO 186B8	Bract from <i>Protea repens</i> infructescens, Malmesbury, South Africa	KF303657	JX091577	JX141597	KF303689
	CBS 135887 = CV 2815 = DTO 184A1	Soil, Malmesbury, South Africa	KF303673	JX091590	JX141606	KF303691
	CBS 135888 = CV 2819 = DTO 184A4	Soil, Malmesbury, South Africa	JX091462	JX091591	JX141607	KF303693
	CBS 135889 = CV 2822 = DTO 184A6	Soil, Malmesbury, South Africa	JX091463	JX091592	JX141608	KF303694
	CBS 135890 = CV 2831 = DTO 184B4	Soil, Malmesbury, South Africa	KF303668	JX091593	JX141612	KF303696
	CBS 135891 = CV 2833 = DTO 184B6	Soil, Malmesbury, South Africa	JX091464	JX091594	JX141613	KF303692
	CBS 135892 = CV 2839 = DTO 184C2	Soil, Malmesbury, South Africa	KF303659	JX091595	JX141615	KF303697
	CBS 135893 = CV 2849 = DTO 184C6	Soil, Malmesbury, South Africa	KF303660	JX091596	JX141616	KF303698
	CBS 135894 = CV 0933 = DTO 185F9	Soil, Malmesbury, South Africa	KF303656	JX091575	JX141591	KF303690
	CBS 135895 = CV 0959 = DTO 182E7 = DAOMC 241059	Soil, Malmesbury, South Africa	JX091456	JX091581	JX141592	KF303688
	CBS 135896 [†] = CV 0979 = DTO 182F6 = DAOMC 241060	Soil, Malmesbury, South Africa	JX091457	JX091580	JX141593	KF303685
	DTO 183E8 = CV 1851	Bract from <i>Protea repens</i> infructescens, Struisbaai, South Africa	JX091461	JX091585	JX141601	KF303687
	<i>P. toxicarium</i>	NRRL 6172	Unknown, Strain of <i>P. toxicarium</i> Miyake <i>vide</i> Serra et al. (2008)	EF198650	EF198620	EF198631
<i>P. tubakianum</i>	CBS 287.66 [†] = DTO 138D9 = MUCL 8519 = IFO 8315	Dead bark from <i>Cyathea</i> , New Zealand	KF303652	KF303611	KF303637	KF303712
<i>P. varratense</i>	CBS 337.97 [†] = DTO 137C8	Humus soil, Varrata National Park near Port Moresby, Papua New Guinea	KF303649	KF303610	KF303630	KF303675
	CBS 129774 [†] = DTO 208A4	Unknown, Williamette National Forest, near Blue River, Oregon, USA	KF303667	KF303622	KF303639	KF303709
<i>P. williamettense</i>	CBS 128279 [†] = DTO 198H7	Unknown, Wisconsin, USA	KF303670	KF303624	KF303641	KF303706
	CBS 128412 = DTO 201B3	Unknown, Wisconsin, USA	KF303674	KF303625	KF303642	KF303707
	CBS 128421 = DTO 198I5	Unknown, Wisconsin, USA	KF303669	KF303626	KF303643	KF303708
<i>P. wollemicola</i>	CBS 137177 [†] = DTO 297E3	<i>Wollemia nobilis</i> needle debris, Australia	KJ174314	KJ174315	KJ174316	KJ174313




Fig. 1 Phylogenetic trees for ITS, *BenA*, *CaM* and *RPB2* datasets of *Penicillium* sect. *Torulomyces*. Branch support in nodes higher than 80 % bs and/or 0.95 pp are indicated above thickened branches (| = ex-type; * = 100 % bs or 1.00 pp; - = support lower than 80 % bs and/or 0.95 pp). Species are indicated by coloured blocks.




Fig. 2 Multigene phylogeny for a combined ITS, *BenA*, *CaM* & *RPB2* dataset of *Penicillium* sect. *Torulomyces*. Branch support in nodes higher than 80 % bs and/or 0.95 pp are indicated above thickened branches († = ex-type; * = 100 % bs or 1.00 pp; - = support lower than 80 % bs and/or 0.95 pp). Species are indicated by coloured blocks.

Taifanglania a reduced form of the verticillate *Acrophialophora* conidiophore, and *Torulomyces* a reduced form of the highly branched *Penicillium* conidiophore.

Following a four gene phylogenetic analysis of *Penicillium*, *Aspergillus* and related genera, Houbraken & Samson (2011) accepted *Penicillium lagena* as the correct classification for Delitsch’s *T. lagena*, and confirmed *Torulomyces* as a synonym of *Penicillium*. They transferred seven species from *Torulomyces* to *Penicillium* sect. *Torulomyces*, making the new combinations *P. laeve*, *P. ovatum*, *P. parviterrucosum* and *P. porphyreum* (for *Monocillium humicola* var. *brunneum* ≡ *T. brunneum*). Although they lack conidiophores with solitary phialides, *P. cryptum* and *P. lassenii* were included in the section. *Torulomyces macrosporus* was not transferred because it probably belongs in *Monocillium* based on the protologue of Matsushima (1987) (Ando et al. 1998). No type material was available for the second original species of *Torulomyces*, *T. viscosus*, and the original description is insufficient for determining the placement of the species (Stolk & Samson 1983, Houbraken & Samson 2011).

In this study, we re-evaluate the taxonomy of *Penicillium* sect. *Torulomyces* following the standardised methods and taxonomic approach advocated by Visagie et al. (2014b). We deline-

ate species using phylogenies based on the nuc rDNA internal transcribed spacer region (ITS), and partial *BenA* (β -tubulin), *CaM* (calmodulin) and *RPB2* (RNA polymerase II second largest subunit) gene sequences, as well as morphological characters. We describe 12 new species and provide descriptions for the six other accepted species, while also correcting the typification of *P. lagena*. Based on phylogenetic results, we classify *P. laeve* and *P. ovatum* in *Penicillium* sect. *Exilicaulis*. The most important morphological characters needed for species identification are summarised in table format.

MATERIALS AND METHODS

Strains

Strains used for this study (summarised in Table 1) were obtained from the CBS-KNAW Fungal Biodiversity Centre, the Netherlands (CBS) and the working collection of the Applied and Industrial Mycology department (DTO) at the same institute. Ex-type and representative cultures were also deposited in the Canadian Collection of Fungal Cultures, Ottawa (DAOMC). Reference material for *P. laeve* and *P. ovatum* was received from the Biological Resource Centre at the National Institute of Technology and Evaluation, Chiba, Japan (NBRC).

Table 2 Length of datasets and models used for phylogenetic analysis.

	ITS	<i>BenA</i>	<i>CaM</i>	<i>RPB2</i>	Concatenated
Length	513 bp	368 bp	551 bp	753 bp	2185 bp
Model Maximum Likelihood	T92+G	K2+G	T92+G+I	TN93+G+I	TN93+G+I
Model Bayesian inference	GTR+G	HKY+G+I	HKY+G+I	GTR+G+I	GTR+G+I

Abbreviations used in this table: T92 = Tamura 3-parameter; K2 = Kimura 2-parameter; TN93 = Tamura-Nei; GTR = General Time Reversible; HKY = Hasegawa-Kishino-Yano; +G = Gamma distribution; +I = Invariant sites.

Table 3 Summary of the most important morphological characters for species identification.

	Growth rate (in mm)											Conidia					
	CYA37°C	CYA30°C	CYAS	CREA	CYA	OA	MEA	YES	SNA	DG18	Brown soluble pigments	Acid on CREA	Conidiophore branching pattern	Ornamentation	Shape	Tubercle size (µm)	Connectives
<i>P. aeris</i>	4–5	9–10	6–7	no growth	11–12	9–10	6–7	10–11	8–9	11–12	absent	absent	solitary phialide	rough	globose	0.21–0.4	long, lack visible rings
<i>P. austrocola</i>	no growth	10–11	8–10	3–4	10–13	9–10	4–8	10–11	9–11	9–11	present	absent	solitary phialide	rough	globose	0.19–0.30	long, lack visible rings
<i>P. centabricum</i>	3–5	12–13	3–4	3–4	13–14	10–11	6–7	9–10	7–8	8–9	absent	weak	solitary phialide	rough	globose	0.21–0.33	present, lack visible rings
<i>P. catalanicum</i>	no growth	no growth	2–3	no growth	6–8	6–7	4–5	5–6	6–7	7–8	absent	absent	solitary phialide	rough, in parallel rings	subglobose to broadly ellipsoidal	0.12–0.18	short, visible rings present
<i>P. cryptum</i>	no growth	6–9	no growth	no growth	6–7	5–6	6–7	5–6	5–6	1–2	absent	absent	solitary phialide to monoverrucillate	smooth	broadly ellipsoidal	n.a.	short, lack visible rings
<i>P. laeve</i>	no growth	4–5	no growth	no growth	8–9	7–8	micro-colonies	8–9	7–8	5–7	absent	absent	solitary phialide	smooth	globose	n.a.	long, lack visible rings
<i>P. legena</i>	no growth	8–10	no growth	no growth	8–10	8–12	8–10	11–13	10–13	11–13	present	absent	solitary phialide	rough	globose	0.17–0.29	long, lack visible rings
<i>P. lasserii</i>	no growth	4–5	4–5	3–4	7–9	5–6	9–10	9–10	5–6	10–11	absent	absent	mono- to biverticillate	smooth	broadly ellipsoidal	n.a.	n.a.
<i>P. marthae-christensense</i>	no growth	5–6	no growth	no growth	7–8	9–10	4–5	7–8	11–12	9–11	absent	absent	solitary phialide	rough	subglobose	0.22–0.32	present, visible rings present
<i>P. oregonense</i>	no growth	no growth	no growth	no growth	4–5	9–10	4–5	5–7	7–8	8–9	absent	absent	solitary phialide	rough, in parallel rings	globose	0.18–0.24	present, lack visible rings
<i>P. ovalum</i>	no growth	12–14	no growth	no growth	10–11	10–11	7–8	13–14	8–10	9–11	absent	absent	solitary phialide	smooth	ellipsoidal	n.a.	present, lack visible rings
<i>P. porphyreum</i>	no growth	11–12	2–3	no growth	10–11	10–11	9–10	10–12	11–12	12–13	present	absent	solitary phialide	rough	globose	0.16–0.23	long, visible rings present
<i>P. riverlandense</i>	4–7	9–14	8–9	3–4	10–14	10–14	6–8	10–15	10–12	10–12	present	absent	solitary phialide	rough	globose	0.19–0.35	long, lack visible rings
<i>P. tubakianum</i>	no growth	5–6	2–4	no growth	8–9	8–9	4–5	7–8	5–7	9–10	absent	absent	solitary phialide	rough	globose	0.17–0.28	present, lack visible rings
<i>P. variatense</i>	no growth	10–12	8–9	2–3	9–10	8–9	4–5	7–8	9–10	7–8	absent	absent	solitary phialide	finely rough	broadly ellipsoidal	0.08–0.14	short, lack visible rings
<i>P. williamtense</i>	no growth	no growth	no growth	no growth	7–8	5–6	4–5	7–8	6–7	7–8	absent	absent	solitary phialide	rough	globose	0.23–0.42	long, lack visible rings
<i>P. wisconsinense</i>	no growth	3–5	no growth, sometimes 2	no growth	7–10	9–11	3–6	7–11	9–10	8–10	absent	absent	solitary phialide	rough	globose	0.24–0.38	short, lack visible rings
<i>P. wolleimicola</i>	no growth	6–7	4–5	no growth	7–8	9–10	4–5	8–9	7–9	9–11	present	absent	solitary phialide	rough	globose	0.22–0.41	long, lack visible rings

DNA extraction, sequencing, and phylogenetic analysis

DNA extractions were made from colonies grown on OA for 8–10 d using the Ultraclean™ Microbial DNA isolation Kit (MoBio, Solana Beach, USA). Subsequent PCR amplification and sequencing of the ITS, *BenA*, *CaM* and *RPB2* gene regions, were done using methods described by Visagie et al. (2014b). Sequence contigs were assembled in DNASTAR® SeqMan v. 9.0.4. Newly generated sequences were deposited in GenBank (Table 1). Sequences were aligned in MAFFT v. 7.164 (Kato & Standley 2013) with the L-INS-I option. For the multigene phylogeny, the datasets were concatenated in SeaView v. 4.4.1 (Gouy et al. 2010). Datasets were analysed using both Maximum Likelihood (ML) and Bayesian tree inference (BI). Trees and alignments were uploaded to TreeBASE (www.treebase.org) with submission ID 16300.

ML analyses were performed using MEGA v. 6.06 (Tamura et al. 2013). The most suitable substitution model for each dataset was selected using the model-test within MEGA, based on the lowest Bayesian information criterion (BIC) value. An initial tree was calculated with the Bio-Neighbour-Joining (BioNJ) option, with the subsequent Heuristic search done with Nearest-Neighbour-Interchange (NNI). A bootstrap analysis with 1 000 replicates was used for calculating node support.

For BI analyses, the most suitable substitution model for each dataset, according to the lowest Akaike information criterion (AIC) value, was selected using MrModeltest v. 2.3 (Nylander et al. 2004). Each analysis was run in MrBayes v. 3.2.1 (Ronquist & Huelsenbeck 2003), with two sets of four chains until the standard deviation of split frequencies reached 0.01. Sample frequency was set at 100 and 25 % of trees removed as burn in. ML phylograms were used for representing phylogenetic results, with both the bootstrap (bs) and posterior probability (pp) values, higher than 80 % bs and/or 0.95 pp given above thickened branch nodes.

Morphology

Standardised techniques for culturing conditions were used as described by Visagie et al. (2014b). Colony characters were recorded from strains grown on CYA (Czapek yeast autolysate agar), MEA (malt extract agar, Oxoid), YES (yeast extract sucrose agar), DG18 (dichloran 18 % glycerol agar), CYAS (CYA supplemented with 5 % NaCl), OA (oatmeal agar), SNA (Spezieller Nährstoffarmer agar) and CREA (creatine sucrose agar). All plates were incubated at 25 °C, with additional CYA plates incubated at 30 and 37 °C. Colour names and codes used for descriptions refer to the Methuen Handbook of Colour (Kornerup & Wanscher 1967).

Microscopic characters were recorded using both light-microscopy and low-temperature scanning electron microscopy (SEM). Microscopic preparations were made from 7–14-d-old cultures grown on OA, with 60 % lactic acid used as mounting fluid. Examinations were done using a Zeiss Axioskop 2 Plus light-microscope fitted with a Nikon microscope camera and pictures captured and analysed using Nikon NIS-elements D v. 4.0 software. In species descriptions, the average measurements and standard deviations are provided in brackets. Two-week-old colonies grown on OA were used for SEM observations. The technique used for SEM is similar to that described in Visagie et al. (2013). Photo plates were prepared using Adobe® Photoshop® Creative Suite v. 6. Photomicrographs were modified for aesthetic purposes using the healing brush tool, without altering areas of scientific significance.

RESULTS

Phylogeny

Phylogenies of all strains were prepared for the ITS region, and the partial *BenA*, *CaM* and *RPB2* gene regions, and compared using the Genealogical Concordance Phylogenetic Species Recognition (GCPSR) concept (Taylor et al. 2000; Fig. 1). A multigene phylogeny of concatenated ITS, *BenA*, *CaM* and *RPB2* sequences (Fig. 2) was prepared to evaluate the overall phylogenetic structure and the strength of support for nodes in sect. *Torulomyces*. The lengths of aligned datasets and substitution models selected for the analyses are provided in Table 2. Tree topologies did not differ between ML and BI analyses and ML trees are used here for presenting results, with BI pp values marked on relevant branches.

Phylogenies of *BenA*, *CaM* and *RPB2* were congruent and variable enough to distinguish all species (Fig. 1b–d). The multigene phylogeny (Fig. 2) indicated the presence of 14 species in the main clade of sect. *Torulomyces*, with *P. cryptum* and *P. lassenii* distantly related. In the main clade, GCPSR confirmed the distinct nature of the previously described species *P. lagena* and *P. porphyreum*. A further 12 clades were recognized, and were congruent across the different phylogenies. These could be distinguished morphologically and are described as new species in the Taxonomy section below.

From the ITS phylogeny (Fig. 1a) it is apparent that the formal barcode for fungi (Schoch et al. 2012) does not work well for species identification in the section, with *P. austriicola*, *P. wisconsinense*, *P. wollemiicola* and some strains of *P. riverlandense* sharing identical ITS sequences. The three alternative genes used here can distinguish between all species in the section, but we follow Visagie et al. (2014b) in proposing the use of *BenA* as an identification marker.

Some degree of variation in gene sequences occurred within *P. austriicola* and *P. riverlandense*, the two species recovered from ecological surveys of the fynbos and *Protea repens* infructescences in South Africa. In the ITS tree, *P. riverlandense* strains had five different genotypes resolved into three main clades (red blocks in Fig. 1a). This conflicts with the other gene trees, where the same clades also occur, but are united in a monophyletic clade that is supported in the *CaM* (Fig. 1c) and *RPB2* (Fig. 1d) phylogenies; all three clades are well-supported in the multigene phylogeny (Fig. 2). In *P. austriicola*, there is one clade in the ITS tree, two in *RPB2* and three in the *CaM* and *BenA* trees. These clades do not conflict, although one of them includes only a single strain (CBS 135903). Although it would be feasible to apply GCPSR and designate these clades within *P. austriicola* and *P. riverlandense* as phylogenetic species, the lack of diagnostic phenotypic characters and their geographical and ecological coherence argues against this. Increased sampling of strains and genes should indicate whether our conservative approach is correct.

ITS sequences of *P. lassenii* and *P. cryptum* show that these species might not belong to sect. *Torulomyces* as defined here. They were basal to the section in the other phylogenies. Similarly, *P. laeve* and *P. ovatum* are shown to belong to sect. *Exilicaulis* with *P. dimorphosporum* as their closest relative.

Morphology

Strains used for phylogenetic analyses were grown for morphological analysis under standardised conditions, and the clades corresponded well with morphological differences. Colony growth rates on different media incubated at different temperatures were the most useful characters for identification. Conidiophores were essentially identical among the species of the section. The exceptions were *P. lassenii*, which produced mono- to biverticillate conidiophores, and *P. cryptum*, which

produced mostly monoverticillate conidiophores, with a small proportion consisting of solitary phialides. Conidial shape varies from globose to ellipsoidal and is taxonomically informative. Conidial ornamentation was informative for species identification, but SEM observation is required for this. *Penicillium cryptum*, *P. lassenii*, *P. laeve* and *P. ovatum*, the species excluded from the section based on molecular results, produces smooth-walled conidia, not observed for other species in sect. *Torulomyces* s.str.

Morphological differences between the species described below are summarised in Table 3.

TAXONOMY

The following species are accepted in *Penicillium* sect. *Torulomyces*. *Penicillium ovatum* and *P. laeve*, previously classified in the section, are classified in *Penicillium* sect. *Exilicaulis*. Art. 9.3. of the Code (McNeill et al. 2012) explicitly includes published illustrations among 'original material', which precludes neotypification (Art. 9.7). Therefore the erroneous neotypification of *Torulomyces lagena* by Stolk & Samson (1983) is corrected by the designation of one of Delitsch's (1943) original figures as lectotype and designation of the 'neotype' strain CBS 185.65 as the epitype. *Penicillium parviverrucosum* is considered a dubious species.

Penicillium aeris Visagie & Samson, *sp. nov.* — MycoBank MB808262; Fig. 3a–g, 4a–c

ITS Barcode. KF303654 (alternative markers: *BenA* = KF303614; *CaM* = KF303627; *RPB2* = KF303681).

Etymology. Latin, meaning of the air, in reference to the type strain isolated from an air sample.

Type specimen. GERMANY, indoor air, 2012, coll. I. Ilyashenko (CBS H-21608 holotype, culture ex-type CBS 135897 = DTO 207D4).

Diagnosis — Growth present on CYA at 37 °C, no growth on CREA, conidia rough-walled, globose, 2–2.5 µm diam.

Description — Colony diam, 7 d, in mm: CYA 11–12; CYA 30 °C 9–10; CYA 37 °C 4–5; MEA 6–7; YES 10–11; OA 9–10; DG18 11–12; CYAS 6–7; SNA 8–9; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse pale orange (5A3), reverse pale orange (5A3); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse greenish white (25A2), reverse brownish orange (5C5); OA obverse greyish green (26E5); DG18 obverse yellowish white (3A2), reverse yellowish white (3A2); CYAS obverse white, reverse white; SNA obverse greenish grey (26B2), reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4–13.5 × 1.5–2 µm; *phialides* ampulliform, 3.5–6.5 × 2–3 µm (4.9 ± 0.5 × 2.4 ± 0.2); *conidia* rough to spiny, globose, 2–2.5 × 2–2.5 µm (2.1 ± 0.2 × 2.1 ± 0.2, n = 34), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives long without visible rings, tubercles 0.21–0.4 µm diam (0.31 ± 0.05).

Penicillium austriicola Visagie & K. Jacobs, *sp. nov.* — MycoBank MB805184; Fig. 3h–n, 4d–f

ITS Barcode. JX091466 (alternative markers: *BenA* = JX091579; *CaM* = JX141600; *RPB2* = KF303705).

Etymology. Latin, meaning resident from the south, in reference to the species isolated close to the most southern tip of Africa at Struisbaai.

Type specimen. SOUTH AFRICA, Struisbaai, from mite inside *Protea repens* infructescence, Aug. 2009, coll. C.M. Visagie (CBS H-21605 holotype, culture ex-type CBS 135900 = CV 1842 = DTO 183E6 = DAOMC 241066).




Fig. 3 Morphology of species characterised in this study. a, h, o. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–g, i–n, p–v. conidiophores and conidia produced on OA. — Scale bar: v = 10 μm, applies to all microscope pictures.




Fig. 4 SEM micrographs showing characters of monophaialidic *Penicillium* species. a, d, g, j, m. Overview of conidiogenous areas; b, e, h, k, n. phialides; c, f, i, l, o. conidia. — Scale bars: a, d, g, j, m = 10 µm; b, e, h, k, n = 5 µm; c, f, i, l, o = 2 µm.

Diagnosis — No growth on CYA at 37 °C, growth on CYA at 30 °C and CREA, colonies typically produces brown soluble pigments, conidia rough-walled, globose, 1.5–2.5 µm diam.

Description — Colony diam, 7 d, in mm: CYA 10–13; CYA 30 °C 10–11; CYA 37 °C no growth; MEA 4–8; YES 10–11; OA 9–10; DG18 9–11; CYAS 8–10; SNA 9–11; CREA 3–4.

Colonies at 25 °C, 7 d: CYA obverse greyish to dull green (25B3–26D3), reverse olive to dark brown (4E3–6F8), brown soluble pigment produced in some isolates; MEA obverse greyish green (25B3–26B3), reverse light to dark brown (6D6–F8), brown soluble pigment produced in some isolates; YES obverse orange white (5A2) when not sporulating, greyish green (25C4) when sporulating, reverse greyish yellow (4B4); OA

obverse greyish green (25D5–E5); DG18 obverse greenish white to pale green (25A2–A3), reverse dull green (29D3); CYAS obverse greenish grey (25B2), reverse brown (6E6); SNA obverse dull green (29D4), reverse greenish grey (29B2); CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4–11 × 1–1.5 µm; *phialides* ampulliform, 5–8 × 2–3 µm (6.1 ± 0.6 × 2.7 ± 0.2); *conidia* rough-walled, globose, 1.5–2.5 × 1.5–2.5 µm (2.1 ± 0.1 × 2.1 ± 0.1, n = 25), average width/length = 0.98; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives long without visible rings, tubercles 0.19–0.30 µm diam (0.23 ± 0.03).

Additional material examined. SOUTH AFRICA, Struisbaai, from mite inside *Protea repens* infructescence, Aug. 2009, coll. C.M. Visagie, cultures CBS 135904 (CV 1808 = DTO 183D8 = DAOMC 241065), CBS 135899 (CV 1811 = DTO 186D8), CBS 135902 (CV 1943 = DTO 183F8 = DAOMC 241067), CBS 135903 (CV 1954 = DTO 183F9), CBS 136253 (CV 1968 = DTO 186G3); Struisbaai, bract from *Protea repens* infructescence, Aug. 2009, coll. C.M. Visagie, culture CBS 135901 (CV 1902 = DTO 183F5).

Penicillium cantabricum Visagie & Samson, *sp. nov.* — MycoBank MB808263; Fig. 3o–v, 4g–i

ITS Barcode. KF303655 (alternative markers: *BenA* = KF303615; *CaM* = KF303646; *RPB2* = KF303682).

Etymology. Latin, named after Cantabria, from where the type was isolated.

Type specimen. SPAIN, Cantabria, Reinosa, from soil, June 1996, isol. A.M. Stchigel (CBS H-21612 holotype, culture ex-type CBS 120415 = DTO 7619 = FMR 9121).

Diagnosis — Growth present on CYA at 37 °C, very weak acid produced within colony periphery on CREA, conidia finely rough-walled, globose, 2–2.5 µm.

Description — Colony diam, 7 d, in mm: CYA 13–14; CYA 30 °C 12–13; CYA 37 °C 3–5; MEA 6–7; YES 9–10; OA 10–11; DG18 8–9; CYAS 3–4; SNA 7–8; CREA 3–4.

Colonies at 25 °C, 7 d: CYA obverse pale orange (6A3), reverse pale orange (5A3), pinkish mycelia produced; MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse orange white (5A2), reverse light orange (5A4); OA obverse greyish green (25B3); DG18 obverse yellowish white (3A2), reverse yellowish white (3A2); CYAS obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid very weakly produced within colony periphery.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4.5–24 × 1.5–2 µm; *phialides* ampulliform, 3.5–5.5 × 2.5–3 µm (4.5 ± 0.5 × 2.6 ± 0.2); *conidia* finely rough-walled, globose, 2–2.5 × 2–2.5 µm (2.2 ± 0.1 × 2.2 ± 0.1, n = 41), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives present without visible rings, tubercles 0.21–0.33 µm diam (0.26 ± 0.03).

Penicillium catalanicum Visagie & Samson, *sp. nov.* — MycoBank MB808265; Fig. 4j–l, 5a–g

ITS Barcode. KF303650 (alternative markers: *BenA* = KF303609; *CaM* = KF303644; *RPB2* = KF303683).

Etymology. Latin, named after Catalonia, from where the type was isolated.

Type specimen. SPAIN, Catalonia, Montseny Natural Park, from soil, unknown date, isol. A.M. Stchigel & M. Caldich (CBS H-21610 holotype, culture ex-type CBS 110532 = DTO 78H5).

Diagnosis — No growth on CYA at 30 °C, growth on CYAS, conidia finely rough-walled, with tubercles forming rings around

conidia under SEM, subglobose to broadly ellipsoidal, 2–2.5 × 2–2.5 µm.

Description — Colony diam, 7 d, in mm: CYA 6–8; CYA 30 °C no growth; CYA 37 °C no growth; MEA 4–5; YES 5–6; OA 6–7; DG18 7–8; CYAS 2–3; SNA 6–7; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse greenish white (25A2), reverse yellowish white (4A2); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse turquoise white to greenish white (24A2–25A2), reverse olive brown (4D5); OA obverse greyish green (26C4); DG18 obverse yellowish white (3A2), reverse yellowish white (3A2); CYAS obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4.5–11 × 1.5–2 µm; *phialides* ampulliform, 5–7 × 2–3 µm (5.9 ± 0.6 × 2.4 ± 0.2); *conidia* finely rough, subglobose to broadly ellipsoidal, 2–2.5 × 2–2.5 µm (2.16 ± 0.1 × 2.11 ± 0.1, n = 41), average width/length = 0.98; *sclerotia* not produced.

SEM observations: Conidia more ellipsoidal than when observed under light microscope, finely rough-walled with very small tubercles present, sometimes in ring like pattern, connections short and not well defined, tubercles 0.12–0.18 µm diam (0.14 ± 0.02).

Penicillium cryptum Goch., Mycotaxon 26: 349. 1986 — MycoBank MB103648; Fig. 4m–o, 5h–p

Basionym. *Eupenicillium cryptum* Goch., Mycotaxon 26: 349. 1986.

ITS Barcode. KF303647 (alternative markers: *BenA* = KF303608; *CaM* = KF303628; *RPB2* = JN121478).

Type specimen. USA, New York, Long Island, from soil in *Quercus-Betula* forest, Oct. 1983, isol. S.E. Gochenaour (NY 769 holotype, culture ex-type CBS 271.89 = DTO 122C9 = ATCC 60138 = IMI 296794 = NRRL 13460).

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, conidiophores typically monophialidic to monoverticillate, conidia smooth-walled, broadly ellipsoidal, 2–2.5 × 1.5–2 µm.

Description — Colony diam, 7 d, in mm: CYA 6–7; CYA 30 °C 6–9; CYA 37 °C no growth; MEA 6–7; YES 5–6; OA 5–6; DG18 1–2; CYAS no growth; SNA 5–6; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse greenish white (25A2), reverse pale; MEA obverse greenish white (25A2), reverse yellowish brown (5D6); YES obverse greenish white (25A2), reverse yellowish white (3A2); OA obverse greenish white (25A2); DG18 small white to pale; SNA obverse white to light greyish, reverse white to light greyish.

Conidiophores monoverticillate with at most 3 phialides, phialides often borne directly on hyphae, conidiophores sometimes as solitary phialides; *stipes* smooth-walled, 5–30 × 1–2.5 µm when monoverticillate, 1–11 × 1–2.5 µm when monophialidic; *phialides* ampulliform, 5–9 × 2.5–3.5 µm (6.8 ± 0.9 × 2.9 ± 0.2); *conidia* smooth-walled, broadly ellipsoidal, 2–2.5 × 1.5–2 µm (2.3 ± 0.1 × 1.9 ± 0.1, n = 35), average width/length = 0.83; *sclerotia* not produced.

SEM observations: Conidia broadly ellipsoidal, smooth-walled, connectives present, but rather obscure.

Penicillium laeve (K. Ando & Manoch) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 — MycoBank MB561960; Fig. 5q–w, 6a–c

Basionym. *Torulomyces laevis* K. Ando & Manoch, Mycoscience 39: 317. 1998.

ITS Barcode. KF667369 (alternative markers: *BenA* = KF667365; *CaM* = KF667367; *RPB2* = KF667371).




Fig. 5 Morphology of species characterised in this study. a, h, q. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–g, i–p, r–w. conidiophores, conidia, asci and ascospores (in w) produced on OA. — Scale bar: w = 10 µm, applies to all microscope pictures.

Type specimen. THAILAND, from soil, Oct. 1992, isol. K. Ando (TNS-F-238517 holotype, culture ex-type CBS 136665 = DTO 270G8 = KY 12727 = NBRC 109724).

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, conidia smooth-walled, globose, 2.5–3 µm diam. Ascomata inconspicuous, asci 3–8 µm diam, ascospores smooth-walled, globose, 1.5–2 µm diam.

Description — Colony diam, 7 d, in mm: CYA 8–9; CYA 30 °C 4–5; CYA 37 °C no growth; MEA microcolonies; YES 8–9; OA

7–8; DG18 5–7; CYAS no growth; SNA 7–8; CREA no growth. Colonies at 25 °C, 7 d: CYA obverse white, reverse greyish yellow (3C3); MEA obverse white, reverse yellowish brown (5D6); YES obverse greenish grey (25B2), reverse brownish orange (5C4); OA obverse greyish white (1B1), reverse greyish yellow (3B3); SNA obverse white to light greyish, reverse white; CREA acid not produced. *Conidiophores* as solitary phialides; *stipes* smooth-walled, 2–6 × 1–1.5 µm; *phialides* ampulliform, 4–6 × 2–3 µm (5.2 ± 0.6




Fig. 6 SEM micrographs showing characters of monophaialidic *Penicillium* species. a, d, g, j, m. Overview of conidiogenous areas; b, e, h, k, n. phialides; c, f, i, l, o. conidia. — Scale bars: a, d, g, j, m = 10 µm; b, e, h, k, n = 5 µm; c, f, i, l, o = 2 µm.

× 2.4 ± 0.1); *conidia* smooth-walled, globose, 2.5–3 × 2.5–3 µm (2.6 ± 0.1 × 2.6 ± 0.1, n = 22), average width/length = 0.98; sexual morph produced, *ascomata* inconspicuous, asci 3–8 µm with smooth-walled, globose *ascospores* 1.5–2 × 1.5–2 µm. SEM observations: Conidia globose, smooth-walled, long connectives present without visible rings.

Penicillium lagena (Delitsch) Stolk & Samson, Stud. Mycol. 23: 100. 1983 — MycoBank MB109162; Fig. 6d–f, 7a–h

Basionym. *Torulomyces lagena* Delitsch, Systematik der Schimmelpilze, Neudamm: 9. 1943.

≡ *Eupenicillium limoneum* Goch. & Zlattner, Stud. Mycol. 23: 100. 1983.

ITS Barcode. KF303665 (alternative markers: *BenA* = KF303619; *CaM* = KF303634; *RPB2* = JN121450).

Lectotype. Fig. 233 (Delitsch, Ergebnisse der theoretischen und angewandten Mikrobiologie: Band I: Systematik der Schimmelpilze. J. Neumann, Neudamm, Germany, Tafel 30, 1943, designated here, MBT203020).

Epitype. CANADA, Guelph, from bog soil under Pacific red cedar (*Thuja plicata*), 1961, isol. G.L. Barron (CBS 185.65 epitype (metabolically inactive), MBT203019, culture ex-epitype CBS 185.65 = MUCL 8221 = JCM10149), designated here.

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, colonies on CYA 10–13 mm, conidia rough-walled, under SEM with small tubercles, connectives lacking visible rings, globose, 2–2.5 µm diam.

Description — Colony diam, 7 d, in mm: CYA 10–13; CYA 30 °C 8–10; CYA 37 °C no growth; MEA 8–10; YES 11–13; OA 8–12; DG18 11–13; CYAS no growth to 3; SNA 10–13; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse greenish grey (25B2), reverse dark brown (6F8), some isolates pale, brown soluble pigment produced in some isolates; MEA obverse greenish white (25A2), reverse yellowish brown (5D6), brown soluble pigment produced in some isolates; YES obverse yellowish white (3A2) when not sporulating, otherwise greyish green (25B3), reverse brownish orange (5C4) in non-sporulating cultures, otherwise olive (2D3); OA obverse greyish green (25C3–26C3), brown soluble pigment produced; DG18 obverse yellowish white (3A2) when not sporulating, otherwise (29C3), reverse brownish orange (5C5), brown soluble pigment produced; CYAS obverse white, reverse white; SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4–20 × 1–1.5 µm; *phialides* ampulliform, 4.5–7 × 2–2.5 µm (6.3 ± 0.7 × 2.2 ± 0.1); *conidia* rough-walled, globose, 2–2.5 × 2–2.5 µm (2.1 ± 0.2 × 2.1 ± 0.2, n = 27), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with pronounced large tubercles present, connectives long without easily visible rings, tubercles 0.17–0.29 µm diam (0.22 ± 0.04).

Penicillium lassenii Paden, Mycopathol. Mycol. Appl. 43: 266. 1971. — MycoBank MB319281; Fig. 7i–o

≡ *Eupenicillium lassenii* Paden, Mycopathol. Mycol. Appl. 43: 266. 1971.

ITS Barcode. KF303648 (alternative markers: *BenA* = KF303607; *CaM* = KF303629; *RPB2* = JN121481).

Type specimen. USA, California, from soil under conifers, May 1969, isol. J.W. Paden (JWP 69-26 (UVIC) holotype, culture ex-type CBS 277.70 = DTO 95D6 = NRRL 5272 = ATCC 22054 = FRR 858 = IMI 148395).

Diagnosis — No growth on CYA at 37 °C, growth on CYA at 30 °C and CREA, conidiophores typically mono- to biverticillate, conidia smooth-walled, broadly ellipsoidal, 2.5–3 × 2–3 µm, produces pale to brown *sclerotia*.

Description — Colony diam, 7 d, in mm: CYA 7–9; CYA 30 °C 4–5; CYA 37 °C no growth; MEA 9–10; YES 9–10; OA 5–6; DG18 10–11; CYAS 4–5; SNA 5–6; CREA 3–4.

Colonies at 25 °C, 7 d: CYA obverse dull green (27D3), reverse yellowish grey (2B2); MEA obverse greyish green (27C3), reverse light brown (6D5); YES obverse dull green (27D3), reverse yellowish grey (2B2); OA obverse dull green (27D3); DG18 obverse dull green (27D3), reverse yellowish grey (2B2); CYAS obverse white, reverse white; SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced. *Conidiophores* monoverticillate to sometimes biverticillate; *stipes* smooth-walled, 9–75 × 1.5–3.5 µm; branches 4–15 × 1.5–3.5 µm; *phialides* ampulliform, 6–7.5 × 2.5–3.5 µm (6.7 ± 0.6 × 2.9 ± 0.2); *conidia* smooth-walled, broadly ellipsoidal, 2.5–3 × 2–3 µm (2.6 ± 0.1 × 2.45 ± 0.1, n = 37), average width/length = 0.89; *sclerotia* pale to brownish, 105–235 × 90–170 µm.

Penicillium marthae-christenseniae Visagie & Samson, *sp. nov.* — MycoBank MB808267; Fig. 4g–i, 7p–w

ITS Barcode. KF303651 (alternative markers: *BenA* = KF303613; *CaM* = KF303645; *RPB2* = KF303711).

Etymology. Latin, named after Martha Christensen, who isolated and deposited the ex-type strain into the CBS collection.

Type specimen. USA, Wisconsin, unknown source and date, isol. *M. Christensen* (CBS H-21613 holotype, culture ex-type CBS 129213 = DTO 201B5).

Diagnosis — No growth on CYA at 37 °C, CREA or CYAS, growth on CYA at 30 °C, colonies on CYA 7–8 mm and SNA 11–12 mm, conidia rough-walled, with large well-defined tubercles (average 0.26 ± 0.03 µm) under SEM, subglobose, 2–2.5 × 2–2.5 µm.

Description — Colony diam, 7 d, in mm: CYA 7–8; CYA 30 °C 5–6; CYA 37 °C no growth; MEA 4–5; YES 7–8; OA 9–10; DG18 9–11; CYAS no growth; SNA 11–12; CREA no growth. Colonies at 25 °C, 7 d: CYA obverse white, reverse pale yellow (3B3); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse greyish green (25C4), reverse yellowish grey to olive brown (4B2–E4); OA obverse dull green (27D4); DG18 obverse very light greyish green, reverse greyish yellow (4B3); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 3–10 × 1–2 µm; *phialides* ampulliform, 4–6.5 × 2–3 µm (5.1 ± 0.6 × 2.4 ± 0.2); *conidia* rough-walled, subglobose, 2–2.5 × 2–2.5 µm (2.2 ± 0.1 × 2.1 ± 0.1, n = 34), average width/length = 0.96; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives present with visible rings, tubercles 0.22–0.32 µm diam (0.26 ± 0.03).

Penicillium oregonense Visagie, M. Chr. & Samson, *sp. nov.* — MycoBank MB808268; Fig. 6j–l, 8a–g

ITS Barcode. KF303668 (alternative markers: *BenA* = KF303623; *CMD* = KF303640; *RPB2* = KF303710).

Etymology. Latin, named after Oregon State, from where the type was isolated.

Type specimen. USA, Oregon, Willamette National Forest near Blue River, unknown source and date, isol. *M. Christensen* (CBS H-21607 holotype, culture ex-type CBS 129775 = DTO 208A5).

Diagnosis — No growth on CYA at 30 °C or CYAS, colonies on CYA at 25 °C 4–5 mm, conidia rough-walled with tubercles in rings under SEM, globose, 2–2.5 × 2–2.5 µm.




Fig. 7 Morphology of species characterised in this study. a, i, p. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–h, j–o, q–w. conidiophores, conidia and sclerotia (q) produced on OA. — Scale bar: w = 10 µm, applies to all microscope pictures.




Fig. 8 Morphology of species characterised in this study. a, h, o. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–g, i–n, p–u. conidiophores and conidia produced on OA. — Scale bar: u = 10 µm, applies to all microscope pictures.

Description — Colony diam, 7 d, in mm: CYA 4–5; CYA 30 °C no growth; CYA 37 °C no growth; MEA 4–5; YES 5–7; OA 9–10; DG18 8–9; CYAS no growth; SNA 7–8; CREA no growth. Colonies at 25 °C, 7 d: CYA obverse white, reverse pale yellow (3B3); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse yellowish white (3A2), reverse yellowish white (3A2); OA obverse dull green (28E4); DG18

obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, $2.5\text{--}14 \times 1.5\text{--}2 \mu\text{m}$; *phialides* ampulliform, $4.5\text{--}6 \times 2.5\text{--}3.5 \mu\text{m}$ ($5.5 \pm 0.4 \times 2.8 \pm 0.2$); *conidia* rough-walled, globose, $2\text{--}2.5 \times 2\text{--}2.5 \mu\text{m}$ ($2.15 \pm 0.1 \times 2.15 \pm 0.1$, $n = 32$), average width/length = 0.99; *sclerotia* not produced.




Fig. 9 SEM micrographs showing characters of monophaealidic *Penicillium* species. a, d, g, j, m. Overview of conidiogenous areas; b, e, h, k, n. phialides; c, f, i, l, o. conidia. — Scale bars: a, d, g, j, m = 10 μm ; b, e, h, k, n = 5 μm ; c, f, i, l, o = 2 μm .

SEM observations: Conidia rough-walled, with tubercles forming conspicuous parallel rings, connectives moderately long and well defined without visible rings, tubercles 0.18–0.24 µm diam (0.22 ± 0.04).

Penicillium ovatum (K. Ando & Nawawi) Houbraken & Samson, *Stud. Mycol.* 70: 48. 2011 — MycoBank MB561961; Fig. 6m–o, 8h–n

Basionym. *Torulomyces ovatus* K. Ando & Nawawi, *Mycoscience* 39: 317. 1998.

ITS Barcode. KF667370 (alternative markers: *BenA* = KF667366; *CaM* = KF667368; *RPB2* = KF667372).

Type specimen. MALAYSIA, Kuala Lumpur, from soil under *Pinus caribaea*, Nov. 1989, isol. K. Ando (TNS-F-238518 holotype, culture ex-type CBS 136664 = KY 12726 = DTO 270G7 = NBRC 109725).

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, conidia smooth-walled, ellipsoidal, 2–3 × 1.5–2 µm.

Description — Colony diam, 7 d, in mm: CYA 10–11; CYA 30 °C 12–14; CYA 37 °C no growth; MEA 7–8; YES 13–14; OA 10–11; DG18 9–11; CYAS no growth; SNA 8–10; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse greyish white (1B1), reverse brownish orange (5C4), colonies producing orange colours after prolonged incubation; MEA obverse greyish white (1B1), reverse yellowish brown (5D6); YES obverse greenish grey (29C2), reverse greyish yellow (4C4); OA obverse greyish white (1B1); DG18 obverse greenish grey (29C2), reverse greyish yellow (3C4); SNA obverse white to light greyish, reverse white; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 3.5–15 × 1–2 µm; *phialides* ampulliform, 4.5–7 × 2–3 µm (5.7 ± 0.5 × 2.5 ± 0.25); *conidia* smooth-walled, ellipsoidal, 2–3 × 1.5–2 µm (2.4 ± 0.2 × 1.8 ± 0.1, n = 27), average width/length = 0.76; *sclerotia* not produced.

SEM observations: Conidia ellipsoidal, smooth-walled, connectives present without visible rings.

Penicillium porphyreum Houbraken & Samson, *Stud. Mycol.* 70: 48. 2011 — MycoBank MB561959; Fig. 8o–u, 9a–c

≡ *Monocillium humicola* G.L. Barron var. *brunneum* M. Chr. & Backus, *Mycologia* 56: 498. 1964.

≡ *Torulomyces brunneus* (M. Chr. & Backus) K. Ando, *Mycoscience* 39: 314. 1998.

ITS Barcode. KF303666 (alternative markers: *BenA* = KF303621; *CaM* = KF303636; *RPB2* = KF303677).

Type specimen. USA, Wisconsin, from soil in community of *Pinus strobus*, July 1955, isol. M. Christensen (NY barcode 00985491 holotype, culture ex-type CBS 382.64 = ATCC 15571 = DTO 78G7 = KY 12723 = WSF 9-C).

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, colonies on CYA 10–11 mm, conidia rough-walled with small tubercles (average 0.19 ± 0.02 µm), connectives have two clearly visible rings.

Description — Colony diam, 7 d, in mm: CYA 10–11; CYA 30 °C 11–12; CYA 37 °C no growth; MEA 9–10; YES 10–12; OA 10–11; DG18 12–13; CYAS 2–3; SNA 11–12; CREA no growth. Colonies at 25 °C, 7 d: CYA obverse yellowish to orange white (4A2–5A2), reverse orange white (5A2), brown soluble pigment typical; MEA obverse yellowish to orange white (4A2–5A2), reverse light brown (6D7), brown soluble pigment typical; YES obverse turquoise white to greenish white (24A2–25A2), reverse brownish orange (5C4); OA obverse greyish green (25C3–26C3), brownish olive soluble pigment produced; DG18

obverse yellowish grey (3B2), reverse greyish orange (5C4); CYAS obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 5–17 × 1–1.5 µm; *phialides* ampulliform, 4.5–7.5 × 2–2.5 µm (6 ± 0.8 × 2.3 ± 0.2); *conidia* rough-walled, globose, 2–2.5 × 2–2.5 µm (2.16 ± 0.1 × 2.16 ± 0.1, n = 43), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with pronounced large to moderately sized tubercles present, connectives long, consistently having two visible rings, tubercles 0.16–0.23 µm diam (0.19 ± 0.02).

Penicillium riverlandense Visagie & K. Jacobs, *sp. nov.* — MycoBank MB808269; Fig. 9d–f, 10a–g

ITS Barcode. JX091457 (alternative markers: *BenA* = JX091580; *CaM* = JX141593; *RPB2* = KF303685).

Etymology. Latin, named after Riverlands Nature Reserve, in reference to the location of the type origin.

Type specimen. SOUTH AFRICA, Malmesbury, Riverlands Nature Reserve, from sandveld Fynbos soil, July 2009, isol. C.M. Visagie (CBS H-21606 holotype, culture ex-type CBS 135896 = CV 0979 = DTO 182F6 = DAOMC 241060).

Diagnosis — Growth observed on CYA at 37 °C, colonies producing brown soluble pigment, conidia rough-walled, globose, mostly smaller than 2 µm diam.

Description — Colony diam, 7 d, in mm: CYA 10–14; CYA 30 °C 9–14; CYA 37 °C 4–7; MEA 6–8; YES 10–15; OA 10–14; DG18 10–12; CYAS 8–9; SNA 10–12; CREA 3–4.

Colonies at 25 °C, 7 d: CYA obverse greyish green (25B2), reverse olive to dark brown (5E5–6F8), brown soluble pigment produced in all isolates; MEA obverse light to greyish green (25A5–C5), reverse dark brown (6F8), brown soluble pigment produced in all isolates; YES obverse greyish green (25B4–26B4), sporulation sometimes absent, then pale to white colonies, brown soluble pigment produced, reverse brownish orange to brown (5C5–5F8); OA obverse dull green (25D4), brownish olive soluble pigment produced; DG18 obverse white to pale yellow (3A3), reverse yellowish white to dull yellow (3A2–B4); CYAS obverse greenish grey (25B2), reverse brown (6E6); SNA obverse dull green (29D4), reverse greenish grey (29B2); CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 3.5–10 × 1–2 µm; *phialides* ampulliform, 4–8 × 2–3 µm (5.5 ± 0.9 × 2.6 ± 0.2); *conidia* rough-walled, globose, 1.5–2 × 1.5–2 µm (1.8 ± 0.1 × 1.8 ± 0.1, n = 35), average width/length = 0.98; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives long without visible rings, tubercles 0.19–0.35 µm diam (0.27 ± 0.04).

Additional material examined. SOUTH AFRICA, Malmesbury, Riverlands Nature Reserve, sandveld Fynbos soil, July 2009, isol. C.M. Visagie, cultures CBS 135894 = CV 0933 = DTO 185F9, CBS 135895 = CV 0959 = DTO 182E7 = DAOMC 241059, CBS 135887 = CV 2815 = DTO 184A1, CBS 135888 = CV 2819 = DTO 184A4, CBS 135889 = CV 2822 = DTO 184A6, CBS 135890 = CV 2831 = DTO 184B4, CBS 135891 = CV 2833 = DTO 184B6, CBS 135892 = CV 2839 = DTO 184C2, CBS 135893 = CV 2849 = DTO 184C6; Riverlands Nature Reserve, bract from *Protea repens* infructescence, July 2009, isol. C.M. Visagie, cultures CBS 135883 = CV 1360 = DTO 183A4 = DAOMC 241061, CBS 135884 = CV 1448 = DTO 183A9 = DAOMC 241063, CBS 135885 = CV 1450 = DTO 183B1 = DAOMC 241064, CBS 135886 = CV 1583 = DTO 186B8; Struisbaai, bract from *Protea repens* infructescence, Aug. 2009, coll. C.M. Visagie, culture DTO 183E8 = CV 1851.




Fig. 10 Morphology of species characterised in this study. a, h, o. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–g, i–n, p–u. conidiophores and conidia produced on OA. — Scale bar: u = 10 µm, applies to all microscope pictures.

Penicillium tubakianum Visagie & Samson, *sp. nov.* — MycoBank MB808270; Fig. 9g–i, 10h–n

ITS Barcode. KF303652 (alternative markers: *BenA* = KF303611; *CaM* = KF303637; *RPB2* = KF303712).

Etymology. Latin, name after K. Tubaki, who isolated the ex-type strain of the species.

Type specimen. NEW ZEALAND, from dead bark of *Cyathea*, Dec. 1963, isol. K. Tubaki (CBS H-21604 holotype, culture ex-type CBS 287.66 = DTO 138D9 = MUCL 8519 = IFO 8315).

Diagnosis — No growth on CYA at 37 °C, CREA, growth on CYA at 30 °C, colonies on CYA 8–9 mm, CYAS 2–4 mm and SNA 5–7 mm, conidia rough-walled, globose, 2–2.5 µm diam, sclerotia not produced.

Description — Colony diam, 7 d, in mm: CYA 8–9; CYA 30 °C 5–6; CYA 37 °C no growth; MEA 4–5; YES 7–8; OA 8–9; DG18 9–10; CYAS 2–4; SNA 5–7; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse yellowish to orange white (4A2–5A2), reverse orange white (5A2); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse pale orange (5A3), reverse pale orange (5A3); OA obverse greyish green (26C4); DG18 obverse yellowish white (3A2), reverse yellowish white (3A2); CYAS obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 2.5–10 × 1–1.5 µm; *phialides* ampulliform, 3.5–6.5 × 2–3 µm (4.6 ± 0.2 × 2.5 ± 0.2); *conidia* rough-walled, globose, 2–2.5 × 2–2.5 µm (2.1 ± 0.1 × 2.1 ± 0.1, n = 34), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with tubercles present, connectives present without visible rings, tubercles 0.17–0.28 µm diam (0.23 ± 0.03).

Penicillium variratense Visagie & Samson, *sp. nov.* — MycoBank MB808271; Fig. 9j–l, 10o–u

ITS Barcode. KF303649 (alternative markers: *BenA* = KF303610; *CaM* = KF303630; *RPB2* = KF303675).

Etymology. Latin, named after Varirata National Park in Papua New Guinea, from where the type was isolated.

Type specimen. PAPUA NEW GUINEA, Varirata National Park near Port Moresby, from humus soil, Dec. 1996, isol. A. Aptroot & A. van Iperen (CBS H-21611 holotype, culture ex-type CBS 337.97 = DTO 137C8).

Diagnosis — No growth on CYA at 37 °C, growth on CYA at 30 °C and CREA, conidia smooth-walled by light microscope, finely roughened under SEM, broadly ellipsoidal, 2–3 × 2–2.5 µm.

Description — Colony diam, 7 d, in mm: CYA 9–10; CYA 30 °C 10–12; CYA 37 °C no growth; MEA 4–5; YES 7–8; OA 8–9; DG18 7–8; CYAS 8–9; SNA 9–10; CREA 2–3.

Colonies at 25 °C, 7 d: CYA obverse greyish green (25B4), reverse greyish yellow to greyish orange (4B4–5B4); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse greenish white (27A2), reverse brownish orange (5C5); OA obverse dark green (27F6); DG18 obverse pale yellow (1A3), reverse light yellow (3A4); CYAS obverse greyish green (25B4), reverse greyish yellow (3B5); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 3.5–11.5 × 1–2 µm; *phialides* ampulliform, 4.5–7 × 2–3.5 µm (5.7 ± 0.6 × 2.6 ± 0.2); *conidia* smooth-walled, broadly ellipsoidal, 2–3 × 2–2.5 µm (2.5 ± 0.2 × 2.3 ± 0.1, n = 30), average width/length = 0.91; *sclerotia* not produced.

SEM observations: Conidia very finely rough-walled, with very small tubercles present, connectives short, not well defined, tubercles 0.08–0.14 µm diam (0.11 ± 0.02).

Penicillium williamettense Visagie, M. Chr. & Samson, *sp. nov.* — MycoBank MB808272; Fig. 9m, n, 11a–h

ITS Barcode. KF303667 (alternative markers: *BenA* = KF303622; *CaM* = KF303639; *RPB2* = KF303709).

Etymology. Latin, named after Williamette National Forest in Oregon, from where the type was isolated.

Type specimen. USA, Oregon, Williamette National Forest near Blue River, unknown source and date, isol. M. Christensen (CBS H-21609 holotype, culture ex-type CBS 129774 = DTO 208A4).

Diagnosis — No growth on CYA at 30 °C or CYAS, colonies on CYA at 25 °C 7–8 mm, conidia rough-walled with big tubercles, visible with SEM, globose, 2–2.5 µm diam.

Description — Colony diam, 7 d, in mm: CYA 7–8; CYA 30 °C no growth; CYA 37 °C no growth; MEA 4–5; YES 7–8; OA 5–6; DG18 7–8; CYAS no growth; SNA 6–7; CREA no growth. Colonies at 25 °C, 7 d: CYA obverse white, reverse pale yellow (3B3); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse yellowish white (3A2), reverse yellowish white (3A2); OA obverse dull green (28E4); DG18 obverse yellowish white (3A2), reverse yellowish white (3A2); SNA obverse white to light greyish, reverse white to light greyish; CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 5–16 × 1–1.5 µm; *phialides* ampulliform, 4–6.5 × 2–3 µm (5.4 ± 0.5 × 2.6 ± 0.2); *conidia* rough-walled, globose, 2–2.5 × 2–2.5 µm (2.2 ± 0.2 × 2.2 ± 0.2, n = 25), average width/length = 0.99; *sclerotia* not produced.

SEM observations: Conidia rough-walled with very big tubercles present, connectives long with rings barely visible, tubercles 0.23–0.42 µm diam (0.33 ± 0.04).

Penicillium wisconsinense Visagie, M. Chr. & Samson, *sp. nov.* — MycoBank MB808273; Fig. 11i–o, 12a–c

ITS Barcode. KF303670 (alternative markers: *BenA* = KF303624; *CaM* = KF303641; *RPB2* = KF303706).

Etymology. Latin, named after Wisconsin, from where the type was isolated.

Type specimen. USA, Wisconsin, unknown source and date, isol. M. Christensen (CBS H-21614 holotype, culture ex-type CBS 128279 = DTO 198H7 = WSF 3132).

Diagnosis — No growth on CYA at 37 °C or CREA, growth on CYA at 30 °C, colonies on CYA 7–10 mm, CYAS 0–2 mm, SNA 9–10 mm, conidia rough-walled with very big tubercles under SEM, globose, 2–2.5 × 2–2.5 µm.

Description — Colony diam, 7 d, in mm: CYA 7–10; CYA 30 °C 3–5; CYA 37 °C no growth; MEA 3–6; YES 7–11; OA 9–11; DG18 8–10; CYAS no growth, sometimes 2; SNA 9–10; CREA no growth.

Colonies at 25 °C, 7 d: CYA obverse yellowish to orange white (4A2–5A2), reverse orange white (5A2), sometimes greyish brown (5D3); MEA obverse orange white (5A2), reverse yellowish brown (5D6); YES obverse greyish turquoise to greyish green (24B3–25B3), reverse olive to olive brown (3D5–4D5); OA obverse dull green (28E4); DG18 obverse greyish turquoise to greyish green (24B3–25B3), reverse greyish green (28C3), some greyish yellow (4C4); CYAS obverse pale when growth, reverse pale when growth; SNA obverse dull green (29D4), reverse greenish grey (29B2); CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 5–45 × 1–2 µm; *phialides* ampulliform, 4.5–7.5 × 2.5–3 µm (5.7 ±




Fig. 11 Morphology of species characterised in this study. a, i, p. Colony morphology from left to right, top row: CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS, OA; from left to right, bottom row: reverse colonies on CYA, CYA 30 °C, CYA 37 °C, MEA, YES, DG18, CYAS and obverse on CREA; b–h, j–o, q–w. conidiophores, conidia, ascocarps (in v) and asci (in w) produced on OA. — Scale bar: w = 10 µm, applies to all microscope pictures, except v = 20 µm.




Fig. 12 SEM micrographs showing characters of monopialidic *Penicillium* species. a, d. Overview of conidiogenous areas; b, e. phialides; c, f. conidia. — Scale bars: a, d = 10 μ m; b, e = 5 μ m; c, f = 2 μ m.

0.6 \times 2.6 \pm 0.1); *conidia* rough-walled, globose, 2–2.5 \times 2–2.5 μ m (2.3 \pm 0.2 \times 2.3 \pm 0.2, n = 40), average width/length = 0.98; *sclerotia* not produced.

SEM observations: Conidia rough-walled with very large tubercles present, connectives rather short with rings not easily visible, tubercles 0.24–0.38 μ m diam (0.31 \pm 0.04).

Penicillium wollemicola Visagie, Summerell, Shenoy & Seifert, *sp. nov.* — MycoBank MB808274; Fig. 11p–v, 12d–f

ITS Barcode. KJ174314 (alternative markers: *BenA* = KJ174315; *CaM* = KJ174316; *RPB2* = KJ174313).

Etymology. Latin, named after the genus of the tree, *Wollemia*, in reference to the species isolated from its pine needles.

Type specimen. AUSTRALIA, from *Wollemia nobilis* needle debris, 12 Dec. 2007, coll. B. Summerell, isol. S. Shenoy (DAOM 675862 holotype, duplicate dried specimen deposited as CBS H-21615, ex-type CBS 137177 = DTO 297E3).

Diagnosis — No growth on CYA at 37 $^{\circ}$ C, CREA, growth on CYA at 30 $^{\circ}$ C, colonies on CYA 7–8 mm, CYAS 4–5 mm and SNA 7–9 mm, distinctly rough to spiny, globose, 2–2.5 μ m diam. Ascomata pale coloured, asci not seen, ascospores lenticular, smooth-walled with two widely spaced ridges, 2.5–3.5 \times 2.5–3.

Description — Colony diam, 7 d, in mm: CYA 7–8; CYA 30 $^{\circ}$ C 6–7; CYA 37 $^{\circ}$ C no growth; MEA 4–5; YES 8–9; OA 9–10; DG18 9–11; CYAS 4–5; SNA 7–9; CREA no growth.

Colonies at 25 $^{\circ}$ C, 7 d: CYA obverse white, reverse brown (5E5–5F5), inconspicuous brown soluble pigment; MEA obverse white, reverse yellowish brown (5D6); YES obverse orange white (5A2), reverse light orange (5A5); OA obverse greyish green (26C4), brown soluble pigment produced; DG18 obverse orange white (5A2), reverse yellowish white (3A2); CYAS obverse orange white (5A2), reverse yellowish white (3A2); SNA obverse dull green (26D3), reverse greenish grey (26B2); CREA acid not produced.

Conidiophores as solitary phialides; *stipes* smooth-walled, 4–13 \times 1–1.5 μ m; *phialides* ampulliform, 4.5–6.5 \times 2.5–3 μ m (5.6 \pm 0.4 \times 2.8 \pm 0.3); *conidia* distinctly rough to spiny, globose, 2–2.5 \times 2–2.5 μ m (2.3 \pm 0.1 \times 2.3 \pm 0.1, n = 25), average width/length = 0.99; *sclerotia* abundant, colourless, 30–94 \times 24–90

μ m; *cleistothecia* observed on old dry plate, similarly sized to sclerotia, wall pseudoparenchymatous, single layer, asci not observed, *ascospores* lenticular with two widely separated ridges, smooth-walled, 2.5–3.5 \times 2.5–3 μ m.

SEM observations: Conidia rough-walled with tubercles present, connectives relatively long without visible rings, tubercles 0.22–0.41 μ m diam, average 0.31 \pm 0.05.

DUBIOUS SPECIES

Penicillium parviterrucosum (K. Ando & Pitt) Houbraken & Samson, *Stud. Mycol.* 70: 48. 2011 — MycoBank MB561962

Basionym. *Torulomyces parviterrucosus* K. Ando & Pitt, *Mycoscience* 39: 317. 1998.

Type specimen. AUSTRALIA, Sydney, Gap Park, from soil, Oct. 1989, isol. K. Ando (TNS-F-0.5203 holotype, culture ex-type KY12720).

Notes — Because of commercial restrictions placed on the use of the ex-type strain, material was unavailable for examination. Morphologically, the species was distinguished from close relatives by its verrucose conidia that have small tubercles under SEM (Ando et al. 1998). Because of the overlapping characters of conidia observed during SEM examinations with other species in this section, it is clear that DNA sequence data is necessary to determine the precise taxonomic placement of *P. parviterrucosum*. Presently, we consider it a dubious species and should be reconsidered when material is made available.

DISCUSSION

Penicillium sect. *Torulomyces* was introduced by Houbraken & Samson (2011) for *P. lagenae* (CBS 185.65) and the phylogenetically related *P. lassenii* and *P. cryptum*, with the consequence that the genus *Torulomyces* was confirmed as a synonym of *Penicillium* based on multigene phylogenetic evidence. This group of species all display restricted growth on media and the majority produce conidiophores consisting of solitary phialides. The aim of our study was to re-evaluate the taxonomy of species currently classified in the section in the context of the discovery of 12 new species, described above in the taxonomy

section. This considerably extends the species sampling for this section, allowing a stringent evaluation of species concepts and the significance of inconspicuous morphological characters, and some consideration of ecological and biogeographic data in the resulting taxonomic structure. Our multigene phylogenies combined with morphological data recognize 16 species in the section. As noted above, identical ITS sequences are shared by several species; *BenA* was proposed as a secondary marker for *Penicillium* by Visagie et al. (2014b) and using *BenA* works well for identification of species in this section.

Species of sect. *Torulomyces* are morphologically challenging to identify. Despite this, their ability or inability to grow on certain media or at higher temperatures makes identification into smaller phenotypic groups possible. Further, within small groups of phylogenetically related species, phenotypic or morphological characters were discerned to distinguish most species pairs. For example, *P. aeris* and *P. cantabricum* are consistently resolved as close relatives in our phylogenies, but *P. aeris* has rougher conidia than *P. cantabricum* (Fig. 4c, i) and does not grow on CREA. *Penicillium tubakianum* and *P. wollemiicola* are close relatives, but differ in growth rates and conidia tubercle sizes (Fig. 9i, 12f). Our multigene phylogeny places *P. marthae-christenseniae*, which grows on CYA 30 °C, closest to *P. catalanicum*, which does not. Conidial shape also differs between the two species. *Penicillium williamettense* and *P. oregonense* are close relatives, but they have different growth rates and *P. oregonense* has conidia that are roughened in parallel rings (Fig. 6l, 9o). When one considers *P. austriicola* and *P. riverlandense*, clearly distinct species under GCPSR, the only morphological difference is that *P. riverlandense* produces smaller colonies at 37 °C. As the most frequently sampled species, the phylogenetic distance and coherence of each species in the single gene phylogenies support the idea that the observed ecological differences between them, discussed below, indicate they are different species rather than different populations of one species.

SEM examination of conidial ornamentations and connectives also reveals differences that distinguish species. Similar observations were made by Ando et al. (1998), who reported that *P. laeve* and *P. ovatum* have smooth conidia, but they were able to distinguish between *P. lagena* and *P. porphyreum* based on the latter's smaller conidial tubercles. In addition to size differences in tubercles, various aspects of conidial ornamentation are informative for some species. For example, *P. varirattense* had very finely roughened conidia compared to the rather large ornamentations in *P. williamettense*. In *P. catalanicum* and *P. oregonense* conidial ornamentations formed parallel rings. Connectives were also sometimes useful, with those of *P. porphyreum* having two distinct, clearly visible rings. Although the combination of SEM and growth characters under different conditions allows phenotype based identification, routine species identification by these means is unrealistic, because SEM is less accessible and more time consuming than routine molecular methods. Although we incorporated these observations for species delineation in this study, we suggest that DNA sequences should be used for reliable identification.

Nine of the new species are based on single isolates. This is not ideal and reflects the difficulty inherent in isolating such slow growing fungi. Although the two species from the heavily sampled fynbos soils, *P. austriicola* and *P. riverlandense*, are represented by many strains, only one culture of *P. wollemiicola* was isolated from Wollemi pine litter among more than 700 strains isolated via dilution to extinction, a method designed to recover slow growing fungi. Future isolations should result in the recovery of more strains that can be used to better document species of sect. *Torulomyces*. In most cases, the genetic distances between our species, including those represented by single strains, corresponds with that observed among phylo-

genetic species in other groups of *Penicillium*. Although strict application of a phylogenetic species concept requires more than one representative of each taxon, our newly proposed species otherwise conform with GCPSR. Unfortunately, the number of singletons in the phylogenies make bootstrap values and posterior probabilities uninformative to some degree.

Determining the distribution of this group of species from the literature is hampered because until recently, most slow growing, brown, monophialidic strains were considered to represent one species, *P. lagena*, which was considered widespread but rarely isolated (Domsch et al. 1980). Many of the species newly described in this paper were deposited into culture collections as that species. The literature mostly reports *P. lagena*, and the three other previously known species, from soil surveys (Christensen & Backus 1962, Barron 1967, Gochenaur 1978, Domsch et al. 1980, Ando et al. 1998, Renker et al. 2005), although *P. lagena* has also been reported from other sources like dead bark and cornmeal (Domsch et al. 1980). In this study, *P. riverlandense* and *P. austriicola* were frequently isolated from *Protea repens* infructescences and the mites living in them. Our single strain of *P. wollemiicola* was isolated from needle debris of *Wollemia nobilis*, the so-called 'dinosaur tree' or 'living fossil' that was discovered in the Blue Mountains of Australia in 1994 (Jones et al. 1995). *Penicillium austriicola* and *P. riverlandense* were isolated from the fynbos biome situated in the Western Cape of South Africa. One of the fynbos sampling sites in the Riverlands Nature Reserve near Malmesbury was surveyed over a period of four years and *P. riverlandense* was consistently recovered from these samples. In addition, *P. austriicola* was only found at a fynbos sampling site at Struisbaai. Our success at isolating these two species makes us optimistic that the development of appropriate sampling or selective isolation techniques will eventually lead to increased sampling of other species, and better understanding of their ecologies.

Culture independent, environmental DNA sequencing might allow enhanced detection of the distribution of this group. Because ITS, the most frequently used marker in such studies, is uninformative at the species level in this group, presently we can only consider data at the sectional level. A BLAST search with the ITS sequence of *T. lagena* (CBS 185.65; KF303665) on GenBank results in only 14 hits that can be assigned to the section. These hits originate from soil (Czech Republic, one unknown location), plant material (Lithuania, Ecuador), marine sediments (China), plant root tissue (Australia, South Africa), a lichen (*Letharia vulpine*, USA) and house dust (Finland). Most of these hits are from cultured isolates, with only two hits (dust from Finland; soil from unreported country) from culture-independent detection techniques. This seems to support that these species are widespread but occur in very low numbers as reported in Domsch et al. (1980). The diversity of identified habitats from these studies suggests an even greater diversity of trophic modes for these species than we have so far explored. Whether the use of a gene with higher species resolution, such as the *BenA*, *CaM* or *RPB2* used here, will allow more sensitive detection remains to be seen.

In our study, only two of the species, *P. laeve* and *P. wollemiicola* produced ascospores in some culture transfers. The ascospores of *P. laeve* are smooth-walled and globose, while those of *P. wollemiicola* are lenticular and have two broadly spaced ridges. The latter are similar to those described for *P. lagena* (Stolk & Samson 1983, as *Eupen. limoneum*), but we did not observe ascospores in that species. *Penicillium lasseni* was originally described in the sexual genus *Eupenicillium*, but we observed only sclerotia in our strains. The paucity of ascomata observed in our study suggests that either the ability to produce the sexual morph is easily lost in culture if the species are homothallic, or that the species are heterothallic. Thus, it is difficult to confidently interpret sexual characters for identi-

fying species, although the lack of ridges on ascospores of *P. laeve*, and the difference in shape from the other two species, presently seems diagnostic.

Our phylogenies reveal that *P. laeve* and *P. ovatum*, previously included in sect. *Torulomyces* by Houbraken & Samson (2011), should be classified in sect. *Exilicaulis* with *P. dimorphosporum* as their closest relative. The placement of *P. lassenii* and *P. cryptum* in sect. *Torulomyces* remains questionable, because our phylogenies resolve them as distant relatives to the well-supported clade that contains the majority of species of sect. *Torulomyces* (Fig. 1, 2). A BLAST search of the *P. lassenii* ITS sequence results in the closest match of 93 % to *P. sublateralium* and *P. cyaneum*, classified in sect. *Ramigena*. For *P. cryptum*, a BLAST search results in a 93 % match to *P. donkii* and *P. boreae*, classified in sect. *Stolkia*. The closest GenBank match for *RPB2* of *P. lassenii* and *P. cryptum*, respectively results in a 84 % similar match to *P. ornatum* and 83 % similar to *P. lagenae*. Houbraken & Samson (2011) used *RPB1*, *RPB2*, *Tsr1* and *Cct8* for their sectional reclassification of *Penicillium*, which placed both species in a clade with *P. lagenae*. However, the branches for these three species in the clade were very long and perhaps provided deceptive support for recognising them as separate and distinct sections. In a similar situation, Visagie et al. (2014a) recently described *P. alfredii*, but were unable to classify it in any of the 25 sections proposed by Houbraken & Samson (2011). Although it seemed that *P. alfredii* represented a new section, they opted to leave it unclassified because of the uncertain phylogenetic position of *P. cryptum* and *P. lassenii*. In their *RPB2* phylogeny, *P. cryptum* made sect. *Torulomyces* polyphyletic with sect. *Fracta*. The fact that these two species are inconsistently placed with seemingly every gene analysed, hints that they may represent two distinct sections that are presently undersampled and thus unrecognizable by cladistic approaches. Conidiophore branching patterns further support the idea that they do not belong in sect. *Torulomyces*, but without isolating additional strains, closely related species and/or analysing more alternative gene sequences, the placement of *P. cryptum*, *P. lassenii* and *P. alfredii* will remain problematic. As a result, for the time being, we believe it best to leave *P. cryptum* and *P. lassenii* in sect. *Torulomyces* as proposed by Houbraken & Samson (2011).

Acknowledgements We acknowledge the South African Biosystematics Initiative (SABI) of the National Research Foundation (NRF) and the Alfred P. Sloan Foundation for financial support. We thank Uwe Braun for providing his assistance on nomenclatural issues. We are also grateful to Martha Christensen who provided many of the strains described here as new species, Brett Summerell for collecting Wollemi pine litter and Damodar Shenoy for isolating *P. wollemiicola* from that substrate, and Katsuhiko Ando who provided the ex-type strains of *P. ovatum* and *P. laeve*.

REFERENCES

Ando K, Nawawi A, Manoch L, et al. 1998. Three new species and a new combination in the genus *Torulomyces* from soil. *Mycoscience* 39: 313–318.

Barron GL. 1967. *Torulomyces* and *Monocillium*. *Mycologia* 59: 716–718.

Christensen M, Backus MP. 1962. A new *Penicillium* from coniferous forest soils. *Mycologia* 54: 573–577.

Delitsch H. 1943. *Ergebnisse der theoretischen und angewandten Mikrobiologie: Band I: Systematik der Schimmelpilze*. Neumann, Neudamm, Germany.

Domsch KH, Gams W, Anderson T-H. 1980. *Compendium of soil fungi*. IHW-Verlag, Eching.

Gams W. 1971. *Cephalosporium-artige Schimmelpilze (Hyphomycetes)*. Gustav Fischer Verlag, Stuttgart.

Gams W. 1978. Connected and disconnected chains of phialoconidia and *Sagenomella* gen. nov. segregated from *Acremonium*. *Persoonia* 10: 97–112.

Gochenaour SE. 1978. Fungi of a Long Island oak-birch forest I. Community Organization and Seasonal Occurrence of the Opportunistic Decomposers of the A Horizon. *Mycologia* 70: 975–994.

Gouy M, Guindon S, Gascuel O. 2010. SeaView Version 4: A Multiplatform Graphical User Interface for Sequence Alignment and Phylogenetic Tree Building. *Molecular Biology and Evolution* 27: 221–224.

Hashmi MH, Kendrick B, Morgan-Jones G. 1972. Conidium ontogeny in hyphomycetes. The genera *Torulomyces* Delitsch and *Monocillium* Saksena. *Canadian Journal of Botany* 50: 1461–1463.

Houbraken J, Samson RA. 2011. Phylogeny of *Penicillium* and the segregation of Trichocomaceae into three families. *Studies in Mycology* 70: 1–51.

Jones WG, Hill KD, Allen JM. 1995. *Wollemia nobilis*, a new living Australian genus and species in the Araucariaceae. *Telopea* 6: 173–176.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30: 772–780.

Kornerup A, Wanscher JH. 1967. *Methuen Handbook of Colour*. Methuen & Co Ltd, Copenhagen.

Liang ZQ, Han YF, Chu HL, et al. 2009. Studies on the genus *Paecilomyces* in China V. *Taifanglania* gen. nov. for some monophialidic species. *Fungal Diversity* 34: 69–77.

Matsushima T. 1987. *Matsushima mycological memoirs no. 5*: 1–100.

McNeill J, Barrie FR, Buck WR, et al. 2012. *International Code of Nomenclature for algae, fungi and plants (Melbourne Code)*. Koeltz Scientific Books Königstein. (Regnum vegetabile no. 154).

Nylander AJJ, Ronquist F, Huelsenbeck JP, et al. 2004. Bayesian phylogenetic analysis of combined data. *Systematic Biology* 53: 47–67.

Onions AHS, Barron GL. 1967. Monophialidic species of *Paecilomyces*. *Mycological Papers* 107: 1–25.

Pitt JI, Hocking AD. 1985. Interfaces among genera related to *Aspergillus* and *Penicillium*. *Mycologia* 77: 810–824.

Pitt JI, Samson RA. 1993. Names in current use in the family Trichocomaceae. In: Greuter W (ed), *Names in current use in the family Trichocomaceae, Cladoniaceae, Pinaceae, and Lemnaceae*: 13–57. Koeltz Scientific Books, Germany.

Pitt JI, Samson RA, Frisvad JC. 2000. List of accepted species and their synonyms in the family Trichocomaceae. In: Samson RA, Pitt JI (eds), *Integration of modern taxonomic methods for Penicillium and Aspergillus classification*: 9–79. Harwood Academic Publishers, Amsterdam, The Netherlands.

Renker C, Otto P, Schneider K, et al. 2005. Oribatid mites as potential vectors for soil microfungi: study of mite-associated fungal species. *Microbial Ecology* 50: 518–528.

Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.

Saksena SB. 1955. A new fungus, *Monocillium indicum* gen. et sp. nov., from soil. *Indian Phytopathology* 8: 9–12.

Samson RA. 1974. A monograph of *Paecilomyces* and allied hyphomycetes. *Studies in Mycology* 6: 1–119.

Samson RA, Visagie CM, Houbraken J, et al. 2014. Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology* 78: 141–173.

Schoch CL, Seifert KA, Huhndorf S, et al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of the United States of America* 109: 6241–6246.

Seifert KA, Morgan-Jones G, Gams W, et al. 2011. The genera of Hyphomycetes. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.

Sigler L, Sutton DA, Gibas CFC, et al. 2010. *Phialosimplex*, a new anamorphic genus associated with infections in dogs and having phylogenetic affinity to the Trichocomaceae. *Medical Mycology* 48: 335–345.

Stolk AC, Samson RA. 1983. The Ascomycete genus *Eupenicillium* and related *Penicillium* anamorphs. *Studies in Mycology* 23: 1–149.

Stolk AC, Samson RA. 1985. A new taxonomic scheme for *Penicillium* anamorphs. In: Samson RA, Pitt JI (eds), *Advances in Penicillium and Aspergillus systematics*: 163–192. Plenum Press, New York, USA.

Tamura K, Stecher G, Peterson D, et al. 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 30: 2725–2729.

Taylor JW, Jacobson DJ, Kroken S, et al. 2000. Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* 31: 21–32.

Visagie CM, Hirooka Y, Tanney JB, et al. 2014a. *Aspergillus*, *Penicillium* and *Talaromyces* isolated from house dust samples collected around the world. *Studies in Mycology* 78: 63–139.

Visagie CM, Houbraken J, Frisvad JC, et al. 2014b. Identification and nomenclature of the genus *Penicillium*. *Studies in Mycology* 78: 343–371.

Visagie CM, Houbraken J, Rodrigues C, et al. 2013. Five new *Penicillium* species in section *Sclerotiora*: a tribute to the Dutch Royal family. *Persoonia* 31: 42–62.

Zhang Y, Liu F, Wu W, et al. 2015. A phylogenetic assessment and taxonomic revision of the thermotolerant hyphomycete genera *Acrophialophora* and *Taifanglania*. *Mycologia* 107: 768–779.