



The species of *Citrus* (Rutaceae) with pinnate leaves

D.J. Mabberley¹

Key words

Aurantiodeae
Citrus
Feroniella
Rutaceae

Abstract *Feroniella* (Rutaceae: Aurantiodeae) is formally transferred to *Citrus* and a new combination, *C. lucida*, made.

Published on 16 April 2010

INTRODUCTION

In a recent molecular analysis (Bayer et al. 2009), based on nine cpDNA sequences, the circumscription of the genus *Citrus* L. (Rutaceae: Aurantiodeae, Aurantieae [Citreae]) was broadened beyond that of Mabberley (1998, 2004, 2008, where *Clymenia* Swingle, *Fortunella* Swingle, *Poncirus* Raf., *Microcitrus* Swingle and *Eremocitrus* Swingle are reunited with the genus), to include *Oxanthera* Montrouz. and, very surprisingly – because its species, unlike those of all the others, have never been included in *Citrus* before – *Feroniella* Swingle. Until this revelation, *Citrus* was considered (Zhang & Mabberley 2008) to comprise species with simple, unifoliolate (often with a markedly winged petiole) or trifoliolate (*C. trifoliata* L., formerly referred to the genus *Poncirus*) leaves, but the species hitherto referred to *Feroniella* has imparipinnate leaves. They are deciduous, like those of the hardy *C. trifoliata*, and both deciduousness and the toughened pericarp seem to be adaptations to the seasonal forests where the species is found.

The range of leaf form in the genus gives support to the hypothesis of Corner (1964: 146, t. 50), who considered that the unifoliolate leaf typical of most species of *Citrus* represents the terminal leaflet of an incompletely developed pinnate leaf typical of Sapindales in general. Unifoliolate or simple and trifoliolate leaves are found within several other genera of subfam. Aurantiodeae, e.g. *Aeglopsis* Swingle, *Balsamocitrus* Stapf, *Burkillanthus* Swingle, *Luvunga* Buch.-Ham. ex Wight & Arn. (from which the unifoliolate/simple-leaved *Paramignya* Wight may not be distinct – Mabberley 1998), *Pleiospermium* (Engl.) Swingle and *Triphasia* Lour.; the full range to imparipinnate is found in *Citropsis* (Engl.) Swingle & Kellerm. and *Naringi* Adans.

As its name suggests, *Feroniella* was formerly considered allied to *Limonia* L. (syn. *Feronia*), being classed as one of the 'wood apples' of Swingle (1943) because of its tough pericarp. *Limonia*, however, is diplostemonous whereas species of *Feroniella* have four times as many stamens as petals as is typical of *Citrus*.

NOMENCLATURE

Citrus L.

Citrus L., Sp. Pl. 2 (1753) 782. — Type: *Citrus medica* L.
Feroniella Swingle (1913) 776, syn. nov. — Type: *Feroniella oblata* Swingle = *Citrus lucida* (Scheff.) Mabb.

According to Swingle (1943: 468–471, q.v. for key and species descriptions), there are three pinnate-leaved species, but one of these (Forman 1958), *F. pubescens* Tanaka, is actually *Harrisonia perforata* (Blanco) Merr. (Rutaceae), while the other two, following Guillaumin (1946: 651), are now considered conspecific. In readiness for an account for *Flora malesiana* the single recognised 'Feroniella' species is here formally transferred to *Citrus*:

1. *Citrus lucida* (Scheff.) Mabb., comb. nov.

Feronia lucida Scheff., Natuurk. Tijdschr. Ned.-Indië 31 (1870) 19; Icon. Bogor. 2 (1904) t. 149.
Feroniella lucida Swingle (1913) 781; (1943) 470, t. 74; Guillaumin (1946) 651. — Type: Anon. s.n. (ex Rembang coll. Teijsmann) (holo BO; iso L, U), Indonesia, Java, cult. Hort. Bogor.
Feroniella oblata Swingle (1913) 779; (1943) 469. — Type: *Pierre* 652 (holo P), Cambodia, Samroing-Aong.

Distribution — Cambodia, Laos, ?Vietnam, Thailand, Java (teak forests)

Notes — I am grateful to J-F Veldkamp for confirmation of L and U sheets as isotypes of *F. lucida*. *Feroniella puberula* Tanaka, Stud. Citrol. 2 (1928) 23 is a *nomen nudum*.

REFERENCES

- Bayer R, Mabberley DJ, Morton C, Miller CH, Sharma IK, Pfeil BE, Rich S, Hitchcock R, Sykes S. 2009. A molecular phylogeny of the orange subfamily (Rutaceae: Aurantiodeae) using nine cpDNA sequences. *American Journal of Botany* 96: 668–685.
Corner EJH. 1964. The life of plants. Weidenfeld & Nicolson, London.
Forman LL. 1958. The identity of *Feroniella pubescens* Tanaka (Rutaceae). *Kew Bulletin* 12: 503–504.
Guillaumin A. 1946. Rutaceae. In: Humbert H, Supplément à la Flore Générale de l'Indochine: 593–657. Muséum National d'Histoire Naturelle Phanérogamie, Paris.
Mabberley DJ. 1998. Australian Citreae with notes on other Aurantiodeae (Rutaceae). *Telopea* 7: 333–344.
Mabberley DJ. 2004. *Citrus* (Rutaceae): a review of recent advances in etymology, systematics and medical applications. *Blumea* 49: 481–498.

¹ Herbarium, Library, Art and Archives, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, United Kingdom; Netherlands Centre for Biodiversity Naturalis (section NHN), Leiden University, P.O. Box 9514, 2300 RA Leiden, The Netherlands; corresponding author e-mail: d.mabberley@kew.org.

- Mabberley DJ. 2008. Mabberley's Plant-book. A portable dictionary of plants, their classification and uses. Third edition. Cambridge University Press, Cambridge, UK.
- Scheffer RHCC. 1870. Observationes phytogeographicae. *Natuurkundig Tijdschrift voor Nederlandsch-Indië* 31: 1–23.
- Swingle WT. 1913. *Feroniella*, genre nouveau de la tribu des Citreae, fondé sur le *F. oblata*, espèce nouvelle de l'Indochine. *Bulletin de la Société Botanique de France* 59: 774–783.
- Swingle WT. 1943. The botany of Citrus and its wild relatives of the orange subfamily (family Rutaceae, subfamily Aurantioideae). In: Webber HJ, Batchelor LD, *The Citrus Industry*, vol. 1. History, botany, and breeding: 129–474. University of California Press, Berkeley & Los Angeles.
- Zhang D-X, Mabberley DJ. 2008. Citrus. *Flora of China* 11: 90–96.