

TREES OF LAOS AND VIETNAM: A FIELD GUIDE TO 100 ECONOMICALLY OR ECOLOGICALLY IMPORTANT SPECIES

HOANG VAN SAM¹, KHAMSENG NANTHAVONG² & P.J.A. KESSLER³

SUMMARY

This field guide to 100 economically or ecologically important tree species from Laos and Vietnam enables the user to identify the included taxa with user-friendly keys. It includes scientific names, botanical descriptions of families, genera, and species. Specific information on distribution, habitat, ecology, and uses has been compiled. All specimens examined have been listed.

Key words: Flora of Laos and Vietnam, field guide, tree-identification, tree flora, Indochina.

INTRODUCTION

Laos and Vietnam's forests are one of the countries' richest natural resources, supporting a huge diversity of plant and animal life, and providing forest products to support local livelihoods. In recent years forest cover in the tropics has decreased drastically and Laos and Vietnam are among the few South East Asian countries where still a substantial part of the land is covered by often unexplored, practically unknown original vegetation. Except 'Flore du Laos, du Cambodge et du Vietnam' and its predecessors there is no publication which enables us to identify plants with an easy to use key. Almost all other publications are descriptive but without any means to identify plants except comparing either descriptions or photographs. Our manual aims at assisting foresters and botanists and their students with user-friendly keys in the identification of some major components of the forests. We hope that our contribution will stimulate further studies of the flora.

SCOPE OF THE FIELD GUIDE

The intended scope of the manual is a Flora containing 100 tree species belonging to 83 genera and 40 families from Vietnam and Laos, important either economically or ecologically, and the means to identify them by field as well as herbarium characters. This manual contains all the information which is needed for identification, including

-
- 1) Vietnam Forestry University, Faculty of Forest Resources & Environment Management, Xuan Mai, Chuong My, Ha Tay, Vietnam.
 - 2) National University of Laos, Faculty of Forestry, Department of Forest Management, P.O. Box 7322, Vientiane, Lao PDR.
 - 3) Nationaal Herbarium Nederland, Universiteit Leiden branch, P.O. Box 9514, 2300 RA Leiden, The Netherlands; e-mail: Kessler@nhn.leidenuniv.nl.

full scientific names, synonyms, literature, vernacular names, botanical descriptions, distribution, habitat & ecology, uses, and specimens examined. Keys to the families, genera, and species are given.

FORMAT OF DESCRIPTIONS

- Families, genera, and species are treated in alphabetical order within the two major groups ‘Gymnospermae’ and ‘Angiospermae’.
- If a family contains more than 1 genus, a key to the genera is given. If a genus has more than one species, a key to the species is given.
- Scientific genera and species names. Nomenclature generally follows recent literature, however, in some cases we have used names that are common use in the Floras of Laos and Vietnam.
- Synonyms are limited to those used in recent literature.
- Vernacular names have been cited using two main languages: Lao (L) and Vietnamese (V).
- Descriptions are based on literature and herbarium specimens kept at Leiden (L) and Paris (P).
- Distribution for the plant families is mentioned including the number of genera and species in the world, Laos, and Vietnam. Distribution within both countries is by province using the names as published in ‘Flore du Cambodge, du Laos et du Vietnam’ 31 (2003) 96.
- Habitat and ecology information include data on forest type, the habitats and altitude where the species are occurring, and the flowering and fruiting time.
- Data on uses.

ACKNOWLEDGEMENTS

This field guide is a reviewed version of the MSc theses of Hoang Van Sam and Khamseng Nanthavong under the supervision of P.J.A. Keßler at the Nationaal Herbarium Leiden, Universiteit Leiden branch.

We would like to thank the co-directors of the ASEAN Regional Centre for Biodiversity Conservation (ARCBC) for enabling HVS and KN to pursue their MSc at Leiden.

The director of the Nationaal Herbarium Nederland, P. Baas, the head of the Project Group Plant diversity of the Indo-Pacific and Tropical Asia, M.C. Roos, and the staff supported our activities. F.A.C.B. Adema, M.M.J. van Balgooy, Ding Hou, J.B. Mols, H.P. Nooteboom, W. Vink, P.C. van Welzen, and W.J.J.O. de Wilde kindly helped us with their expertise on various plant families.

Hoang Van Sam would like to thank the rector, the dean of the Forest Resources and Environment Management Faculty, the head of the Forest Plants department and his colleagues at the Vietnam Forestry University to enable him to study abroad and to support all his activities.

Khamseng Nanthavong received much support from the Ministry of Education, the rector of the National University of Laos and the dean of the Faculty of Forestry enabling him to pursue a MSc study in Leiden.

The director of the herbarium at P kindly allowed us to study the Indochinese collections during a visit and to borrow quite a lot of material. Th. Deroin supported us in various ways during our stay. S. Hul provided us with information concerning localities and general data on the Indochinese Flora. J.E. Vidal helped us in finding relevant literature especially for Laos.

We are very grateful to F.A.C.B. Adema for critically reading earlier versions of the manuscript. Ms. C.G.G. Baak kindly formatted the manuscript for which she is gratefully acknowledged.

Miss Joanne Porck (NHN-Leiden) and Mr. Priono (The International Tropenbos Kalimantan project) made several original drawings for which we are thankful. The authors wish to express their gratitude to the curators of P and BKF for the permission to use illustrations from *Flore du Cambodge, du Laos et du Vietnam* and *Thai Forest Bulletin*.

SELECTED LITERATURE

- Backer, C.A. & R.C. Bakhuizen van den Brink Jr. 1964–1965. *Flora of Java*. Vol. 1, 2. Noordhoff, Groningen, The Netherlands.
- Bân, N.T. 1996. *Red data book of Vietnam*. Vol. 2. Plants. Science & Technics Publishing House, Hanoi, Vietnam.
- Bân, N.T. 2000. *Flora of Vietnam*. Vol. 1. Science & Technics Publishing House, Hanoi, Vietnam. (In Vietnamese.)
- Bodegom, S., P.B. Pelser & P.J.A. Keßler. 1999. Seedlings of secondary forest tree species of East Kalimantan, Indonesia. *Tropenbos-Kalimantan Series 1*. Balikpapan.
- Callaghan, R.M. 1995. *Plants of Laos*. A list of tree species. Vientiane, Laos.
- Chan, L.M. & L.T. Huyen. 2000. *Forest plants of Vietnam*. Hanoi, Vietnam. (In Vietnamese.)
- Corner, E.J.H. 1940. *Wayside Trees of Malaya*. Singapore.
- De Vogel, E.F. (ed.). 1987. *Manual of herbarium taxonomy. Theory and practice*. UNESCO. Jakarta, Indonesia.
- Dung, V.V. 1996. *Vietnam Forest Trees*. Hanoi, Vietnam.
- Hooker, J.D. 1872–1890. *Flora of British India*. Vol. 3–5, London, United Kingdom.
- Flora of China*. 1999–2000. Vol. 4–24, Beijing, China.
- Flora Malesiana*. 1948–2000. Ser. 1, Vol. 4–14. Nationaal Herbarium Nederland, Universiteit Leiden branch, The Netherlands.
- Flora of Thailand*. 1970–2002. Vol. 2–7. The Forest Herbarium, Royal Forest Department, Bangkok, Thailand.
- Flore du Laos, du Cambodge et du Vietnam*. 1960–2003. Vol. 1–31. Muséum National d'Histoire Naturelle, Laboratoire de Phanérogamie, Paris, France. (In French.)
- Gardner, S., P. Sidisunthorn & V. Anusarnsunthorn. 2000. *A field guide to forest trees of northern Thailand*. Kobfai Publishing Project, Bangkok, Thailand.
- Hô, P.H. 1991–1992. *Flora of Vietnam*, Vol. 1, 2. Paris. (In Vietnamese.)
- Ke, L.K. 1969–1977. *The common plants in Vietnam*. Vol. 1–6. Hanoi. (In Vietnamese.)
- Keßler, P.J.A. (ed.) 2000. *Secondary forest trees of Kalimantan, Indonesia*. MOFEC Tropenbos-Kalimantan Series 3. The Netherlands.
- Lecomte, H. (ed.). 1907–1912. *Flore Générale de l'Indo-Chine*. Vol. 1–5. Muséum National d'Histoire Naturelle, Laboratoire de Phanérogamie, Paris, France.
- Lehmann, L., M. Greijmans & D. Shenman. 2003. *Forests and trees of the central highlands of Xieng Khouang Lao PDR, A field guide*. Lao Tree Seed Project, Vientiane, Laos.
- PROSEA (Plant Resources of South-East Asia). 1993–2003. Vol. 5–17. PUDOC Scientific Publishers, Wageningen, The Netherlands.
- Ridley, H.N. 1923–1925. *Flora of Malay Peninsula*. Vol. 1–15. Reeve & Co, London, United Kingdom.
- Tree Flora of Malaya*. 1972–1989. Vol. 1–4. Forest Research Institute Malaysia, Kepong, Malaysia.
- Tree Flora of Sabah and Sarawak*. 1995–2002. Vol. 1–4. Forest Research Institute Malaysia, Kepong, Malaysia.
- Trung, V.T. 1970. *The vegetation of Vietnam*. Hanoi, Vietnam. (In Vietnamese.)
- Vidal, J.E. 1959. Noms vernaculaires de plantes (LAO, MËO, KHA) en usage au Laos. *Bull. de l'École Française d'Extrême-Orient* 49: 435–608.
- Vidal, J.E. 2000. *Vegetation-types and plants of the Indochinese Peninsula*. Ecocart, Paris, France. (CD-ROM).

FLORISTIC TREATMENT

SPOT CHARACTERS FOR THE GENERA

List of spot characters

- | | |
|---|---|
| 1 – Gymnosperms | 28 – Leaves peltate |
| 2 – Bark with hissing sound when cut | 29 – Leaves trifoliolate |
| 3 – Armed plants | 30 – Leaflets alternate |
| 4 – Stem flanged | 31 – Leaflets opposite |
| 5 – Twigs with ridges | 32 – Leaf or leaflet margin entire |
| 6 – Sap white | 33 – Leaf or leaflet margin toothed |
| 7 – Sap yellow | 34 – Leaves hairy |
| 8 – Sap black or brown | 35 – Leaves or leaflets with dots |
| 9 – Sap red or orange | 36 – Leaves or leaflets with domatia |
| 10 – Living parts of plants with smell | 37 – Leaves 3-veined |
| 11 – Stellate hairs or scales present | 38 – Intramarginal vein present |
| 12 – Glands on petiole | 39 – Leaves with very close parallel veins |
| 13 – Glands on leaf lamina | 40 – Inflorescences a spike |
| 14 – Stipules present | 41 – Inflorescences in a head |
| 15 – Stipules absent | 42 – Flowers large and showy (at least
1.5 cm diam.) |
| 16 – Petioles swollen | 43 – Ovary inferior |
| 17 – Petioles wrinkled | 44 – Ovary superior |
| 18 – Winged rhachis or petiole | 45 – Fruits compound |
| 19 – Rhachis with swollen nodes | 46 – Fruits dehiscent |
| 20 – Leaves opposite | 47 – Fruits indehiscent |
| 21 – Leaves whorled | 48 – Fruits with wing(s) |
| 22 – Leaves spirally arranged | 49 – Fruit a berry |
| 23 – Leaves alternate (in two rows) | 50 – Fruit a cone |
| 24 – Leaves simple | 51 – Fruit a drupe |
| 25 – Leaves palmately compound or
digitate | 52 – Fruit a nut |
| 26 – Leaves pinnately compound | 53 – Fruit a capsule |
| 27 – Leaves bipinnately compound | 54 – Fruit a pod |
| | 55 – Seeds winged |

Spot characters

1. Gymnosperms
Cupressus, Dacrycarpus, Dacrydium, Fokienia, Keteleeria, Nageia, Pinus, Podocarpus
2. Bark with hissing sound when cut
Dillenia
3. Armed plants
Caesalpinia p.p., Erythrina p.p., Gmelina p.p., Pithecellobium p.p.

4. Stem flanged
Archidendron p.p.
5. Twigs with ridges
Duabanga
6. Sap white
*Aglai*a p.p., *Alstonia*, *Broussonetia*, *Canarium*, *Eberhardtia*, *Madhuca*, *Wrightia*
7. Sap yellow
Mesua
8. Sap black or brown
Canarium p.p., *Choerospondias*, *Dracontomelon*
9. Sap red or orange
Bischofia, *Dalbergia*, *Endospermum* p.p., *Horsfieldia*, *Knema*, *Pterocarpus*
10. Living parts of plants with smell
Cananga, *Canarium*, *Cinnamomum*, *Cupressus*, *Dracontomelon*, *Fokienia*, *Michelia*
11. Stellate hairs or scales present
*Aglai*a, *Aleurites*, *Bombax* p.p., *Endospermum*, *Lagerstroemia*, *Pterospermum*
12. Glands on petiole
Archidendron, *Terminalia* p.p.
13. Glands on leaf lamina
Diospyros, *Endospermum*, *Fagraea*, *Gmelina*, *Terminalia* p.p., *Vatica*
14. Stipules present
Acrocarpus, *Adenanthera*, *Afzelia*, *Albizia*, *Aleurites*, *Altingia*, *Anisoptera*, *Archidendron*, *Baccaurea*, *Berrya*, *Betula*, *Bischofia*, *Bombax*, *Broussonetia*, *Caesalpinia*, *Canarium* p.p., *Cassia*, *Castanea*, *Castanopsis*, *Dalbergia*, *Delonix*, *Dialium*, *Dipterocarpus*, *Duabanga*, *Eberhardtia*, *Erythrina*, *Fagraea*, *Hopea*, *Irvingia*, *Liquidambar*, *Madhuca*, *Michelia*, *Parashorea*, *Parkia*, *Peltophorum*, *Pithecellobium*, *Pterocarpus*, *Pterospermum*, *Pyrus*, *Samanea*, *Senna*, *Sindora*, *Tamarindus*, *Vatica*, *Xylia*
15. Stipules absent
Alstonia, *Cananga*, *Canarium* p.p., *Chisocheton*, *Choerospondias*, *Chukrasia*, *Dillenia*, *Diospyros*, *Dracontomelon*, *Gmelina*, *Horsfieldia*, *Knema*, *Lagerstroemia*, *Mesua*, *Oroxylum*, *Polyalthia*, *Sapindus*, *Schima*, *Syzygium*, *Tectona*, *Terminalia*, *Tetrameles*, *Wrightia*, *Xylopi*a
16. Petioles swollen
*Aglai*a p.p., *Aleurites*, *Anisoptera*, *Baccaurea*, *Dipterocarpus*, *Endospermum*, *Pterospermum*

17. Petioles wrinkled
Diospyros p.p., *Syzygium* p.p.
18. Winged rhachis or petiole
Dillenia p.p., *Tectona*
19. Rhachis with swollen nodes
Archidendron, *Oroxylum indicum*
20. Leaves opposite
Duabanga, *Fagraea*, *Gmelina*, *Lagerstroemia*, *Markhamia*, *Mesua*, *Oroxylum*, *Syzygium*, *Tectona*, *Terminalia*, *Wrightia*
21. Leaves whorled
Alstonia, *Terminalia* p.p.
22. Leaves spirally arranged
Aleurites, *Dillenia*, *Dracontomelon*, *Endospermum*, *Senna*, *Terminalia* p.p.
23. Leaves alternate (in two rows)
Aquilaria, *Bombax*, *Caesalpinia*, *Cananga*, *Diospyros*, *Dipterocarpus*, *Duabanga*, *Erythrina*, *Polyalthia*, *Terminalia*, *Xylopia*
24. Leaves simple
Aleurites, *Alstonia*, *Altingia*, *Anisoptera*, *Aquilaria*, *Baccaurea*, *Berrya*, *Bischofia*, *Broussonetia*, *Cananga*, *Castanea*, *Castanopsis*, *Dillenia*, *Diospyros*, *Dipterocarpus*, *Duabanga*, *Eberhardia*, *Endospermum*, *Fagraea*, *Gmelina*, *Horsfieldia*, *Knema*, *Lagerstroemia*, *Liquidambar*, *Irvingia*, *Madhuca*, *Mesua*, *Michelia*, *Parashorea*, *Polyalthia*, *Pterospermum*, *Schima*, *Shorea*, *Syzygium*, *Tectona*, *Terminalia*, *Vatica*, *Wrightia*, *Xylopia*
25. Leaves palmately compound or digitate
Bombax
26. Leaves pinnately compound
Aglaia, *Canarium*, *Cassia*, *Chisocheton*, *Choerospondias*, *Chukrasia*, *Dracontomelon*, *Markhamia*, *Pterocarpus*, *Sapindus*, *Senna*
27. Leaves bipinnately compound
Acrocarpus, *Archidendron*, *Caesalpinia*, *Oroxylum* (sometimes 3-pinnate), *Parkia*, *Peltophorum*
28. Leaves peltate
Endospermum p.p., *Pterospermum*
29. Leaves trifoliolate
Bischofia, *Erythrina*
30. Leaflets alternate
Chukrasia, *Dalium*, *Dracontomelon*, *Sapindus*

31. Leaflets opposite

Acrocarpus, Afzelia, Aglaia, Caesalpinia, Canarium, Chisocheton, Choerospondias, Delonix, Dracontomelon, Markhamia, Oroxyllum, Peltophorum, Sindora, Tamarindus

32. Leaf or leaflet margin entire

Acrocarpus, Afzelia, Aglaia, Aleurites, Alstonia, Anisoptera, Aquilaria, Archidendron, Caesalpinia, Cananga, Cassia, Cinnamomum, Delonix, Dialium, Diospyros, Dipterocarpus, Dracontomelon, Duabanga, Eberhardtia, Endospermum, Fagraea, Gmelina, Hopea, Horsfieldia, Knema, Lagerstroemia, Madhuca, Michelia, Oroxyllum indicum, Parashorea, Parkia, Peltophorum, Polyalthia, Pterocarpus, Pterospermum, Senna, Shorea, Sindora, Syzygium, Tamarindus, Vatica, Wrightia, Xylopia

33. Leaf or leaflet margin toothed

Altingia, Betula, Bischofia, Castanea, Castanopsis, Choerospondias, Dillenia p.p., Gmelina p.p., Markhamia, Schima, Terminalia, Tetrameles

34. Leaves hairy

Broussonetia, Wrightia

35. Leaves or leaflets with dots

Aglaia p.p., Caesalpinia, Cananga, Cinnamomun, Diospyros, Syzygium, Terminalia

36. Leaves or leaflets with domatia

Chukrasia, Dracontomelon, Hopea, Terminalia

37. Leaves 3-veined

Cinnamomun

38. Intramarginal vein present

Duabanga, Syzygium

39. Leaves with very close parallel veins

Alstonia, Mesua

40. Inflorescences a spike

Castanea, Castanopsis, Betula, Broussonetia (male), Keteleeria (male), Liquidambar (male), Pinus (male), Podocarpus (male)

41. Inflorescences in a head

Altingia (female), Broussonetia (female), Liquidambar (female)

42. Flowers large and showy (at least 1.5 cm diam.)

Bombax, Cananga, Delonix, Dillenia, Gmelina, Markhamia, Mesua, Michelia, Oroxyllum, Senna

43. Ovary inferior

Altingia, Anisoptera, Anogeissus, Betula, Castanea, Castanopsis, Lagerstroemia, Liquidambar, Pyrus, Syzygium, Terminalia, Tetrameles

44. Ovary superior

Acrocarpus, Adenanthera, Afzelia, Aglaia, Albizia, Aleurites, Alstonia, Aquilaria, Archidendron, Baccaurea, Berrya, Bischofia, Bombax, Broussonetia, Caesalpinia, Cananga, Canarium, Cassia, Chisocheton, Choerospondias, Chukrasia, Cinnamomum, Dalbergia, Delonix, Dialium, Dillenia, Diospyros, Dipterocarpus, Dracontomelon, Duabanga, Eberhardtia, Endospermum, Erythrina, Fagraea, Gmelina, Hopea, Horsfieldia, Irvingia, Knema, Lagerstroemia, Madhuca, Markhamia, Mesua, Michelia, Oroxyllum, Parashorea, Parkia, Peltophorum, Pithecellobium, Polyalthia, Pterocarpus, Pterospermum, Samanea, Sapindus, Schima, Senna, Shorea, Sindora, Tamarindus, Tectona, Vatica, Wrightia, Xylia, Xylopi

45. Fruits compound

Broussonetia

46. Fruits dehiscent

Acrocarpus, Adenanthera, Afzelia, Aglaia, Albizia, Alstonia, Altingia, Aquilaria, Archidendron, Berrya, Bombax, Caesalpinia, Chisocheton, Chukrasia, Delonix, Eberhardtia, Erythrina, Horsfieldia, Knema, Lagerstroemia, Liquidambar, Markhamia, Mesua, Oroxyllum, Peltophorum, Pithecellobium, Pterospermum, Schima, Sindora, Tetrameles, Wrightia, Xylia

47. Fruits indehiscent

Anisoptera, Anogeissus, Cananga, Cassia, Choerospondias, Dalbergia, Dialium, Dillenia, Dipterocarpus, Dracontomelon, Gmelina, Hopea, Parashorea, Parkia, Polyalthia, Pterocarpus, Samanea, Senna, Shorea, Tamarindus, Tectona grandis, Terminalia, Vatica, Xylopi p.p.

48. Fruit with wing(s)

Anisoptera, Anogeissus, Betula, Dipterocarpus, Hopea, Parashorea, Pterocarpus, Shorea, Terminalia, Vatica

49. Fruit a berry

Baccaurea, Diospyros, Duabanga, Madhuca, Pyrus, Syzygium

50. Fruit a cone

Cupressus, Dacrycarpus, Dacrydium, Fokienia, Keteleeria, Nageia, Pinus, Podocarpus

51. Fruit a drupe

Aleurites, Baccaurea, Bischofia, Cananga, Canarium, Choerospondias, Cinnamomum, Dracontomelon, Endospermum, Fagraea, Gmelina, Irvingia, Polyalthia, Sapindus, Tectona grandis, Xylopi

52. Fruit a nut

Anisoptera, Betula, Castanea, Castanopsis, Dipterocarpus, Hopea, Parashorea, Shorea p.p., *Vatica* p.p.

53. Fruit a capsule

Aglaiia, Alstonia, Altingia, Aquilaria, Berrya, Bombax, Chisocheton, Chukrasia, Eberhardtia, Horsfieldia, Knema, Lagerstroemia, Liquidambar, Mesua, Pterospermum, Schima, Tetrameles, Wrightia

54. Fruit a pod

Acrocarpus, Adenanthera, Afzelia, Albizia, Alstonia, Archidendron, Bombax, Caesalpinia, Cassia, Dalbergia, Delonix, Dialium, Erythrina, Markhamia, Oroxyllum, Parkia, Peltophorum, Pithecellobium, Pterocarpus, Samanea, Senna, Sindora, Tamarindus, Wrightia, Xylia

55. Seeds winged

Chukrasia, Keteleeria, Lagerstroemia, Markhamia, Oroxyllum, Pinus, Pterospermum, Schima

KEYS

1. KEYS TO THE GROUPS

- 1a. Secondary wood without true vessels, ovules born on the surface of open carpels or scales, usually naked **A. Gymnosperms**
- b. Secondary wood with true vessels, ovules surrounded by carpels, forming a closed chamber **B. Angiosperms**
- 2a. Leaves compound **Group 1**
- b. Leaves simple 3
- 3a. Leaves opposite, rarely subopposite or whorled **Group 2**
- b. Leaves spirally arranged or alternate 4
- 4a. Stipules absent **Group 3**
- b. Stipules present **Group 4**

2. KEYS TO THE FAMILIES

A. GYMNOSPERMS

- 1a. Sap absent, if present not resinous. Leaves solitary 2
- b. Resin present. Leaves solitary, paired or tufted. — Pollen sacs 2. Female bracts in a spiral, seeds usually winged **Pinaceae**
- 2a. Leaves usually scale-like, decussate or whorled. Pollen sacs usually 3 or more, rarely 2. Mature female cones with 2–15 seeds. Seeds not winged or with 1–3 wings **Cupressaceae**
- b. Leaves needle-, scale-, or leaf-like, usually spirally arranged, alternate or opposite. Pollen sacs 2. Mature female cone with only 1 seed. Seeds not winged **Podocarpaceae**

B. ANGIOSPERMS

Group 1: Leaves compound

- 1a. Leaves opposite 2
 b. Leaves spirally arranged or alternate 3
- 2a. Leaves pinnate, bipinnate, or tripinnate. Fruit pod-like **Bignoniaceae**
 b. Leaves trifoliolate or palmate. Fruit a drupe or capsule **Verbenaceae**
- 3a. Stipules absent, rarely present 4
 b. Stipules present 7
- 4a. Sap present, living parts usually with aromatic or turpentine smell 5
 b. Sap usually absent, rarely present, living parts without smell 6
- 5a. Sap black when exposed to the air. Stipules absent. Living parts with turpentine smell **Anacardiaceae**
 b. Sap usually white or watery, rarely black. Stipules present or absent. Living parts usually with aromatic, resinous smell **Burseraceae**
- 6a. Bark usually smooth. True stipules absent but pseudostipules sometimes present. Stamens free. Seeds completely or partially covered by an aril . . . **Sapindaceae**
 b. Bark smooth, fissured, scaly or flaky. Stipules absent. Stamens usually partly or completely united into a tube or globose head. Seeds winged or covered by an aril **Meliaceae**
- 7a. Leaves trifoliolate, palmate, or digitate. Fruit a capsule 8
 b. Leaves usually pinnate or bipinnate. Fruit a pod 9
- 8a. Bark usually with sharp conical thorns. Sap absent. Leaves scaly. Flowers large and showy. Fruit an elliptic capsule, splitting in 5 parts **Bombacaceae** (*Bombax*)
 b. Thorns absent. Sap red. Leaves with simple hairs. Fruit a globose capsule, splitting in 3 parts **Euphorbiaceae** (*Bischofia*)
- 9a. Leaves usually pinnate, rarely bipinnate or digitate, usually without glands on the rachis. Flowers often irregular, rarely regular, usually in racemes, panicles or pseudoracemes 10
 b. Leaves usually bipinnate, often with glands on the rachis. Flowers regular, in globose heads or spikes, rarely racemously arranged **Mimosaceae**
- 10a. Flowers mostly irregular (pea-flower shaped), stamens usually 10, all united in a tube or in 2 groups (1+9) or (5+5), sometimes free **Fabaceae**
 b. Flowers regular or irregular. Stamens 10, or fewer sometimes 1, free, rarely united at the base **Caesalpiniaceae**

Group 2: Leaves simple: opposite, rarely subopposite or whorled

- 1a. Stipules present 2
 b. Stipules absent 3
- 2a. Twigs angular. Flowers large and showy, sepals connate into tube, leathery. Fruit indehiscent resting on the calyx tube **Sonneratiaceae** (*Duabanga*)
 b. Twigs terete. Flowers usually not large and showy, sepals united or free. Fruit indehiscent or dehiscent, calyx not persistent **Loganiaceae** (*Fagraea*)
- 3a. Sap present 4
 b. Sap absent 5

- 4a. Sap white. Leaves opposite or whorled, without dots. Calyx not persistent in fruit. Fruits usually paired, seeds winged or with tuft(s) of hairs at one or both ends **Apocynaceae**
- b. Sap usually yellow, rarely white. Leaves always opposite, with dots. Calyx persistent in fruit. Fruit single, seeds not winged. **Clusiaceae**
- 5a. Young twigs quadrangular. Corolla usually 2-lipped. Stamens 5, 4, or 2 **Verbenaceae**
- b. Young twigs usually terete. Corolla variously shaped, stamens 5 or more 6
- 6a. Leaves usually either with domatia or oil dots. Ovary superior or inferior. Fruit a berry, a capsule or winged, rarely a drupe 7
- b. Leaves usually without domatia or oil dots. Ovary half inferior. Fruit a capsule **Lythraceae**
- 7a. Leaves without intramarginal vein and oil glands, but often conspicuously scaly, domatia usually present. Ovary superior. Fruit usually a 2–5-winged drupe **Combretaceae**
- b. Leaves often with intramarginal vein, usually finely dotted with oil glands, not scaly, domatia absent. Ovary inferior. Fruit a berry, a capsule, rarely a drupe, never winged **Myrtaceae**

Group 3: Leaves simple: spirally arranged or alternate, stipules absent

- 1a. Sap present 2
- b. Sap absent 3
- 2a. Sap usually white. Leaves spirally arranged and frequently crowded at the end of twigs. Fruit usually a berry, rarely a drupe or capsule. Sepals persistent at base and styles persistent at apex. Aril absent **Sapotaceae**
- b. Inner bark with red sap. Leaves alternate, usually distichous. Fruit a capsule. Sepals and style not persistent. Seeds with red or orange-red, fleshy aril **Myristicaceae**
- 3a. Leaf margin serrate 4
- b. Leaf margin entire 6
- 4a. Petiole sometimes winged. Secondary veins straight, parallel toward the margin. Fruit fleshy, covered by persistent sepals **Dilleniaceae**
- b. Petiole never winged. Secondary veins curving towards the margin. Fruit a capsule. 5
- 5a. Flowers usually large and showy, petals large. Stamens numerous. Ovary superior. Fruit with persistent calyx at base and style at apex **Theaceae**
- b. Flowers usually small, petals absent. Stamens 4. Ovary inferior. Fruit with calyx persistent at apex **Datisaceae (Tetrameles)**
- 6a. Living parts usually with smell. Flowers 3-merous. 7
- b. Living parts without smell. Flowers 3–7-merous 8
- 7a. Inner bark with conspicuous wedge-shaped fibre bundles. Living parts usually with smell, but not strongly aromatic. Stamens usually numerous. Carpels many, free. Fruit monocarps **Annonaceae**
- b. Inner bark without conspicuous wedge-shaped fibre bundles. Living parts often with strong aromatic smell. Stamens 9–12. Carpels connate. Fruit a drupe, usually with persistent perianth at base **Lauraceae**

- 8a. Bark black, usually not tough or fibrous. Leaves usually with black dots on one or both surfaces. Stamens usually 10–20. Fruit a berry with enlarged sepals at base **Ebenaceae** (*Diospyros*)
- b. Bark not black, usually tough or fibrous. Leaves without black dots. Stamens 2–10. Fruit a capsule **Thymelaeaceae** (*Aquilaria*)

Group 4: Leaves simple: spirally arranged or alternate, stipules present

- 1a. Sap or resin present 2
- b. Sap or resin absent 6
- 2a. Sap present 3
- b. Resin present 5
- 3a. White sap in all parts of trees. 4
- b. Sap clear, watery. — Fruit usually 3-lobed **Euphorbiaceae** (*Aleurites*, *Endospermum*)
- 4a. Inner bark pink or red. Leaves spirally arranged and frequently crowded at the end of twigs. Fruit usually a berry, rarely a drupe or capsule. Sepals persistent at base and styles persistent at apex. **Sapotaceae**
- b. Inner bark usually cream or white. Leaves mostly alternate. Fruit united into a large fleshy compound structure **Moraceae**
- 5a. Buds often enclosed in scales. Leaves usually palmately lobed, domatia absent. Sepals and petals 4 or 5, but usually small, sometimes absent. Fruit a head. **Altingiaceae**
- b. Buds not enclosed in scales. Leaves entire, often with domatia. Sepals and petals 5. Fruits usually a 2–5-winged (from calyx) nut. **Dipterocarpaceae**
- 6a. Inner bark usually with spicy smell. Stipules large, enclosing the young bud, soon falling and leaving ring-like scars on the twigs **Magnoliaceae**
- b. Inner bark without spicy smell. Stipule not enclosing the young bud, not leaving ring-like scars on the twigs 7
- 7a. Leaves usually crowded at top of twigs. Raised glands usually present on lower surface. Stipules mainly triangular. Fruit a berry. . **Euphorbiaceae** (*Baccaurea*)
- b. Leaves usually not crowded at top of twigs. Raised glands absent on lower surface. Stipules usually linear, ovate, sometimes splitted. Fruit usually a nut or a capsule 8
- 8a. Stellate hairs or scales usually present. Bark usually fibrous 9
- b. Stellate hairs or scales usually absent, rarely present. Bark usually not fibrous 11
- 9a. Petiole often long and kneed **Sterculiaceae**
- b. Petiole usually not long and rarely kneed 10
- 10a. Bark often with sharp conical thorns. Flowers large and showy, stamens many and connate into bundles. Fruit a woody capsule, never winged. **Bombacaceae** (*Bombax*)
- b. Bark without sharp conical thorns. Flowers usually small, stamens many and free or connate only at base. Fruit a (winged) capsule **Tiliaceae** (*Berrya*)
- 11a. Stipules sword-like, free or sometimes adnate to the petiole. Fruit a drupe or pear-like 12
- b. Stipules usually small, free. Fruit a nut 13

- 12a. Stipules sword-like, free, enclosing the terminal bud. Leaf margin entire. Petiole grooved above. Stamens 10. Fruit a drupe. **Irvingiaceae**
- b. Stipules filiform or linear, sometimes adnate to the petiole, not enclosing the terminal bud. Leaf margin usually serrate. Stamens numerous. Fruit pear-like **Rosaceae** (*Pyrus*)
- 13a. Leaves usually double serrate, domatia usually present. Inner bark without tannin, but contains oil, cambium smooth. Fruit a winged nut **Betulaceae** (*Betula*)
- b. Leaves entire or serrate but not double serrate, domatia absent. Inner bark with much tannin, darkening on exposure, cambium furrowed. Fruit a nut **Fagaceae**

A. GYMNOSPERMAE

CUPRESSACEAE

Trees or shrubs, monoecious or dioecious, predominantly evergreen (deciduous in *Taxodium*), resinous and aromatic in all or some parts. *Leaves* simple, usually scale-like, decussate or whorled. *Male cones* ovoid, solitary, axillary or terminal, usually comprising only a few microsporophylls, with stamiferous scales, pollen sacs 2–7. *Female cones* terminal or axillary, solitary or in fascicles, consisting of scales with 2–15 ovules. Ripe female cones woody, or sometimes fleshy and berrylike, scales with apical tip, splitting open or not. *Seed* variable, winged or not (seed wings derived from the seed coat), resin glands present, cotyledons 2, sometimes 4, vein 1.

Distribution — Tropical, subtropical and temperate areas: 22 genera, 150 species. In Laos: 2 genera, 2 species. In Vietnam: 6 genera, 8 species.

KEY TO THE GENERA

- 1a. Upper and lower surface of leaves green, with a resinous furrowed gland at lower surface. Male cones usually with 8–12 pairs of stamens. Female cones with 6–8(–14) pairs of scales. Seeds 6–8 per scale, with 2 equal wings **1. Cupressus**
- b. Upper surface of leaves green, lower one white due to waxy stomatal depressions, gland absent at lower surface. Male cones with 14–16 pairs of stamens. Female cones with 5–8 pairs of scales. Seeds 2 per scale, with 2 very unequal wings. **2. Fokienia**

1. CUPRESSUS L.

Cupressus funebris Endl. — Fig. 1

Synonym — *Cupressus tonkinensis* Silba

Vernacular name — Hoàng đàn (V).

Evergreen, medium sized trees up to 25 m high, 40–90 cm diameter. *Bark* greyish brown, with longitudinal fissures. *Twigs* cylindrical, nearly quadrangular, two sides of leaves green. *Branches* in whorls. *Crown* large, oval in shape. *Leaves* scaly, in 4 rows, imbricate, apex obtuse, margins denticulate, with a resinous furrowed gland at lower




Fig. 1. *Cupressus funebris* Endl. (Cupressaceae).

surface. *Cones* unisexual. *Male cones* cylindrical-oblong, 5–6 mm long, pedicel of cones short, 3–10 mm long, usually consisting of 8–12 pairs of stamens with an ovate connective, and 3 or 4 pollen sacs. *Female cones* subglobose or ovate, less than 1.5 cm diam., formed by 4–8 scales, tip mucronate. *Seeds* 6–8 per scale, subglobose, c. 4 mm long, green when young, dark brown when mature, compressed laterally, sometimes triquetrous, with 2 equal wings.

Distribution — West Himalaya (Nepal, Bhutan), South China, Vietnam. An endangered species in Vietnam, only found in a narrow belt in North Vietnam, belonging to Lang Son (Huu Lung, Chi Lang), Tuyên Quang (Na Hang), and Hà Giang (Dong Van) provinces. This species does not occur in Laos.

Habitat & Ecology — Found in hot and wet tropical and subtropical forests, altitude 250–3000 m, mixed with *Burretiodendron hsienmu* and *Markhamia stipulata* on limestone mountains, sometimes forming pure stands on mountain slopes, ridges, and tops. Rather slow growing, natural regeneration very limited. *Cones*: February, March; mature ones: May, June of the next year.

Uses — Wood with straight grain and fine texture, resistant to termites and insects, with aromatic smell. Used for cabinetwork, furniture, and fine art articles. Wood, especially the root wood, contains essential oils that are used as medicine to cure inflammatory wounds, or as an antiseptic; it is also used in the cosmetic industry.

Note — Older references for Vietnamese conifers, identify *Cupressus* trees found in Lang Son and north-eastern Vietnam as *C. torulosa* D. Don. This species is restricted to the Himalayas although it is cultivated in highland areas of Vietnam and may be naturalised in some places (Lâm Đông). Natural and many cultivated trees from Lang Son are definitely not *C. torulosa*.

Specimens examined: *Cooper* 5793 (P); *Falconer* 994 (P); *Jacquemont* 1999, s.n. (P); *Pierre* 159 (P); *Strachey* s.n. (P).

2. FOKIENIA A. Henry & H.H. Thomas

Fokienia hodginsii (Dunn) A. Henry & H.H. Thomas

Synonyms — *Cupressus hodginsii* Dunn; *Fokienia maclurei* Merr.

Vernacular names — Lang len, Leng le (L); Pơ mu (V).

Trees up to 35 m high, up to 200 cm diam., bole straight, crown pyramidal. *Bark* brown grey, peeling off when young, later longitudinally fissured, aromatic. *Leaves* dimorphic, scaly, arranged in flattened tripinnate branchlet systems, the pinnae disposed in one plane, the branchlets tapering above. Leaves on adult trees arising in whorls of 4 at the same level, subacute, about 2 mm long, lateral leaves ovate, compressed, with white stomatal depressions on the ventral surfaces, facial leaves narrow obovate with a triangular apex, furrowed above. Internodes longer on older branchlets, the leaves rising at different levels in alternately opposite pairs. Leaves on young plants larger, about 8 mm long with spine-like points. *Male cones* oval to cylindrical, axillary, solitary, pedicle 5–20 mm long, cones about 7–12 mm long, with 3–5 pairs of scales, with 14–16 orbicular stamens each with 3 or 4 pollen sacs. *Female cones* globose or subglobose, 1.5–2.5 cm diam., shortly stalked, composed of 6–8 pairs of scales, each with a central spine or short process. *Seeds* 2 on each fertile scale, about 4 mm long, angular, pointed, with 2 large resin blisters on the upper and lower surface, wings lateral, very unequal.

Distribution — China (Zhejiang, Fujian, Guizhou, Yunnan) and Indochina. In Laos: Houa Phan, Xieng Khouang, and Bolikhamsai provinces. In Vietnam: Dac Lac, Gia Lai, Đông Nai, Khanh Hoa, Lâm Đông, Hà Bac, Hà Giang, Hà Tĩnh, Hoa Binh, Kon Tum, Lai Châu, Lào Cai, Nghệ An, Sơn La, Thanh Hoa, Tuyên Quang, Kiên Gian, and Yên Bái provinces.

Habitat & Ecology — *Fokienia hodginsii* is a shade-intolerant species, well adapted to mild climates with abundant rainfall, occurring naturally on humid soils in high mountain areas, slopes or plains. In Vietnam it is found at altitudes above 900 m on granite or limestone mountains, forming pure stands or mixed with *Dacrydium elatum*, *Pinus dalatensis*, and other broad-leaved tree species of the families Fagaceae, Lauraceae, and Magnoliaceae. Sufficient natural regeneration occurs in open areas, e.g. along streams, at forest edges, and in clearings in young forests. Cones: March, April; mature ones: September to December of the next year.

Uses — Wood is light, fine, and aromatic with straight grains and can be made into valuable furniture, art articles, and produces charcoal of high heat value. The distillation of wood, especially that of the roots, gives a high value essential oil, used for cosmetics.

Specimens examined: *Chevalier* 29391, 29493 (P); *Hiep et al.* 403, 1437 (P); *Krempe* 1599 (P); *Pételot s.n.* (P); *Poilane* 3462, 3521, 3539, 4397, 6500, 6527, 18742, 31056, 32545 (P); *Schmid* 858 (P), 859 (L).

PINACEAE

Normally big, monoecious trees, with resin. Branches whorled, scaly. *Leaves* usually persistent, alternate or spirally arranged, linear or awl-shaped. *Male cones* axillary, sometimes in groups at the end of twigs. *Male flowers* with 2-locular anthers. *Female cones* axillary. *Female flowers* in axils of scales and bracts. *Fruit* cone scales woody. *Seeds* usually with a large wing. Cotyledons many, above the ground.

Distribution — In all temperate regions: c. 11 genera, 200 species. In Laos: 2 genera, 3 species. In Vietnam: 4 genera, 10 species.

KEY TO THE GENERA

- 1a. Leaves single, linear, thin, flattened, twisted at base, 2–4 mm wide **1. Keteleeria**
 b. Leaves 2–5 in fascicles, needle-like, 0.5–1 mm wide, enclosed around the base by a sheath of scale leaves **2. Pinus**

1. KETELEERIA Carrière

***Keteleeria evelyniana* Mast.**

Synonyms — *Keteleeria davidiana* Beissn.; *K. delavayi* Tiegh.; *K. dopiana* Flous; *K. roulettii* (A. Chev.) Flous; *Tsuga roulettii* A. Chev.

Vernacular names — Hing (L); Du sam, Ngo tùng (V).

Big trees up to 40 m high, up to 130 cm diameter. *Bark* grey-brown scaly. *Bud* ovate, short, glabrous. *Leaves* alternate or spirally arranged, single, persistent, linear, flattened, usually (20–)30–70(–90) by 20–40 mm, base cuneate, slightly twisted, apex acute or rounded. Midrib raised above, at both sides with a white stripe of stomata. *Male flowers* in lateral or terminal fascicles. *Female flowers* in erect, solitary cones. *Cones* lateral or subterminal, ovoid, cylindrical, 12–20 by 3–6 cm, ripening every year, woody, persistent; scales rhomboid, more or less auriculate at the base, densely striate on the scales, margin denticulate. *Seeds* shiny, ovate, up to 26 mm long, with large resin cavities, winged, 15 by 10 mm, yellow-brown.

Distribution — South China, Myanmar, Thailand, Indochina. In Laos: Houa Phan, Xieng Khouang, Bolikhamsai (Khamkeut District), Khammouane, Saravane, and Champassak (Bolaven) provinces. In Vietnam: Lai Châu, Sơn La, Hòa Bình, Thừa Thiên Huế, Kon Tum (Ngọc Linh), Nghệ An, Lâm Đông (Dalat city), Đăk Lăk, Khánh Hòa, and Ninh Thuận provinces.

Habitat & Ecology — In primary, secondary forest, in moist evergreen and mixed conifer-broadleaved forest together with Fagaceae and *Alstonia* species. Sometimes near streams, in Dipterocarp forest, dry evergreen hill forests, at high altitudes of (500–)800–1600(–2000) m.

Uses — Wood is used for construction, house posts, boards and roof shingles, household furniture; the resin of the bark is used for medicine; oil of the seeds is used for burning as incense and in soap manufacturing; young shoots are used for paper-making.

Specimens examined: *Chevalier 30025* (P); *Colani s.n.* (P); *Evrard 1048* (P); *Hayata 642* (P); *Jacquet 578* (P); *Kerr 20971* (P); *Krempf s.n.* (P); *Lecomte 1542, 1584* (P); *Mieville 27070, 37068* (P); *Pételot s.n.* (P); *Poilane 1959, 16188, 20064* (P), *16188* (L); *Spire 494, 1505* (P); *Tixier 20* (P).

2. PINUS L.

Evergreen monoecious trees or rarely shrubs. *Bark* smooth or rough, particularly in older trees, peeling in flakes with very irregular shape. *Leaves* linear, often with minute serrations, spirally arranged, soon replaced by scales in the axils of which appear reduced shoots in the form of bundle-like leaves enclosed around the base by a sheath of membranous scale leaves. Adult leaves in bundles, linear, sharp, needle-like, 0.5–1 mm wide. *Pollen cones* numerous, axillary, cylindrical, subtended by a cluster of overlapping scales similar to those of the foliage bud, microsporophylls scale-like with two inverted pollen sacs. *Seed cones* terminal on short scaly shoots, more or less cylindrical, ovoid, consisting of numerous fertile scales which become woody, ripening in the second or third year, the apiculate bract fused with the scale, two inverted ovules on each scale. *Seeds* egg-shaped (ovate), expanded wing is attached to the broad base of the seed.

KEY TO THE SPECIES

- 1a. Leaves three per bundle. Bark reddish brown when old. Mature seed cone globular. **1. P. kesiya**
- b. Leaves two per bundle. Bark dark brown or blackish when old. Mature seed cone ovate. **2. P. merkusii**

1. Pinus kesiya Royle ex Gordon

Synonyms — *Pinus insularis* Endl.; *P. khasya* Hook.f.; *P. langbianensis* A. Chev.

Vernacular names — Paek sam bai (L); Thông ba lá (V).

Big trees up to 45 m high, 60–80 cm diameter. *Trunk* straight, cylindrical, resinous. *Bark* thick, reddish brown when old, deeply fissured longitudinally, breaking off in small thick irregular plates and thus sometimes becoming smoother and plate-like. *Crown* pyramidal when young, rounded when old. *Branchlets* smooth, bright brown. *Foliage buds* cylindrical and non-resinous with brown awl-shaped scales. *Needles* three together, 12–24 by 0.5 mm, acuminate, stomata on both surfaces, caducous after two years. Basal sheath 5–18 mm long, greyish brown. *Male cones* 18–30 by 5 mm.

Female cones ovoid to conical before opening, 4.5–10 by 3–5 cm, usually persistent. *Seeds* 5–8 by 3 mm, with a thin wing, 20 by 8 mm.

Distribution — East India, East Myanmar, China, Thailand, Indochina, Malaysia, Philippines. In Laos: Louang Namtha (Sing Distr.), Houa Phan, Xieng Khouang (Paek, Phoukoud, Paxay districts), Vientiane and Bolikhamsai (Phou Khao Khouay NBCA), Khammouane (Nakai), Saravane, Sekong, Champassak (Houay Ho, Bolaven Distr.) and Attopeu provinces. In Vietnam: Lai Châu, Yên Bái, Hà Giang, Dac Lac, Son La, Lang Son, Cao Bang, Quang Ninh, Kon Tum, Khanh Hoa, Ninh Thuận, Lâm Đồng (Dalat city), and Đông Nai provinces.

Habitat & Ecology — Light demanding, frost-tolerant tree, can grow on well-drained, acid, mineral soils. Usually growing in pure stands or mixed with other species in conifer forests or with broad-leaved trees, preferring high rainfall, with distinct dry and rainy seasons, air humidity not too low. Altitude (800–)1000–2300 m. Young trees are slow-growing during the first five years, later rather fast-growing. 15 years after planting the resin can be extracted. Natural regeneration is good, especially in open areas. In Laos: flowering: February, March; fruiting: January to March the year after. In Vietnam: flowering: April, May; fruiting: the second year after flowering.

Uses — The soft and light wood is commonly used as timber, in housing implements and constructions, for boxes, matches, paper pulp, board-making, window or door frames, old- and slow-growing trees which develop red heartwood are chipped and used as torches sold in the local market, used for furniture, temporary electric poles, sometimes used as firewood. Resin good but not abundant, therefore not yet economically exploited.

Note — The diameter of the branches depends on the stand density, in open forest the diameter of branches is larger than that in dense forest.

Specimens examined: *Averyanov et al. VH 162 (P); Capus 6 (P); Chevalier 38480 (P); Chinh 993 (P); Hayata 891 (P); Hennipman 3601 (L); Lecomte 1444, 1511, 1601 (P); Massie s.n. (P); Pételot 4383, 4385 (P); Poilane 1972, 2050, 3785, 3950, 4082, 4131, 15527, 15992, 16159, s.n. (P); Schmid 863, 864 (P); Spire 554 (P).*

2. *Pinus merkusii* Jungh. & de Vriese

Synonyms — *Pinus finlaysoniana* Wall. ex Blume; *P. latteri* Mason; *P. merkusiana* Cooling & Gausсен; *P. merkusii* var. *tonkinensis* (A. Chev.) Gausсен; *P. sumatrana* Jungh.; *P. sylvestris* auct. non L.

Vernacular names — Paek songbai, Paek yang, Khoua (L); Thông nhựa, Thông hai lá (V).

Big trees up to 50 m high, 60–80 cm diameter. *Trunk* straight, cylindrical, resinous. *Bark* thick, reddish when young, dark brown or blackish when old, deeply fissured longitudinally. *Crown* pyramidal with heavy horizontal branches. First year branches brownish, glabrous, without white powder. *Foliage buds* long and narrow with awl-shaped scales. *Needles* in pairs of 2, 15–28 by 1 mm, abruptly pointed, stomata on both surfaces, falling in the second year. Basal sheath 10–20 mm long, reddish. *Male cones* 18–25 by 5 mm. *Female cones* cylindrical before opening, 5–11 by 3 cm, usually falling soon after shedding seeds. *Seeds* ovate, slightly flat, 7.5 by 4.5 mm, with a thin wing 25 by 8 mm.

Distribution — Scattered throughout South East Asia from East Myanmar to South China, Malaysia, Indonesia, Philippines. In Laos: Louang Namtha (Sing Distr.), Oudomsai (Hua Namkat Distr.), Houa Phan, Louang Prabang (Chomphet Distr.), Xieng Khouang, Bolikhamsai (Khamkeut Distr.), Khammouane (Nakai), Savannakhet, Saravane (Bolaven), Sekong, Champassak (Houay Ho Distr.), Attopeu provinces, and Vientiane Municipality (Phou Khao Khouay NBCA). In Vietnam: Lai Châu, Son La, Lang Son, Bac Thai, Hà Bac, Quang Ninh, Thanh Hoa, Nghệ An (Vinh), Hà Tĩnh, Nam Hà, Thai-Nguyen, Khanh Hoa, Ninh Thuận, Đông Nai, Soc Trang, Quang Binh, Thua Thien Hue, Kon Tum, and Lâm Đông (Dalat city) provinces.

Habitat & Ecology — Light demanding, heat- and drought-tolerant tree. Generally on poor quality acid soils over sandstone or fresh volcanic ash. Most stands show a clear relationship to fire or other disturbance. Occurs in pure stands or mixed with other species in conifer forest. Altitude 800–1200(–2000) m. Young trees slow-growing during the first five years, later rather fast-growing. 15 years after planting the resin can be tapped from the trees. Natural regeneration is good, especially in open areas. In Laos and Vietnam after the Indochina war pure stands were growing in the bomb craters. In Laos: flowering: January, February; cones mature: February to May the year after. In Vietnam: flowering: May, June; cones mature: October, November the year after.

Uses — Pinewood is mainly used for house construction. Used for board-making, window or door frames, old trees are chipped for resinous wood used as torches, also for matches, paper pulp, furniture, pit props, electronic poles, ships and vehicle-building. High content of resin, each tree gives 3–4 kg of resin per year which is used for medicine, paints, printing, and in the perfume industry.

Specimens examined: *D'Alleizette* s.n. (P); *Averyanov* 2210 (P); *Balansa* 4204 (P); *Beauchaine* 107 (P); *Bon* 3162, s.n. (P); *Capus* 7 (P); *Chevalier* 29154, 29768, 29825, 30954, 37588, 37675, (P), 38231 (L); *Fleury* 37968, 38014 (P); *Hiep* 944 (P); *Lecomte* 1114, 1460 (P); *Magnen* 46 (P); *Pierre* 547 (P), s.n. (L); *Poilane* 2292, 2342, 5659, 6232, 8613, 12184, 15926 (P), 28632 (L, P); *Service Forestier* 38230, 38231 (P); *Schmid* 862 (P); *Thorel* 416 (P).

PODOCARPACEAE

Trees or shrubs, monoecious or dioecious, usually with straight trunk and more or less horizontal branches. *Leaves* simple, usually spirally arranged, sometimes opposite, alternate, scale-like, needle-like, linear to lanceolate, flat or leaf-like. *Male cones* axillary or terminal, solitary or in groups of 3–5. *Stamens* numerous, close together, imbricate, each with 2 sporangia, pollen grains usually winged. *Female cones* terminal or axillary, solitary or in fascicles; much reduced to a few fleshy bracts or scales, pendant, usually born on a thin peduncle, containing a single inverted ovule. *Seeds* wingless, completely covered by a fleshy structure referred to as an epimatium, epimatium and integument sometimes connate and forming a leathery testa. *Cotyledons* 2, with 2 parallel vascular bundles.

Distribution — In tropical and subtropical areas: 17 genera, 125 species. In Laos: c. 4 genera, c. 4 species. In Vietnam: c. 4 genera, c. 7 species.

KEY TO THE GENERA

- 1a. Leaves dimorphic. Adult leaves needle- or scale-like, less than 5 mm long. Young leaves needle-like or linear 2
 b. Leaves monomorphic. Adult leaves neither needle- nor scale-like, more than 5 mm long. Young leaves similar to adult leaves in shape, but often larger. . . . 3
 2a. Young leaves 2-ranked, linear. Adult leaves needle- or scale-like, falcate, 0.8–1.5 cm long **1. Dacrycarpus**
 b. Young leaves not 2-ranked, spreading, linear to needle-like or subulate. Adult leaves hard and scale-like, 2–5 mm long. **2. Dacrydium**
 3a. Leaves with a single, obvious, often raised midvein visible on 1 or both surfaces **4. Podocarpus**
 b. Leaves without an obvious midvein but with many, parallel veins. . . . **3. Nageia**

1. DACRYCARPUS (Endl.) de Laub.

Dacrycarpus imbricatus (Blume) de Laub.

Synonyms — *Podocarpus imbricatus* Blume; *P. kawaii* Hayata

Vernacular names — Long len (L); Thông nàng, Thông lông gà, Bành tùng (V).

Trees up to 35 m high, up to 200 cm diam., bole straight, cylindrical. *Bark* dark brown or blackish, inner bark orange with brownish resin. *Branchlets* stiff, erect. Two types of leaves present, leaves on young branchlets and young trees linear, penniformly arranged, 0.6–1.5 by 0.08–0.12 cm, stomata arranged in 2 whitish rows on lower surface, base decurrent, margin entire, apex obliquely incurved apiculate, apiculus 0.2–0.3 mm long. *Leaves* on the old branches and fruiting branches, imbricate, scale-shaped, falcate, base keeled, apex acute, 1.5–2.5 by 0.4–0.6 mm. *Male cones* ovoid or ellipsoid, axillary, 0.6–1.2 cm long. *Female cones* solitary or paired at the tip of twigs, usually one fertile, receptacle glaucous, obovoid, 3–4 by 1–2.5 mm. *Seeds* globose, 5–6 by 4–6 mm, reddish brown when ripe.

Distribution — India, Myanmar, China, Indochina, Thailand, Malaysia, Indonesia, Philippines, Papua New Guinea. In Laos: Xieng Khouang, Savannakhet, Saravane, and Attopeu provinces. In Vietnam: Tuyên Quang (Na Hang Distr.), Yên Bái, Lào Cai, Hà Bắc, Hà Giang, Sơn La, Hòa Bình, Ninh Bình, Thanh Hóa, Nghệ An, Hà Tĩnh, Kom Tum, Quang Bình, Quang Tri, Đắc Lắc, and Bình Thuận provinces.

Habitat & Ecology — Found in tropical forests, altitude 300–1000 m, usually mixed with *Altingia siamensis*, *Celtis australis*, *Cinnamomum* spp., *Gironniera subaequalis*, *Lithocarpus* spp., and *Mallotus yunnanensis*. Light demanding tree, but shade tolerant when young, prefers fertile, humid and sandy soil. Natural regeneration is good. Cones: February to April; mature ones: October to December.

Uses — Wood with fine grain, resistant to termites and insects, easy to work. Used for construction, box making, cabinetwork, furniture, and fine art articles.

Specimens examined: *Chevalier* 29391, 29493 (P); *Hiep et al.* 403, 1437 (P); *Krempe* 1599 (P); *Pételot s.n.* (P); *Poilane* 3462, 3521, 3539, 4397, 6500, 6527, 18742, 31056, 32545 (P).

2. DACRYDIUM Sol. ex G. Forst.

Dacrydium elatum (Roxb.) Wall. ex Hook.

Synonyms — *Dacrydium pierrei* Hickel; *Juniperus elata* Roxb.

Vernacular names — Long len (L); Hoàng đàn già, Dương tùng, Dương liễu, Bạnh tùng (V).

Trees up to 40 m high, up to 100 cm diam., bole straight. *Crown* a large billowy dome with tufts of more or less erect branches. *Bark* brown. *Leaves* dimorphic, spirally arranged. Young leaves and leaves on young trees awl-shaped, imbricate, 0.8–2 cm long. Leaves on twigs with cones scale-like and triangular, hard, base decurrent, apex often curved. *Male cones* solitary, terminal, cylindrical, 0.7–0.9 cm long. *Female cones* solitary or arranged in small groups, axillary or terminal, only one ovule develop. *Seeds* ovoid, 0.4–0.5 by 0.2–0.3 cm, suberect, 1/3 of the base covered by epimatium.

Distribution — India, Myanmar, China, Indochina, Thailand, Malaysia, Indonesia, and Philippines. In Laos: Xieng Khouang, Savannakhet, Saravane, and Attopeu provinces. In Vietnam: Hà Bac, Hà Giang, Hà Tĩnh, Hoa Binh, Nghê An, Son La, Thanh Hoa, Tuyên Quang, Yên Bái, Kom Tum, Quang Binh, Quang Tri, Dac Lac, and Binh Thuân provinces.

Habitat & Ecology — Found in tropical forests, altitude 700–2000 m, usually mixed with *Castanopsis* spp., *Cupressus* spp., *Dacrycarpus imbricatus*, and *Fokienia hodginsii*. Light demanding tree, prefers a cool climate, humid and yellow soil. Cones: February to April; mature ones: October, November.

Uses — Wood with fine grain, resistant to bending and pressing. Used for construction, boat and ship making, cabinet work, furniture, and fine art articles. Essential oil distilled from the wood is used as medicine, especially to treat stomach ache.

Specimens examined: *Chevalier* 29391, 29493 (P); *Forestier* 38217 (L); *Hiep et al.* 403, 1437 (P); *Krempe* 1599 (P); *Pételot s.n.* (P); *Pierre* 1396 (L); *Poilane* 3462, 3521, 3539, 4397, 6500, 6527, 18742, 31056, 32545 (P).

3. NAGEIA Gaertn.

Nageia fleuryi (Hickel) de Laub.

Synonyms — *Decussocarpus fleuryi* (Hickel) de Laub.; *Podocarpus fleuryi* Hickel

Vernacular names — Kim giao, Kim giao núi, Báng sủng (V).

Medium sized trees up to 20 m high, 80 cm diameter. *Trunk* straight. *Crown* pyramidal, branches horizontal or slightly pendulous. *Bark* greyish brown, peeling in large thin irregular shaped plates with scattered lenticels. *Foliage buds* a compact cluster of lanceolate deciduous scales abruptly wider than the shoot and distinctly acute. *Twigs* green. *Leaves* simple, opposite or subopposite, distichous, elliptic or broadly lanceolate, 8–20 by 3–6 cm, thick and leathery, base acute, margin entire, apex acute or acuminate, venation parallel. *Petiole* twisted, 5–10 mm long. *Male cones* cylindrical, solitary or in groups of 3–6, axillary, sessile, 2–3 cm long. *Female cones* solitary, axillary, ovules 1–3, sessile in axils of subterminal bracts, only 1 ovule maturing; peduncles 1.5–2 cm long; mature cones globose, 1.5–2.5 cm diam., brownish violet, green when young.

Distribution — South China and Indochina. In Laos: Vientiane province. In Vietnam: North and Central Vietnam, in Hai Phong (Cat Na NP), Ninh Binh (Cuc Phong NP), Thanh Hoa, and Nghệ An provinces.

Habitat & Ecology — Tropical forests, altitude 200–1000 m, usually mixed with *Madhuca pasquieri*, *Quercus bambusaefolia*, and *Vatica odorata*. Light-demanding tree, rather slow growing, natural regeneration very limited. Cones: March, April; mature ones: October, November.

Uses — Wood white with fine texture, resistant to termites and insects. Used for cabinetwork, furniture, fine art articles, and chopsticks. Leaves used as medicine to cure cough. The shape of the tree is beautiful, used as an ornamental in parks, along avenues and at pagodas and temples.

Note — *Nageia fleuryi* is an endangered tree species of Vietnam, depleted from its former numbers because of the demand for its timber. It should be strictly protected at Cat Ba and Cuc Phuong National Parks. It is listed (as *Podocarpus fleuryi*) as threatened species in Vietnam.

Specimens examined: *Chevalier 37512* (P); *Clemens 4190* (P); *Fleury 38017* (P); *Poilane 29808* (P); *Tixier s.n.* (P); *Vidal 624* (P).

4. PODOCARPUS L'Hér. ex Pers.

Podocarpus neriifolius D. Don — Fig. 2

Synonyms — *Nageia neriifolia* (D. Don) Kuntze; *Podocarpus annamensis* N.E. Gray

Vernacular names — Ka dong (L); Thông tre, Thông trúc đào, Bạnh niên tùng, Thông tre nam (V).

Trees up to 35 m high, up to 100 cm diam., bole straight. *Bark* greyish brown, peeling off in longitudinally flakes. *Branches* spreading or ascending. *Foliage bud* scales erect, triangular, 1–1.5 mm wide, apex acute. *Petiole* 3–6 mm long. *Leaves* alternate, linear or lanceolate, usually slightly curved, 2.5–18 by 0.5–1.5 cm, leathery, midvein raised on beneath, flat or slightly raised on above, base wedge-shaped, apex long acuminate. Juvenile leaves wider, with obtuse to mucronate apex. *Cones* ovoid, 0.8–1.6 cm high. *Male cones* solitary or in clusters of 2–4, normally sessile, 2–4.5 cm long, with several spirally arranged, basal bracts. *Female cones* solitary, axillary, peduncle 0.4–2.2 cm long. *Receptacle* orange red when ripe, obconical ellipsoid, 4–10 by 2–6 mm, base with 2 subulate bracts, 2–6 mm long. *Mature seed* green to purple, outer layer fleshy to coriaceous, middle layer stony, and inner layer papyraceous.

Distribution — India, Nepal, Myanmar, Indochina, China, Thailand, Malaysia, Indonesia, Philippines, Papua New Guinea. In Laos: Houa Phan, Xieng Khouang, and Bolikhamsai provinces. In Vietnam: North and Central Vietnam, such as Hà Bac, Hà Giang, Hà Tĩnh, Hoa Binh, Lai Châu, Lào Cai, Nghệ An, Son La, Thanh Hoa, Tuyên Quang, Kien Gian, and Yên Bái provinces.

Habitat & Ecology — Found in tropical forests, altitude 400–1500 m, usually mixed with *Castanopsis* spp., *Dacrycarpus imbricatus*, *Fokienia hodginsii*, *Lithocarpus* spp., and *Pometia* species. Shape tolerant tree, prefers fertile, humid, and humus rich soil. Sometimes it regenerates by seed under dense forest cover. Cones: April, May; mature ones: August to November.



Fig. 2. *Podocarpus neriifolius* D. Don (Podocarpaceae).

Uses — Wood with fine texture, resistant to termites and insects. Used for construction, boat making, cabinetwork, furniture, and fine art articles. The shape of the tree is beautiful, used as an ornamental tree.

Specimens examined: *Averyanov et al.* 972 (P); *Bodeneunc* 36731 (P); *Chevalier* 36585, 38693, 39241 (P); *Clemens* 3475 (P); *Evrard* 336 (L, P), 1101, 2390 (P); *Gray* 48 (P); *Pierre* 354 (P), 5532 (L, P); *Poilane* 1561, 3541, 6532, 21828, 24178, 24439, 29031, 32515, 32566, 35674 (P), 24443, 30856, 30934 (L, P).

B. ANGIOSPERMAE

ALTINGIACEAE

Trees or shrubs with aromatic resin in bark and wood. *Buds* often enclosed in scales. Stellate or tufted hairs or scales frequently present. *Stipules* present (except *Rhodoleia*). *Leaves* simple, alternate, spirally arranged, often palmately lobed, entire to serrate, venation pinnate or palmate. *Flowers* small, in small clusters, spikes or heads, bisexual, polygamous or unisexual, monoecious, rarely dioecious. *Sepals* and *petals* 4 or 5, latter often small or absent. *Stamen(s)* 1 or more, in 1 whorl. *Ovary* superior, half-inferior or inferior, carpels 2, often free at the apex, 2-locular, ovule(s) 1—many per locule. *Styles* 2, slender. *Stigmas* 2, apical or adaxially. *Fruits* forming globose

heads, woody capsules, 2–4-valved, often tipped by persistent styles. *Seed(s)* 1–many, small, sometimes winged.

Distribution — Africa, Australia, but mainly in Asia: c. 28 genera, c. 130 species. In Laos: c. 6 genera, 6 species. In Vietnam: c. 7 genera, 16 species.

KEY TO THE GENERA

- 1a. Aromatic resin present or absent. Leaves elliptic or obovate, base usually obtuse, stellate hairs absent, venation pinnate. Male inflorescences in globose heads **1. *Altingia***
- b. Aromatic resin always copious. Leaves 3–5-lobed, base cordate, stellately hairy when young, venation palmate. Male inflorescences in spikes . . . **2. *Liquidambar***

1. *ALTINGIA* Noroña

***Altingia siamensis* Craib — Fig. 3**

Synonyms — *Altingia gracilipes* Hemsl.; *A. takhtajanii* Thai

Vernacular names — Sop (L); Tô hạp, Tô hạp điện biên, Mây xa hon (V).

Trees up to 45 m high, up to 1 m diameter. *Trunk* straight, cylindrical. *Bark* greyish pink, smooth, splitting into large pieces when old. *Stipules* caducous. *Petioles* slender, 2–3 cm long. *Leaves* simple, alternate, obovate, 5–8 by 2–3 cm, base obtuse, margin slightly roundish serrate, rarely entire, apex long pointed, venation pinnate, evident at lower surface, secondary veins 5–7 pairs. Leaves red when old. *Flowers* unisexual, monoecious. *Male inflorescences* in globose heads, sepals and petals absent, stamen 1, filaments short. *Female flowers* in globose heads, sepals scale-like, petals absent, ovary half inferior. *Fruits* forming globose heads, woody capsules, style persistent. *Seed(s)* very tiny, oblong, winged.

Distribution — India, Myanmar, Thailand, Indochina. In Laos: Vientiane province. In Vietnam: Lai Châu, Lào Cai, Sơn La, Gia Lai, Kon Tum, Đăk Lăk, and Ninh Thuận.




Fig. 3. *Altingia siamensis* Craib (Altingiaceae).

Habitat & Ecology — Primary and secondary forest, altitude below 700 m. Light demanding tree, growing very well in deep soils along streams. Usually mixed with *Gironniera subaequalis*, *Machilus* spp., *Phoebe* spp., and *Quercus* species. Natural regeneration is good at low forest cover, along streams. Flowering: February, March; fruiting: June to August.

Uses — Wood resistant to termites and insects. Used for construction, ship and boat building. Essential oil from resin is used in the perfume industry.

Note — Tardieu-Blot in Lecomte, Flore du Cambodge, du Laos et du Vietnam 4 (1965) 98–99 considered *Altingia takhtajanensis* Thai (= *A. takhtajanii* Thai) and *Altingia siamensis* Craib different species. Dung (1996) considered *Altingia takhtajanii* Thai a synonym of *Altingia siamensis* Craib. After examining the available specimens and literature we believe that *Altingia takhtajanii* Thai is indeed a synonym of *Altingia siamensis* Craib as no major differences can be found.

Specimens examined: *Kerr* 21255 (L, P); *Maxwell* 74886 (L); *Phon* 1147 (P); *Pierre s.n.* (P).

2. LIQUIDAMBAR L.

Liquidambar formosana Hance — Fig. 4

Synonyms — *Liquidambar acerifolia* Maxim.; *L. maximoviczii* Miq.; *L. tonkinensis* A. Chev.

Vernacular names — Sau sau (V), Chao, Cha phai (Mường, V).

Trees up to 40 m high, up to 1.5 cm diameter. *Trunk* straight, cylindrical. *Bark* blackish brown, thick, longitudinally fissured, aromatic resin copious. *Twigs* slender. *Stipules* needle-shaped, 1–2 cm long, caducous. *Petioles* 8–10 cm long. *Leaves* simple, alternate, 3- or 5-lobed, usually 3-lobed, base cordate, margin serrulate, stellately hairy when young. *Flowers* unisexual, monoecious. *Male flowers* in spikes, sepals and petals absent, stamen 1. *Female flowers* in globose heads, sepals scale-like, petals absent, staminodes 4–10, ovary half inferior, 2-locular, ovules many per locule, styles 2, elongated, stigmas bifid. *Fruits* in globose heads, 2–3 cm diam., formed by many capsules, spiny from the persistent styles and calyx. *Seed(s)* very tiny, winged.

Distribution — China and Vietnam. In Laos so far not yet recorded, but most probably present. In Vietnam widely distributed in the north, mainly in Hoa Binh, Hà Bac, Hà Tây, Bac Thai, Lang Son, Cao Bang, Quang Ninh, Ninh Binh, Thanh Hoa, and Nghệ An provinces.




Fig. 4. *Liquidambar formosana* Hance (Altingiaceae).

Habitat & Ecology — Primary and secondary forest, altitude below 600 m. Usually mixed with *Canarium album*, *Engelhardtia chrysolepsis*, and *Erythrophloeum fordii*. Shade tolerant when young, light demanding when mature, can grow on poor soils. Natural and coppice regeneration is very good. This species is a pioneer species after forest fire and in shifting cultivation areas. Flowering: March, April; fruiting: October to December.

Uses — Wood light, but strong after being processed. Used for construction, railway sleepers, boat- and ship-building, furniture. Resin used in production of perfume and medicine. Leaves used as vegetable.

Specimens examined: *Balansa* 1157 (P), 3170 (P); *Bon* 5118, 5394 (P); *Boom* 2406, 10755, 11885, 13166 (L); *Butreau* 35 (P); *Cadays* 38263 (P); *Chevalier* 29654, 30170, 37299, 40980 (P); *Cuong* 93 (L); *Dodo s.n.* (P); *Domat* 38201 (P); *Faurie* 45 (P); *Pételot* 5747 (L).

ANACARDIACEAE

Trees or shrubs, rarely climbers or epiphytic shrubs, usually with acid, often turpentine smell, sap becoming black when exposed to the air. *Buttresses* sometimes present. *Bark* smooth, cracked, fissured, scaly, flaky or dipped, sometimes blotched by black exudate. Inner bark pinkish or yellowish, turning black or brown. Sapwood whitish. *Stipules* absent. *Petioles* often thickened at the base. *Leaves* often crowded at the end of twigs, usually compound, often imparipinnate, less frequently simple or paripinnate, alternate, margin usually entire. *Inflorescences* mostly large, terminal or axillary panicles, rarely racemes or spike-like, bracts and bracteoles usually caducous, sometimes persistent. *Flowers* generally small, regular, bisexual or unisexual, then plants polygamous, monoecious or dioecious. *Sepals* 4 or 5, free or connate. *Petals* absent or 4 or 5, free, valvate or imbricate. *Stamens* 8–10—numerous, sometimes partly staminodal, inserted on the disc. *Disc* mostly well-developed. *Ovary* superior, 2–5(–12)-celled, rarely inferior, ovule 1 per cell, stigma(s) 1–5(–12)-lobed or entire; placentation axillary, but mostly seemingly parietal by abortion of locules. *Fruit* a drupe or nut, fleshy and resinous.

Distribution — Throughout the tropical regions of the world: c. 75 genera, 850 species. In Laos: c. 13 genera, c. 23 species. In Vietnam: c. 15 genera, c. 55 species.

KEY TO THE GENERA

- 1a. Bole not fluted, inner bark brown. Leaves without sour taste, domatia absent. Flowers dark red, polygamous. Seeds with 5 holes at the top. **1. Choerospondias**
- b. Bole often fluted, inner bark pink. Leaves with sour taste, domatia present. Flowers green-yellow or white, bisexual. Seeds without holes at the top. **2. Dracontomelon**

1. CHOEROSPONDIAS Burt & Hill

Choerospondias axillaris (Roxb.) Burt & Hill — Fig. 5

Synonyms — *Spondias acuminata* Gamble; *S. axillaris* Roxb.; *Poupartia fordii* Hemsl.

Vernacular names — Mu (L); Xoan nhừ, Lát xoan, Xuyên cóc (V).

Deciduous trees up to 20 m high, 30–40 cm diam., buttresses present. *Bark* grey-brown, cracked and peeling in vertical flakes, inner bark pink. *Petioles* 4–8 mm long, 4–7 pairs of leaflets. *Leaves* spirally arranged, becoming red before falling, imparipinnately compound, alternate, 30–40 cm long, main stalk 5–12 cm long, grooved. *Leaflets* opposite, 4–12.5 by 2–6 cm, elliptic, base oblique, tips tapering, young leaves with scattered teeth, mature ones usually entire, venation pinnate, 6–16 veins per side, domatia absent. *Inflorescences* panicles, flowers dark red, polygamous, male flowers in large branched clusters at the end of twigs, bisexual flowers in small groups, sepals connate, 2 mm long, 5-lobed, dark red glabrous outside, glandular-hairy inside, petals 5, pointed, glabrous, imbricate, stamens 10, alternating with disc lobes, ovary superior, 5-celled. *Fruit* an ovoid or globose drupe, 1.5–2.5 cm diameter. *Seeds* with 5 holes at the top.

Distribution — India, China, Japan, Indochina. In Laos: Xieng Khouang and Houa Phan (Sam Nua Distr.) provinces. In Vietnam: from the north to Nghê An province in Central Vietnam.

Habitat & Ecology — A deciduous fast growing tree in secondary forests, altitude below 600 m. This species occurs mainly on wet, deep soils or on heavy clay or sandy clay soils. Flowering: April to June; fruiting: July to September.

Uses — Wood soft, used for general furniture. Fruits edible, also used for wine production. Bark fibrous, can be used for making rope, the pericarp is used as medicine, to promote the flow of blood.

Specimens examined: *Chevalier* 29660 (P); *Cuong* 257 (L); *Fleury* 30155, 32537 (P); *Hoi* 412 (P); *Pételot* 5441 (P); *Poilane* 2083 (L, P); *Smitinand* 1740 (L); *Soejarto et al.* 11616 (L); *Sukkri* 32 (L); *Tsang* 29308 (P); *Van Beusekom* 440, 4325 (L).

2. DRACONTOMELON Blume

Dracontomelon dao (Blanco) Merr. & Rolfe — Fig. 6

Synonyms — *Dracontomelon duperreanum* Pierre; *D. mangiferum* Blume; *D. sinensis* Stapf

Vernacular name — Sầu (V).

Large deciduous trees up to 35 m high, 1–1.2 m diam., usually with large buttresses. *Bark* grey brown, splitted into pieces, inner bark pink, latex absent, sometimes gum



Fig. 5. *Choerospondias axillaris* (Roxb.) Burt & Hill (Anacardiaceae).

drops present. *Twigs* with large leaf scars, twigs, buds and bases of the leaf-stalks light fawn-brown, finely hairy. *Leaves* spirally arranged, imparipinnate compound, clustered near tips of twigs, more or less hairy, at least on the stalk and on the underside of the veins. *Leaflets* alternate or opposite, 4–9 pairs, leaf axis 30–50 cm long, main stalks 8–10 cm long, grooved at base; leaflets narrow oval with oblique bases, tapering apex, margin entire, apex acute, lower leaflets 5–9 by 2–4 cm, upper leaflets 10–15 by 3–5 cm, venation pinnate, veins 7–14 pairs, usually with fulvous hairs at axils (domatia), upper surface glabrous or subglabrous, lower surface glabrous or pubescent, petiolules 3–4 mm long. *Inflorescences* a terminal or axillary panicle, up to 60 cm long, pubescent. *Flowers* minute, yellowish green or white, bisexual; sepals 5, fused at base, triangular, hairy outside but glabrous inside; petals 5, narrow and pointed, slightly longer than stamens and style, glabrous. *Stamens* 10, the ones opposite sepals longer than those opposite petals. *Ovary* superior, pubescent at base, 5-celled, styles 5. *Fruit* a globose




Fig. 6. *Dracontomelon dao* (Blanco) Merr. & Rolfe (Anacardiaceae).

drupe, 2–4 cm diam., green turning yellowish, large with the 5-lobed calyx at the base. Stone very hard, flattened at the base.

Distribution — From India to the Solomon islands. This species occurs from North to Central Vietnam in Hà Tĩnh, Nghệ An, Thanh Hoa, Kien Gian, Hoa Binh, Ninh Binh, Lang Son, and Hà Tây provinces.

Habitat & Ecology — This species prefers wet and very light conditions in primary and secondary forests. In Vietnam it is often mixed with *Aglaia gigantea* and *Parashorea chinensis*. Flowering: April, May; mature fruits: June to September.

Uses — The wood is used for construction and making furniture. Leaves and fruit are used for cooking, mature fruits are edible and used for dental treatment.

Specimens examined: *Balansa* 3424, 3427, 3428, 4378, 4401, 4604 (P); *Blanche* 4339 (P); *Chevalier* 37292, 37296 (P), 38440 (L, P); *Cuong* 276 (L); *Fleury* 37973 (L, P); *Forestier* 38440 (P); *Himavong* 32 (L); *Kerr* 12110, 12802, 14196, 14834, 15663 (P); *Pételot* 3978 (P); *Pierre* 1623 (P); *Sangkachand* 1563 (L); *Schmid* 1156 (P); *Smitinand* 11017 (L); *Soejarto et al.* 10953 (L).

ANNONACEAE

Trees, shrubs, or climbers with aromatic tissues. *Crown* typically narrowly conical with a single main stem and horizontal twigs. *Bark* usually pale grey, buff-brown or orange-brown, smooth or elongate shallowly fissured; inner bark with conspicuous wedge-shaped fibre bundles. *Twigs* often hairy when young, and smooth when mature. *Stipules* absent. *Leaves* simple, alternate, often planar or drooping, margin entire, with short stalks. *Flowers* axillary or opposite the leaves or on old leafless stems, often pendulous. *Sepals* 3, valvate or imbricate, usually smaller than petals, pubescent or tomentose outside, glabrous inside. *Petals* 6 in 2 whorls, the inner whorl sometimes arching inwards and covering the sexual organs like a hood. *Stamens* minute, usually numerous, densely clustered in a central circular clump surrounding the ovaries. *Ovaries* usually many, free, pubescent, style present or absent. *Monocarps* fleshy, generally indehiscent, usually in clusters radiating from a common woody stalk. *Seeds* with ruminant endosperm.

Distribution — A large family found throughout the tropics: 130 genera, 2300 species. In Laos: c. 15 genera, 100 species. In Vietnam: c. 27 genera, 168 species.

KEY TO THE GENERA

- 1a. Leaves ovate-elliptic. Petals elliptic. Monocarps elliptic or obovoid, black when mature 2
- b. Leaves elliptic. Petals ovate. Monocarps round or ovoid, red when mature 2. **Polyalthia**
- 2a. Leaf margins strongly wavy, base strongly asymmetric. Inner and outer whorls of petals spreading. Connective of stamens with a narrowly elliptic acute apex. Monocarps elliptic, symmetrical or slightly asymmetrical developed, stalks long 1. **Cananga**
- b. Leaf margins flat or slightly wavy, base symmetric. Inner whorl of petals arching at base. Connective of stamens with a truncate dilate apex. Monocarps obovoid to cylindrical, strongly asymmetrical developed, stalks short 3. **Xylopia**

1. CANANGA Hook.f. & Thomson

Cananga odorata (Lam.) Hook.f. & Thomson — Fig. 7

Basionym — *Uvaria odorata* Lam.

Vernacular names — Ka dan nga (L); Ngọc lan tây (V).

Evergreen medium unbuttressed trees up to 30 m high. *Trunk* straight. *Bark* grey or silvery, smooth. *Young twigs* minutely pubescent, becoming glabrous, dark, striate. *Crown* with branches drooping, or slightly erect with dangling leafy twigs. *Leaves* ovate to broadly elliptic, 5–20 by 3–10 cm, base rounded or heart-shaped, margin entire, but strongly wavy, apex acuminate, dark green above, green beneath, secondary veins 8 or 9 pairs, clearly visible on both sides, often with domatia, pubescent or glabrous on midrib and veins on both sides. *Flowers* hanging in clusters from the older branches, greenish turning light dull yellow, very fragrant. *Sepals* triangular. *Petals* elliptic, 5–9 by 0.5–1.5 cm, with a purple or red spot at the base inside. *Stamens* many, connec-



Fig. 7. *Cananga odorata* (Lam.) Hook.f. & Thomson (Annonaceae).

tive with a lanceolate acute apex. *Carpels* many, ovules many. *Monocarps* in clusters, symmetrical or slightly asymmetrical, elliptic in outline, 1.5–2.5 by 1–1.5 cm, green when young becoming blackish when mature, stalks c. 1.5 cm long, *Seeds* 2–12 in two rows, pale brown.

Distribution — Very widespread from India all over Malesia to Fiji. In Laos: Vientiane and Champassak provinces. In Vietnam: Hà Tây (Sontá, Mt Ba Vi), Hoa Binh, Hà Nội, and Hồ Chí Minh provinces.

Habitat & Ecology — Common in secondary forest and on forest edges, abundant in villages where often dwarf forms are cultivated.

Uses — The plants are used for ornamental purposes and the flowers produce Ylang-Ylang essential oil which is an important ingredient in the perfume industry.

Specimens examined: *D'Alleizette* 81 (L); *Chevalier* 29766, 37750, 37801 (P); *Counillon s.n.* (P); *Eberhardt* 3426 (P); *Germain* 65 (P); *Hiep* 13 (P); *Pierre* 1744 (P); *Poilane* 27489 (P); *Svengsuksa* 32 (P); *Thorel* 2126, 2136 (P).

2. POLYALTHIA Blume

***Polyalthia cerasoides* (Roxb.) Bedd.**

Vernacular names — Chan dong, Ka dan nga paa (L); Duôi trâu, Nhọc, Nhọc lá bé (V).

Medium deciduous trees up to 20 m high, 20–50 cm diameter. *Trunk* straight. *Bark* greyish or pale brown, inner bark yellow, aromatic. *Young twigs* slender and yellowish tomentose, old twigs glabrous. *Leaves* ovate, ovate-elliptic, 7–12 by 5–6 cm, base rounded or obtuse, margin slightly wavy, apex acuminate, secondary veins 7–9 pairs, glabrous above, pubescent beneath. *Flower* aromatic, solitary or in clusters of 2 or 3 flowers on short woody shoots, axillary, light green, stalks long and slender, 1.2–2.5 cm, bracts 2 or 3. *Sepals* ovate or triangular, 0.6–0.9 cm long. *Petals* thick, ovate, the 3 outer petals 1.3 by 0.7 cm, inner ones narrower and shorter. *Stamens* numerous, connective disc-like. *Carpels* many, tomentose, same length as stamens. *Stigma* head-shaped, tomentose. *Monocarps* in clusters, fruitlet round or ovoid, 8 by 5 mm, stalk long, young fruits green later yellow and red when mature, silky. *Seed* one.

Distribution — India, China, Indochina. In Laos: Louang Namtha, Sayabouri, Louang Prabang, Bolikhamsay, Saravane, Savannakhet, and Attopeu provinces. In Vietnam: Hoa Binh, Thanh Hoa, Ninh Thuận, Bac Giang, Nghệ An, Hà Tĩnh, Gia Lai, Kon Tum, Dac Lac, Nhatrang, and Binh Thuận provinces.

Habitat & Ecology — Tropical rain forest, in bushy savannah, edges of secondary forest, altitude below 700 m, previously logged forest with an open canopy, and little disturbed closed canopy forest with *Camellia*, *Lagerstroemia*, and *Liquidambar* association. Flowering: January to March; fruiting: April to July.

Uses — Timber yellowish or greyish, annual rings obscure, texture fine, wood hard and heavy, usually splitting and curving when dry. Used in construction and for making agricultural tools. Used as traditional medicine.

Specimens examined: *Averyanov* 1024, 1038 (P); *Fleury* 38951 (L); *Harmand* 1169 (P); *Pierre* 469, 496, 1756 (P); *Poilane* 14, 140, 2779, 3195, 4854, 5588, 6034, 6064, 6090, 6256, 6356, 8537, 9808, 9849, 13721, 16627, 17406, 17809, 17863, 26292, 26295 (P); *Soejarto et al.* 11408 (L); *Thorel* 3167 (P); *Vichit Lamxay et al.* 256 (L).

3. XYLOPIA L.

Xylopia vielana Pierre

Vernacular names — Sa thang (L); Dền, Sai, Thối ruột (V).

Medium, deciduous trees up to 20–25 m high, 20–30 cm diameter. *Trunk* straight, cylindrical. *Bark* dark orange, easily separated from wood, with many tough fibrous layers. Inner bark pale brown. *Young twigs* slender, green, covered with fulvous tomentose hairs. *Petiole* fulvous hairy, 5–8 mm long. *Leaves* broadly ovate, 6–10 by 3–5 cm; base rounded, margins flat or slightly wavy, apex acuminate, secondary veins 7–12 pairs, evident beneath, both surfaces of leaves finely yellow hairy. *Flowers* solitary or in clusters with 1–3 flowers at axils of leaves near the end of twigs, flower buds pale green, yellow, aromatic. *Sepals* ovate or triangular, 4–5 mm long. *Petals* inner whorl arching at the base, outer whorl spreading, yellow tomentose outside. *Stamens* numerous in many whorls, connective with a truncate widely apex. *Staminodes* similar to fertile ones without anther sacs. *Receptacle* concave. *Carpels* few, tomentose. *Monocarps* in clusters, 4–6 fruitlets, obovoid to cylindrical, 3–4 by 1–1.5 cm, arranged on a globose receptacle, peduncle 1.5–3.5 cm long, strongly irregular, stalks short, green when young, slightly hairy, mature orange-yellow, black when dry, indehiscent. *Seeds* 2–5, black.

Distribution — India, Sri Lanka, China, Thailand, Indochina. In Laos: Vientiane, Bolikhamsay, and Champassak provinces. In Vietnam: Tuyên Quang, Bac Thai, Quang Ninh, Hà Bac, Hà Tây, Thanh Hoa, Nghệ An, Đông Nai (Bien hoa city), Tây Ninh, and Khanh Hoa provinces.

Habitat & Ecology — Tropical lowland rain forest, or secondary forest. Usually mixed with *Acronychia laurina*, *Engelhardtia*, and *Ormosia balansae*. Flowering: February to July; fruiting: April to November.

Uses — Timber slightly yellow, aromatic, used for common furniture, fuelwood, and mining poles. Bark and leaves used as tonic for the elderly and women after giving birth.

Specimens examined: *D'Alleizette s.n.* (P); *Balansa* 4165, 4169 (P); *Eberhardt* 4343 (P); *Gourgand* 4, 6, 8, 11 (P); *Hahn s.n.* (P); *Harmand* 646 (P); *Muller* 751 (P); *Perry* 2021 (P); *Pierre* 73, 205, 1759, 2020, 2021 (P); *Pirey* 41217 (P); *Poilane* 4638, 6866, 7454, 7773, 15157, 23620 (P); *Soejarto et. al.* 11410, 11451 (L); *Thorel* 2078 (P); *Vidal* 939, 4457 (P).

APOCYNACEAE

Small to large trees, climbers, rarely herbs, milksap present. *Stipules* absent. *Leaves* simple, mostly opposite, sometimes whorled, rarely spirally arranged. *Inflorescence* cymes or cymose panicles, rarely fasciculate or solitary, terminal or axillary. *Flowers* regular, rarely irregular, large, bisexual. *Sepals* tubular, often with colleters inside. *Petals* tubular, overlapping in bud. *Stamens* inserted on the corolla tube, completely included or exerted, anthers sagittate or ovate, free or adnate to the stigma, sometimes with the base and apex sterile. *Disc* present or absent. *Ovary* superior, rarely semi-inferior, with 2 separate carpels united into a common style, with one or two locules, pistil head-like. *Fruits* usually paired, a drupe, berry, capsule, or follicle. *Seeds* simple, aril absent, winged, with a ciliate margin or often with tuft of hairs at one or both ends.

Distribution — Worldwide: c. 500 genera, c. 4500 species (incl. Asclepiadaceae). In Thailand: 36 genera, 118 species. In Laos: 30 genera, 65 species. In Vietnam: c. 36 genera, 109 species.

KEY TO THE GENERA

- 1a. Leaves whorled, 4–8, glabrous, secondary veins 20–45 pairs, straight to the margin.
Stamens inserted in upper half of corolla tube **1. *Alstonia***
- b. Leaves opposite, tomentose, secondary veins 8–15 pairs, curved to the margin.
Stamens inserted at corolla mouth **2. *Wrightia***

1. ALSTONIA R.Br.

***Alstonia scholaris* (L.) R.Br.** — Fig. 8

Synonym — *Echites scholaris* L.

Vernacular names — Tin pet (L); Sũa, Mò cua (V).

Medium-sized to big deciduous trees 10–50(–60) m high, 20–80 cm diam.; trunk straight and often fluted at base. *Bark* smooth, scaly or shallowly fissured and peeling off in rectangular flakes, brown; inner bark granular, creamy, brown to yellow with white sap. *Branches* smooth or sparsely rough, scaly, with sparse or dense lenticels; branchlets terete, glabrous, grey-yellowish brown or sometimes dark brown, crown pagoda-shaped. *Leaves* in whorls of 4–8; petiole 5–20 mm long, glabrous, coriaceous, flattened and shallowly grooved above, slightly winged; colleters narrowly triangular and densely packed together in the axils; blade narrowly elliptic, obovate, 4.4–32 by 1.2–8.7 cm, glossy and dark green above, pale or grey-green beneath, subcoriaceous when dry, base cuneate, apex acute, obtuse to rounded, or rarely retuse. Secondary veins 20–45(–55) pairs, straight and parallel to the margin, rarely close together or branched, forming an angle of 80–90° with the midrib, venation reticulate, prominent above, glabrous. *Inflorescences* terminal cymes, many flowers in branched clusters (compound umbels) grouped at the end of twigs, 3–10 cm long, axes finely hairy; peduncle 1.5–6 cm long, glabrescent or pubescent; bracts and bracteoles scale-like, ovate or narrowly ovate, acuminate or sometimes trilobed or irregularly lobed at the apex, very shortly ciliate. *Flowers* fragrant; pedicel more or less 1 mm long. *Sepals* pale green, connate at the base, bell-shaped, with 5 teeth, ovate, 1–2.4 by 0.8–2.5 mm, apex acute to rounded, pubescent, ciliate. *Petals* white-yellow or cream, 5-lobed, overlapping to the left in bud, elliptic, 1.7–4.5 by 2–4.5 mm, tube 5.3–10 mm long; outside pubescent in the top of tube and lobes, inside pubescent except at the base of tube. *Stamens* 5, inserted in upper half of corolla tube or below the mouth of the corolla tube, not projecting; filaments 0.5–0.7 mm long; anthers ovate, 0.9–1.2 by 0.4–0.6 mm, apex obtuse; disc very small and annular or absent. *Ovary* superior, ovoid or broadly ovoid, 0.9–1.5 by 0.7–1 mm, carpels 2, pubescent, style (incl. pistil head) 2.8–8 mm long. *Fruit* a pair of follicles, 20–56 by 0.2–0.3 cm, glabrous, splitting into 2 sections with many seeds. *Seeds* dark brown, elliptic, ends rounded, 3–7 by 1.5–2 mm, margin slightly thickened. Tufts of cilia at both ends, longest cilia 8–18 mm long, becoming much shorter or glabrous towards the margins, hilum linear, 0.7–1 mm long.




Fig. 8. *Alstonia scholaris* (L.) R.Br. (Apocynaceae).

Distribution — Pakistan, India, Sri Lanka, Nepal, Bhutan, Bangladesh, Myanmar, China, Cambodia, Thailand, Malesia, Australia, Solomon Island. In Laos: Vientiane Municipality. In Vietnam: from north to south.

Habitat & Ecology — In evergreen, deciduous or mixed forest, in scrub or at the edge of forest, on wet places, along streams or rivers, altitude 50–1200 m. Easily dispersed by seeds. Strong coppice-regeneration. Flowering: August to September; mature fruiting: January, February.

Uses — The bark is used to treat malaria and male genital pains. The sap can be applied to open sores to keep maggots out. The wood can be used as plywood, for boxes, coffins, frames, furniture, and handicrafts. Trees can be planted for ornamental purposes providing excellent shade.

Specimens examined: *Balansa* 613, 4729 (P); *Bon* 6190 (P); *Chevalier* 36502 (P); *Clemens* 3869 (P); *Cuong* 905 (L); *Donnat* 38137 (P); *Fleury* 30177, 37764, 38009 (P); *Lam* 27 (P); *Muller* 725 (P); *Pételot* 5985 (P); *Pierre* 2093, 4410 (P); *Poilane* 771, 1223, 18859 (P); *Spire* 507 (P); *Tsang* 27098, 27483 (P).

2. WRIGHTIA R.Br.

Wrightia pubescens R.Br.

Synonyms — *Wrightia annamensis* Eberh. & Dubard; *W. tomentosa* Roem. & Schult. var. *cochin-chinensis* Pierre ex Pit.

Vernacular names — Mouk (L); Múc lông mềm, Thường mực lông (V).

Shrub or trees 10–20 m high, 50–60 cm diameter. *Bark* 3 mm thick, grey-white to grey-yellow, slightly rough, peeling off in thin pieces. *Inner bark* yellowish with much milky sap and soft and durable fibres. *Branchlets* cylindrical, tomentose; becoming glabrous and lenticellate. *Petiole* 3–8 mm long. *Leaves* opposite, usually with glands in the axils; blade papery to coriaceous, elliptic to obovate, 2.5–11.6 by 1.4–5.2 cm, base cuneate to obtuse, apex acuminate, secondary veins 8–15 pairs, curved from midrib to the margin, densely tomentose or slightly pubescent. *Inflorescences* a terminal cyme, few- to many-flowered, 3–8 cm long; peduncle 1–2 cm long; tomentose or slightly pubescent. *Bracts* ovate, 2.5–3 by 0.8–1 mm. *Flowers* white, sometimes pinkish; pedicel 4.3–10 mm long. *Bracteoles* broadly ovate, 2 by 1.8 mm. *Sepals* 5, ovate, 1–3.5 by 1.2–1.7 mm, apex obtuse to acute, tomentose or slightly pubescent. *Petals* lobes overlapping to the left in bud, mature rotate, narrowly elliptic, 8.5–27.2 mm long, apex obtuse or rounded, tube 3–7 mm long, pubescent-papillose on lobes inside and outside, scales 10, linear, as long as stamens, the ones opposite sepals slightly shorter and forked. *Stamens* inserted at corolla mouth, filaments 0.7–1.8 mm long, anthers 6.2–7 by 1.3–1.4 mm, pubescent on both surfaces. *Ovary* superior, 1.6–2.1 mm long, glabrous, carpels medially connate, style (incl. pistil head) 8.5–10.2 mm long. *Fruit* connate follicles, cylindrical, 9.3–28 by 0.9–1.7 cm, splitting into two at dehiscence, minutely slightly pubescent or rarely glabrous. *Seeds* linear, 10.7–15.7 by 1.1–2.6 mm, with a tuft of hairs at tip, attached to the testa of the seed, 1.2–4.1 cm long.

Distribution — South China, Thailand, Cambodia, Malesia. In Laos: Champassak (Lakhone) provinces. In Vietnam: Vinh Phu, Nghê An, Khanh Hoa, Thanh Hoa, Gia Lai, Kon Tum, and Dac Lac provinces.

Habitat & Ecology — Light demanding and drought-tolerant tree, in disturbed open area, on dry and poor degraded but prefers deep soil, altitude 280–800 m. Flowering: April to August; mature fruiting: August to February.

Uses — Wood is used for general construction, pencils, musical instruments, house appliances, windows, doors, and carving. The latex has been used against severe dysentery.

Specimens examined: *Balansa* 600, 2119, 2120, 2121, 2122 (P); *Bon* 1193, 1194, 1543, 1561, 1566, 1568, 1814, 1815, 1816 (P); *Chevalier* 29624, 31403, 32064, 32117, 39495, 40954, 41000 (P), 31403 (L); *Cuong* 1716 (P); *Eberhardt* 4345 (P); *Evrard* 1666 (P); *Fleury* 30172, 32080, 32117, 37620, 37728 (P); *Harmand* 694, 1149 (P); *Hayata* 245, 425 (P); *Hiep* 446, 514, 993 (P); *Mouret* 290 (P); *Muller* 1051 (P); *Pételot* 955, 958, 1411, 2446, 4917, 5988, 5991, 5994, 6003 (P); *Phung Van Dieu* 106 (P); *Poilane* 4992, 5315, 5584, 8773, 13006, 13323, 18544, 19792, 19932 (P); *Spire* 120, 570 (P); *Thorel* 696 (P).

BETULACEAE

Deciduous trees or shrubs. *Pith* triangular in cross section. *Stipules* present, usually early caducous. *Leaves* simple, alternate, petiolate. *Margin* usually doubly serrate, rarely simply serrate, lobed or entire, secondary veins pinnate, mostly with domatia in the axils of secondary veins with the midrib. *Flowers* unisexual. *Male inflorescence* precocious, elongate, pendulous. *Bracts* numerous, overlapping; each bract usually subtending a small cymose inflorescence with 1–3 flowers. *Sepals* scale-like. *Petals* absent. *Stamens* many, opposite sepals, filaments very short, connate or nearly so; anthers 2-loculed, opening by longitudinal slits. *Female inflorescence* pendulous or erect, with numerous overlapping bracts; each bract usually subtending a small cymose inflorescences with 2 or 3 flowers. *Sepals* 1–6 scale-like, or much reduced. *Petals* absent. *Ovary* inferior, 2-loculed, styles 2, free, ovule(s) 1 or 2, pendulous from near apex of each locule. *Fruit* a nut or nutlet, winged or not. *Seed* 1, with straight embryo and flat or thickened cotyledons, endosperm absent.

Distribution — Mainly in Asia, Europe, North and South America: 6 genera, c. 200 species. In Laos: 2 genera, 2 species. In Vietnam: c. 2 genera, 2 species.

BETULA L.

Betula alnoides Buch.-Ham.

Synonyms — *Betula acuminata* Wall.; *Betulaster acuminata* Spach

Vernacular names — Sakai (Xieng Khouang), Ketsana (Champassak, Bolaven) (L); Cáng lò (V).

Deciduous trees up to 30 m high, up to 60 cm diameter. *Trunk* straight and cylindrical, with rather open crown and slightly drooping branches. *Young bark* shiny brown, pelling in very thin horizontal flakes, medium-sizes trees with deeply fissured bark, rough and irregular flaking vertically when old, greyish or red-brown and easily rotting. *Inner bark* pale brown, contains oil, strongly aromatic. *Petioles* 0.6–1.3 cm long, slender. *Leaves* ovate 4–14 by 2.5–5 cm, base cuneate or nearly rounded, margin sharply and irregularly toothed, apex acuminate. Young leaves densely silvery hairy, mature leaves thin, smooth or with tufts of hairs (domatia) in the axils of secondary veins, lower surface with many tiny resinous dots, secondary veins 9–13 pairs. *Flowers* tiny, greenish, in drooping catkins along twigs or behind young leaves, male and female in different catkins but on same tree. *Male flowers* in slender catkins, up to 18 cm long, individual flowers in groups of 3 at the axil of a hairy bract. *Sepals* 4, linear, hairy. *Stamens* 4–6, with short filament, anthers 2. *Female flowers* in catkins, up to 9 cm long, thicker and denser, in groups of 2–5. Individual flowers solitary in the axil of a 3-lobed bract. *Fruit* a small nut, 1.5–4 mm long, with 2 thin wings 2 or 3 times broader than nut, clusters covered with spiky bracts that persists on a tree long after fruits have blown away by the wind.

Distribution — Subtropical forests of Nepal, India, Myanmar, South China, Thailand. In Laos: all northern provinces and in the southern provinces Saravane, Savannakhet (Se Pon), Champassak (Bolaven). In Vietnam: Tuyên Quang, Hà Giang, Lào Cai, Lang Son, Cao Bang, and Kon Tum provinces.

Habitat & Ecology — Light demanding and fast-growing tree. Can be found in pure stands in secondary forests, in moist evergreen forests on rich and deep loamy soil, in

dry evergreen hill forests on upper hill slopes and ridges on yellow-red soils. Mixed with *Calophyllum*, *Canarium*, *Castanopsis*, *Machilus*, *Pinus*, *Quercus*, and *Tristaniopsis* species. Altitude 300–2100 m. Natural regeneration on bare soil, even in erosion gullies, along roads and fallow forests. Flowering: usually October to February, rarely in July; fruiting: March to June.

Uses — Wood very hard and durable, used for house construction, household utensil-making, fuel. Young bark used as additive for drinking water. The aromatic oil can be used in the leather industry.

Specimens examined: *Averyanov et al.* VH 142, VH 955 (P); *Pocs & Tiep* 505 (P); *Poilane* 2072, 12163, 15423, 16162 (P);

BIGNONIACEAE

Trees, shrubs, or lianas. *Twigs* often lenticellate and nodes with gland fields, spines very rare. *Stipules* absent, sometimes pseudostipules present (*Markhamia*). *Leaves* generally opposite, compound, 1–3 times pinnate, or in some cases simple and whorled or simple and spirally arranged, often with glands on the leaflets, margin entire to serrate, venation pinnate to palmate, domatia sometimes present. *Inflorescences* generally terminal, bracts present, cymose rarely racemose or even reduced to solitary flowers. *Flowers* bisexual, bilateral, usually large and showy, trumpet or funnel-shaped. *Calyx* tubular with or without 5 lobes or teeth, often closed in the bud and filled with water. *Corolla* zygomorphic, united basally to form a dorsiventrally flattened tube, with 5 lobes, 2 on the upper side, 3 on the lower. *Stamens* 4 or 5, mostly 4 and didynamous (2 long and 2 short) attached to the corolla. *Ovary* superior, 2-celled, stipe long, with 2-lobed stigma. *Fruit* a capsule, pod-like, but divided by a longitudinal partition into 2 compartments, splitting open when ripe. *Seeds* many, flat, winged or fringed with hairs, endosperm lacking.

Distribution — Widely distributed in tropical and subtropical regions: c. 120 genera, 650 species. In Laos: c. 9 genera, c. 15 species. In Vietnam: c. 8 genera, c. 22 species.

KEY TO THE GENERA

- 1a. Leaves imparipinnate, 25–55 cm long, the lowest pair of leaflets at the base of the petiole much smaller than the others and stipule-like (pseudostipules). Stamens 4, didynamous. Fruits 40–55 cm long, covered with fulvous hairs. Seeds with wings on both ends. **1. Markhamia**
- b. Leaves 2 or 3 times pinnate, up to 150 cm long, pseudostipules absent. Stamens 5, equal or nearly equal. Fruits 40–120 cm long, glabrous. Seeds with a wing all around **2. Oroxyllum**

1. MARKHAMIA Seem.

Markhamia stipulata (Wall.) Seem. ex K. Schum.

Synonyms — *Bignonia stipulata* (Wall.) Roxb.; *Markhamia indica* P.H. Hô; *Spathodea stipulata* Wall.; *S. velutina* Kurz

Vernacular names — Khai (L); Đĩnh, Thiết đĩnh (V).

Medium or large trees, 15–25 m high, 60–80 cm diameter. *Twigs* densely covered with caducous fulvous hairs. *Bark* grey-yellow, thin, slightly fissured, inner bark consisting of alternate pale orange and dark orange layers. *Petioles* 4–12 cm long. *Leaves* imparipinnate, 25–55 cm long, petiolules very short 0–3 mm long; leaflets elliptic, 5–9 pairs, 8–20 by 3–7 cm, base rounded or obtuse, margin entire or denticulate, apex acuminate or attenuate, lower surface of mature leaflets with pale brown hairs, the lowest pair of sessile leaflets at the base of petiole much smaller than others and stipule-like. *Inflorescence* a raceme. *Flowers* 7–12 cm long, cream or brownish yellow. *Calyx* 3–5.5 cm long, deeply split on one side only, curved sharply backwards at the top, without lobes or teeth. *Corolla* broadly funnel-shaped large, wart-like glands outside. *Stamens* 4, didynamous, filaments brownish violet. *Fruit* flat 40–70 cm long, 3–4.5 cm wide, covered with fulvous hairs. *Seeds* 3–5 cm long, rectangular with thin semi-transparent wings on both ends.

Distribution — In Laos: Louang Prabang, Vientiane, and Savannakhet provinces. In Vietnam: from North to South Vietnam, Kien Gian, Hoa Binh, Hà Tây, Thanh Hoa, Nghê An, Quang Binh, Quang Tri, Phu Yên, and Khanh Hoa provinces, but concentrated in Tuyên Quang and Lang Son provinces.

Habitat & Ecology — Growing in evergreen forest on limestone mountains, mixed with *Aglaiia gigantea*, *Cinnamomum obtusiflorum*, and *Dracontomelon dao*. Flowering: October to January next year; fruiting: February to May.

Uses — The wood is hard and heavy and resistant to insects, especially termites; used for valuable furniture and construction. Young shoots are sold on local markets in Laos as vegetable.

Note — Santisuk & Vidal, *Flore du Cambodge, du Laos et du Vietnam* 22 (1985) 52 considered *Markhamia cauda-felina* (Hance) Sprague a synonym of *Markhamia stipulata* var. *kerrii* Sprague. We think that *Markhamia cauda-felina* is a good species, differing from *M. stipulata* in being a small tree, never reaching more than 15 m high, with much longer petiolules and with much longer fruits.

Specimens examined: *Bejoud* 95 (L); *Bon* 4702, 5218, 5407 (L); *Fleury* 30911 (L); *Hiep* 311 (L); *Kostermans* 355 (L); *Larsen* 8791 (L); *Linh Gioc* 13 (L); *Muller* 199 (L); *Pételot* 330 (L); *Poilane* 25003, 28050 (L); *Soejarto* 5870 (L).

2. OROXYLUM Vent.

Oroxylum indicum (L.) Kurz — Fig. 9

Synonyms — *Bignonia indica* L.; *B. pentandra* Lour.; *Calosanthos indica* (L.) Blume; *Spathodea indica* (L.) Pers.

Vernacular names — Lin maiz (L), Ing ka (Xieng Khouang, L); Núc nác (V), Tonok (Gia Lai, Kom Tum, V).

Evergreen or partly deciduous trees up to 18 m high. Young trees with a single main stem with the leaves clustered at the top like in most palms. *Bark* pale creamy brown or pale grey, twigs very stout, pale grey, with large scars. *Leaves* up to 150 cm long, 2 or 3 times pinnate with terminal leaflets, swollen at the nodes of main and lateral rachis. Petiolules 5–20 mm long. *Leaflets* 5–20 mm long, broadly elliptic, 6–15 by 4–9 cm, base rounded or cuneate, margin entire, apex acuminate or pointed. *Inflorescences* in




Fig. 9. *Oroxylum indicum* (L.) Kurz (Bignoniaceae).

terminal racemes, 60–18 cm long. *Flowers* large with a foxy stink. *Calyx* tubular, 2.5–4 cm long, without lobes, filled with water in bud. *Corolla* funnel-shaped, with 5 lobes, c. 13.5 cm long, c. 12 cm wide, fleshy, lurid reddish purple to liver brown outside, greenish white and pinkish yellow inside, lobes wrinkled. *Stamens* 5, equal or nearly equal, hairy at base. *Fruits* 40–120 cm long, dark brown, flattened, slightly curved at base, with a fine ridge on each side, woody, splitting into 2 sections lengthwise. *Seeds* many, 4–8 by 3–5 cm, with a thin, semi-transparent wing all around.

Distribution — India, China, Indochina, Malaysia. Very common in Laos and Vietnam.

Habitat & Ecology — A fast growing species, found along the forest fringes, roadsides, and planted around the houses in midland regions. Prefers deep, wet, and well-drained soil. Usually mixed with *Alangium kurzii* and *Girroniera subaequalis*. Flowering: April to August; fruiting: October to December.

Uses — Wood soft, can be used for making matches, firewood, and for paper as the fibres are long. Young fruits edible. Seeds used to cure stomachache, lasting cough, pleuro-pneumonia and pimples. The bark is rich in tannin.

Specimens examined: *Balansa* 800 (P); *Bon* 1728 (P); *Chevalier* 8595, 19587, 38197 (L); *Domat* 38197 (P); *Dournes s.n.* (P); *Dupuy* 201 (P); *Dussaud* 77 (P); *Fleury* 30134 (P); *Geesink et al.* 6077 (L); *Hahn* 1297 (P); *Kostermans* 94, 1456 (L); *Larsen et al.* 2304 (L); *Massie s.n.* (P); *Palee* 44 (L); *Pierre* 232 (P); *Poilane* 661, 1808, 10015 (P), 8595, 19587 (L, P); *Pottier* 223 (P); *Soejarto et al.* 5886, 10544, 11528 (L); *Thorel* 1130 (P); *Vidal* 916 (P); *Welzen* 33 (L).

BOMBACACEAE

Trees, trunk straight, usually with buttresses at base and large spreading crown. *Bark* with sharp conical thorns. *Stipules* present, early caducous. *Leaves* alternate, simple or digitately compound, usually with scales. *Inflorescences* terminal or axillary racemes, cymose, fasciculate or solitary. *Flowers* large and showy, bisexual. *Sepals* cup-shaped, apex irregularly 3–5-lobed, lobes not overlapping. *Petals* large, free, overlapping. *Stamens* long, many, filaments are united in bundles or in a tube (monodelphous) or arranged into five groups or clusters (pentadelphous), anthers reniform. *Ovary* superior, 2–5 locules, ovules 2 to many in each locule. *Fruits* a big woody capsule, smooth, usually splitting into 5 parts filled with fluffy silky hairs. *Seeds* ovoid, or round, glabrous or enclosed by wool.

Distribution — Worldwide, mainly in tropical America: c. 20 genera, 250 species. Laos: 1 genus, 1 species. Vietnam: c. 2 genera, 6 species.

BOMBAX L.

Bombax ceiba L.

Synonyms — *Bombax malabaricum* DC.; *B. thorelii* Gagnep.

Vernacular names — Ngiu pa, Ngiu deng (L); Gạo, Mộc miên (V).

Large, deciduous trees up to 35 m high, up to 340 cm diameter. *Trunk* straight, buttresses prominent and large at the base when old. *Bark* pale grey or cream, studded with sharp conical thorns when young, smooth when old. In young trees the lateral branches are horizontal and straight. *Crown* large and lax. *Stipules* minute. *Petiole* 10–20 cm long. *Leaves* digitately compound, alternate, petiole 10–19 cm long. *Petiolule* 1.5–2.5 cm long. *Leaflets* 5–7, ovate, or narrowly elliptic, 8–15 by 4–5 cm, base broad or tapering, margin entire, apex acuminate, glabrous, secondary veins 9–15 pairs. *Flowers* bright orange-red, solitary or fasciculate, 1–3 flowers on a twig, 8–12 cm long, appearing after the old leaves fall. *Pedicel* 1–2 cm long, thick, glabrous or puberulous. *Calyx* 5, bright green, 1.5–3 cm long, glabrous outside, tomentose inside. *Petals* 5, red or

orange red, thick and fleshy, spreading, obovate, 8–10 by 3–4 cm, tomentose on both sides. *Stamens* pale orange, many, in two whorls, fused together into 10 bundles, anthers, attached in the middle of the lower surface (peltate). *Ovary* conical, style slender, dark red, spreading, longer than stamens, stigma with 5 short branches. *Fruit* a woody capsule, elliptic, 8–17 by 4–6 cm, opening longitudinally into 5 segments, inner walls covered with silky cotton-like hairs. *Seeds* numerous, ovoid, glossy, with silky cotton-like hairs.

Distribution — India, South China, Thailand, Cambodia, Malaysia, Indonesia. In Laos: along the Mekong. In Vietnam: Tây Ninh province.

Habitat & Ecology — In natural forests, altitude up to 800 m, or planted in all parts of the countries. Suitable for growing on deep, dry, sandy soil. The species grows very fast. Can be propagated by seedlings or by stumps. Resistant to slight frost, not affected by drought and not killed by fires. Flowering: February to April; fruiting: May to July.

Uses — Wood is used for timber, plywood, common implement-making as boxes, matchsticks, sawing boards, wooden shoes, coffins, and firewood. The floss is used as stuffing in cushions, mats, and pillows. Inner bark yields good fibres, suitable for cordage. The tree is planted as an ornamental.

Note — We have seen only two specimens from Laos and Vietnam available at the Paris Herbarium. The description is mainly based on specimens from Thailand and Sri Lanka. — Gagnepain, Fl. Gen. Indochine (1910) 397 recognizes two species, i.e. *B. malabaricum* and *B. thorelii*. After studying the type material of both we decided to unite them with *B. ceiba* as there are no major differences.

Specimens examined: Maxwell 88-108 (L); Muller 732 (P); Robyns 7189, 7190, 7316, 7326, 7332 (P); Thorel 3003, s.n. (P).

BURSERACEAE

Medium to large buttressed trees, rarely shrubs, crushed living parts with strong resinous smell. *Bark* pale grey to brown, smooth or scaly, often lenticellate, surface sometimes with brown, black, or white resin patches; inner bark usually pinkish, laminated or mottled with droplets of clear or white resinous gum, with strong resinous smell in most species (especially those of *Canarium* and *Triomma*). Sapwood often white, shiny, sometimes pinkish or pale yellowish brown, vessels usually visible with the naked eyes. (Pseudo)stipules present in all species of *Garuga* and in most species of *Canarium*. *Leaves* compound, usually imparipinnate, rarely simple. Leaflets margin entire, except in some *Canarium* and *Garuga* species. Leaves and leaflets distinctly stalked, the stalks kned at both ends, the leaf-stalk swollen at the nodes even when dry. *Inflorescences* panicles, sometimes racemes, thyrses or spikes, axillary or terminal. *Flowers* bisexual, unisexual, or polygamous, dioecious, 3- or 5-merous, usually greenish to cream or white, the remains of the other sex persisting, sepals valvate, mostly united, petals valvate, free, stamens usually twice as many as petals, filaments free or united, sometimes fused to the disc, anthers elongate, dehiscing inwards. *Disc* present, inside stamens (except in *Triomma*). *Ovary* superior, with 3–5 cells, each with 2 descending ovules, style simple, stigma globular often slightly lobed. *Fruit* a drupe with a fleshy to leathery resinous wall and a more or less woody stone, or in *Triomma* a woody capsule.

Stone 3-celled or less by reduction. *Seed* one, with fleshy cotyledons, containing oil, endosperm absent. *Cotyledons* emerging from the ground on germination.

Distribution — Throughout tropics and subtropics: 16 genera, 550 species. In Laos: so far not yet recorded. In Vietnam: c. 4 genera, c.13 species.

CANARIUM L.

KEY TO THE SPECIES

- 1a. Slash of the bark with white sticky sap. Stipules present. Inflorescences shorter than the leaves, flowers polygamous. Fruits yellowish green when ripe, stone with 6 ribs on the surface, cuneate at both ends **1. *C. album***
- b. Slash of the bark with black sap. Stipules absent. Inflorescences longer than the leaves, flowers unisexual. Fruits dark violet when ripe, stone with smooth surface, rounded at both ends **2. *C. pimela***

1. *Canarium album* (Lour.) Raeusch.

Synonym — *Canarium tonkinensis* Engl.

Vernacular name — Trám trắng (V).

Large trees up to 30 m high, 80 cm diameter. *Trunk* cylindrical. *Bark* light brown, fissured with many lenticels. Outer bark thin, inner bark pinkish brown, laminated, aromatic with white sticky sap. *Stipules* awl-shaped, caducous. *Petioles* 5–8 cm long, swollen at base. *Leaves* imparipinnately compound. *Leaflets* opposite, 4–8 pairs, elliptic, 6–15 by 2.5–6 cm, crustaceous, with many shining white scales beneath, base slightly oblique, margin entire, apex distinctly acuminate, petiolules 5–10 mm long, basal pair of leaflets usually smaller. *Inflorescences* panicles, usually shorter than the leaves. *Flowers* polygamous. *Pedicels* 2–7 mm long. *Calyx* cupular, lobes deltoid, outside hairy or glabrous, inside densely silky-hairy. *Petals* 3, cream-white, free, usually overlapping in bud, outside hairy in centre, inside usually glabrous. *Stamens* 6, in one whorl, free, united at base, filaments flattened. *Disc* within the stamens, 6-lobed, strongly developed in bisexual flowers. *Ovary* ovoid, 3-celled. *Fruit* an oblong drupe, 3.5–4 by 2–2.5 cm, round at both ends, yellowish green when ripe; calyx persistent in fruit. *Stone* hard, woody, with 6 ribs on the surface, cuneate at both ends, 3-celled. *Seed* one per cell.

Distribution — China, Japan, Vietnam, Malaysia. In Vietnam: most provinces, particularly in Tuyên Quang, Quang Ninh, Bac Thai, Kien Gian, Vinh Phu, and Thanh Hoa.

Habitat & Ecology — Distributed in primary and secondary forest, altitude below 500 m. Usually mixed with *Erythrophloeum fordii*, *Gironniera subaequalis*, *Hopea odorata*, *Peltophorum tonkinensis*. This species is fast-growing and light demanding. Flowering: March to May; fruiting: June to October.

Uses — Wood used for sawing board, house construction, and fuel. The resin is used for incense or as perfume oil, turpentine, and raw material for paint and printing industries. Fruit edible and used against diarrhoea and rheumatism. Seeds contain edible oil.

Specimens examined: *Balansa* 3706 (P); *Brillet* 5 (P); *Cuong* 391 (L); *Fleury* 30122, 37542, 37960 (L, P), 30141, 37921, 38026 (L); *Pételot* 5751 (P); *Poilane* 29914 (P); *Thirlot* s.n. (P); *Tsang* 39334 (L).

2. *Canarium pimela* Leenh.

Synonym — *Canarium tramdenum* C.D. Dai & Yakovlev

Vernacular names — Bai (L); Trám đen (V).

Large trees up to 30 m high, 90 cm diameter. *Trunk* cylindrical. *Bark* light brown, fissured with many lenticels, inner bark pinkish brown, aromatic with black sap. *Stipules* absent. *Petioles* 7–9 cm long, swollen at base. *Leaves* imparipinnately compound, 30–65 cm long. *Leaflets* opposite, 2–4 pairs, elliptic, 7–18 by 3–6.6 cm, stiff, crustaceous, smooth above, dark coloured beneath, apex distinctly acuminate or pointed, margin entire, oblique at base, petiolules 3–6 mm long. *Inflorescences* panicles, longer than the leaves. *Flowers* unisexual, dioecious, pedicel 6–8.5 cm long. *Calyx* cupular, lobes deltoid, outside hairy or glabrous, inside densely silky-hairy. *Petals* 3, cream-white, free, usually overlapping in bud, outside hairy in centre, inside usually glabrous. *Stamens* 6, in one whorl, united at base. *Filaments* flattened. *Disc* within stamens, 6-lobed, strongly developed in male flower. *Ovary* ovoid, 3-celled. *Fruit* a long-ovoid drupe, 3–4 by 1.5–2 cm, 3 unequal loculi, dark violet when ripe; calyx persistent in fruit. *Stone* hard, woody, surface smooth, rounded at both ends, 3-celled. *Seed* one per cell.

Distribution — South East China, Indochina. In most provinces of Laos, mainly in Xieng Khouang, Houa Phan, and Vientiane. In almost all provinces in northern Vietnam, mainly in Bac Thai, Tuyên Quang, Kiên Giang, Vinh Phu, Hà Tây, Hoa Binh, Ninh Binh, Thanh Hoa, Nghệ An, and Quang Binh.

Habitat & Ecology — In primary and secondary forests, altitude below 500 m. Usually mixed with *Aglaiia gigantea*, *Eberhardtia tonkinensis*, *Erythrophloeum for-dii*, *Garcinia* spp., *Gironniera subaequalis*, and *Pygeum arboreum*. This species is fast-growing and light demanding, preferring deep humid soil and moderately steep topography. Natural regeneration is good under a forest cover of 30–40%. Flowering: April, May; fruiting: May to September.

Uses — The wood is used for sawing board, house building, and furniture. Fruits are edible. Fruits and leaves are used against diarrhoea and rheumatism. Seeds contain edible oil.

Specimens examined: *Balansa* 3694, 4602 (P); *Bon* 1545 (P); *Cuong* 103 (L); *Fleury* 30091, 30141, 37833 (P); *Forestier* 7, 8 (P); *Kerr* 20890 (P); *Poilane* 2013 (P).

CAESALPINIACEAE

Trees, shrubs, sometimes lianas, rarely herbs. *Buttresses* sometimes present. *Stipules* nearly always present, stipes usually lacking, minute when present. *Leaves* usually spirally arranged, rarely opposite or alternate, pinnate, bipinnate, or unifoliolate, rarely simple; sometimes with translucent dots; leaflets entire. *Inflorescences* axillary and/or terminal, usually in racemes, panicles or spikes, rarely solitary flowers. *Flowers* regular or irregular, large to small, often showy, bisexual. *Sepals* adnate, normally

5-lobed, overlapping or separate. *Petals* usually 5, sometimes imperfectly developed or absent, sometimes only one (*Afzelia xylocarpa*, *Sindora*), slightly unequal, the upper petal distinctive and innermost in bud; stamens free or joined, 10 or fewer, sometimes staminodes present; anthers dehiscent by lateral slits or terminal pores, ovary superior. *Fruit* a pod, sometimes indehiscent, sometimes winged, or fleshy and drupe-like.

Distribution — In tropical and subtropical regions of the world: c. 180 genera, 2500–3000 species. In Vietnam: 20 genera, 97 species. In Laos: 18 genera, 71 species.

KEY TO THE GENERA

- 1a. Leaves bipinnate 2
- b. Leaves pinnate 5
- 2a. Stipules lobed or branched. 3
- b. Stipules entire 4
- 3a. Petals with long claw. Pods dehiscent, glabrous **5. Delonix**
- b. Petals not clawed. Pods indehiscent, glabrous or hairy **7. Peltophorum**
- 4a. Plants armed with spines. Bracteoles absent. Fertile stamens 10. Pods dehiscent or indehiscent **3. Caesalpinia**
- b. Plants not armed. Bracteoles present. Fertile stamens 5. Pods dehiscent **1. Acrocarpus**
- 5a. Exudate present, red. 6
- b. Exudate absent 7
- 6a. Leaves paripinnate. Sepals 4. Petal 1. Fertile stamens 7. Pods dehiscent **2. Afzelia**
- b. Leaves imparipinnate. Sepals 5 (rarely 3). Petals absent. Fertile stamens 2 (rarely 6). Pods indehiscent **6. Dialium**
- 7a. Sepals 5 8
- b. Sepals 4 9
- 8a. Bracteoles present. Filaments straight or curved. Anthers dehiscent by apical pores. **4. Cassia**
- b. Bracteoles absent. Filaments straight. Anthers dehiscent by slits or basal pores **8. Senna**
- 9a. Leaflets 2–5 (rarely more) pairs, 30–150 by 20–80 mm. Sepals spiny outside. Petal 1. Fertile stamens 10. Pods dehiscent, flattened. **9. Sindora**
- b. Leaflets 10–20 pairs, 8–30 by 3–10 mm. Sepals not spiny outside. Petals 5. Fertile stamens 3. Pods indehiscent, thickened **10. Tamarindus**

1. ACROCARPUS Wight & Arn.

Acrocarpus fraxinifolius Wight & Arn.

Synonym — *Acrocarpus combretiflorus* Teijsm. & Binn.

Vernacular names — Ket hoy (Sayabouri), Khan khak (Louang Prabang) (L).

Trees up to 50 m high, deciduous at the beginning of the cold season, often with buttresses when older. *Bark* pale grey with large brown lenticels, inner bark pinkish. Heartwood dark red. Young shoots appressed yellowish tomentose. *Stipules* triangular, small, 1–2 mm long, usually early caducous. *Leaves* up to 100 cm long, bipinnate with

3–5 pairs of pinnae. *Leaflets* 4–9 pairs per pinna, ovate, 4–14 by 2–7 cm, slightly oblique at the base, apex acuminate, subsessile; young leaves pink, pubescent beneath; mature leaves pale green, glabrous. *Inflorescences* racemes 1–3 together, dense, 20–25 cm long. *Pedicels* 6–8 mm long. *Flowers* reflexed after anthesis. *Receptacle* pubescent. *Sepals* 5, 3–4 mm long, apex rounded, bright green, pubescent. *Petals* 5, red, obovate, 5–10 by 1–2.5 mm, apex acute, pubescent on both surface. *Stamens* 5, yellowish orange, green at base, glabrous, twice as long as petals. *Style* short, curved, pale green, stigma small, ovules 15–20. *Pods* long-stalked, 8–15 by 1–2 cm, flat, slightly torulose, with a 3–5 mm broad wing along the upper suture, dehiscent. *Seeds* 12–18, 6.5 by 5 mm, lens-shaped, pale brown.

Distribution — India (Sikkim, Assam), Myanmar, Thailand, Indonesia (Sumatra, Java). In Laos: Louang Prabang (Phou Phung) and Sayabouri (Pak Lep) provinces. In Vietnam not yet recorded.

Habitat & Ecology — In evergreen gallery-forests, altitude 400–1100 m. Flowering: January, February after shedding the leaves (pers. obs.).

Uses — Timber used for indoor constructions and for plywood.

Note — We have seen only one specimen from Laos at the Paris Herbarium. The description is mainly based on specimens from Thailand.

Specimens examined: *Dussaud* 84 (P); *Poilane* 20290 (L); *Maxwell* 90-369, 96-545, 97-143 (L); *Teijsman s.n.* (L); *Wight* 845 (P).

2. AFZELIA Sm.

Afzelia xylocarpa (Kurz) Craib — Fig. 10

Synonyms — *Afzelia cochinchinensis* (Pierre) Leonard; *A. siamica* Craib; *Pahudia cochinchinensis* Pierre; *P. xylocarpa* Kurz

Vernacular names — Tae kha, Kha, Go ca te (L); Gỏ đỏ (V).

Deciduous trees up to 30 m high, up to 100 cm diam. or sometimes even more. *Crown* broad, rounded. *Trunk* stout, usually dividing near base into large, spreading branches. *Bark* smooth or slightly rough, grey or yellowish, inner bark reddish, forming burls when cut. *Stipules* minute, usually early caducous. *Leaves* 18–25 cm long, paripinnate with 3–5 pairs opposite leaflets. *Leaflets* elliptic, 5–9 by 4–5 cm, base rounded or truncate, apex obtuse to acute, slightly emarginated, young leaflets pubescent, mature leaves completely glabrous, sometimes pubescent below. *Inflorescences* panicles 5–15 cm long, greyish pubescent. *Pedicels* 7–10 mm long. *Bracts* ovate to narrowly elliptic, 6–9 mm long, greyish pubescent; bracteoles 2, as the bracts, persistent. *Sepals* 4, narrowly elliptic, 10–13 mm long, finely velvety outside. *Petal* 1, broadly elliptic, 7–9 mm long, pink or reddish, claw thin, 5–12 mm long, apex emarginate, hairy at the base. *Fertile stamens* 7 or 8, filaments 30 mm long, pubescent at base; anthers 3–4 mm long, opening by a slit. *Sterile stamens* 3, shorter than the fertile ones, filiform. *Ovary* hairy along both sutures, c. 7 mm long, stipe 7 mm long, style glabrous, 20–25 mm long, slightly enlarged towards the stigma. *Pods* brown, woody, 15–20 by 7–9 cm, dehiscent, valves 5–7 mm thick. *Seeds* 2.5–3 by 2–2.5 by 0.8–1.2 cm, aril orange, cup-shaped, c. 1.5 cm long.




Fig. 10. *Afzelia xylocarpa* (Kurz) Craib (Caesalpinaceae).

Distribution — Myanmar, Indochina, Thailand. In Laos: Vientiane (Ban Na Kham) and Savannakhet (Muong Tin) provinces. In Vietnam: Gia Lai, Bac Thai, Đông Nai, Ninh Thuận, Tây Ninh, and Hồ Chí Minh provinces.

Habitat & Ecology — In mixed deciduous and dry evergreen forests at lowland altitudes, in transitional zones between evergreen and dry open Dipterocarp forest. Flowering: March, April; mature fruits: October to December.

Uses — Timber used for construction, cabinet work, furniture, and in wood carving industries. Young seeds edible.

Specimens examined: *Chevalier* 29937, 32031 (L), 4, 30074, 31315, 32031 (P); *Dong* 168 (P); *Hiep* 458 (P); *Muller* 907 (P); *Pierre* 497 (P); *Poilane* 5540, 6803, 8477 (L, P), 10064 (L), 7912, 9794, 10046, 12500, 40791 (P); *Tilley* 29937 (P).

3. CAESALPINIA L.

Caesalpinia sappan L. — Fig. 11

Vernacular names — Fang deng, Fang pa (L); Vang, Vang nhuộm (V).

Small trees or shrubs up to 20 m high, 20–30 cm diameter. *Bark* pale grey studded with pale brown lenticels, young twigs and buds finely brown hairy. Spines small, strong, a pair at each leaf-base and several scattered. *Stipules* 3–4 mm long, usually caducous, often in 3 on the twigs, often a few on the underside of the petiole. *Leaves* bipinnate, 15–40 cm long; pinnae 8–16 pairs. *Leaflets* 6–14 pairs per pinna, opposite, subsessile, ovate, 10–20 by 6–10 mm, base asymmetrical, apex slightly emarginate.




Fig. 11. *Caesalpinia sappan* L. (Caesalpinaceae).

Inflorescences racemes, above the axil of leaves, or combined into terminal panicles. *Bracts* narrowly ovate, 6 mm long, acuminate, caducous. *Pedicels* 15–20 mm long, pubescent. *Sepals* 5, overlapping, glabrous, the lowest one more concave and larger. *Petals* 5, the 4 outer ones rounded, large, the fifth petal (standard) smaller. *Stamens* exserted; filaments c. 15 mm long, hairy at base; anthers c. 1.5 mm long, glabrous. *Ovary* greyish pubescent, with 3–6 ovules, style filiform. *Pods* pale green when young, reddish brown when mature, pubescent at both sutures, woody, irregular obovate-elliptic, flattened, 7–12 by 1.5–2 cm, sessile on the receptacle. *Seeds* 2–4, elliptic in outline, 18–20 by 10–12 mm.

Distribution — India, Myanmar, South China, Indochina, Thailand. In Laos: Louang Prabang, Sayabouri (Pak Lay), and Savannakhet provinces. In Vietnam: Hà Nội, Quang Ninh, Thanh Hoa, Đông Nai, Gia Lai, Hồ Chí Minh, and Sông Bé provinces.

Habitat & Ecology — In secondary forest, near roadsides, at the forest-edges, limestone hills, and cultivated around villages, buildings, or in parks. Flowering: June to December; fruiting: December to May.

Uses — Wood is used for wood-carving. Bark is used as yellow dye and wood as red dye. The wood is also used to cure haemostatic and diarrhoea.

Specimens examined: Balansa 1294, 1295, 2148 (P); Bon 2952, 4570, 5471 (P); Chevalier 32516 (L); Cuong 506 (L); Evrard 1744 (P); Fleury 37770 (P); Hiep 134 (P); Lefèvre 100 (P); Poilane 1772, 5184 (L, P), 130, 11515, 12040 (P); Thorel 102 (P); Vidal 2054 (P).

4. CASSIA L.

Cassia fistula L.

Vernacular names — Khoum (General), Lom leng (Sayabouri Prov.) (L); Muồng hoàng yến, Bò cạp nước (V).

Deciduous trees up to 20 m high, up to 50 cm diameter. *Branches* glabrous, spreading and drooping. *Bark* pale brown or dark grey, smooth or slightly cracked; inner bark reddish brown. *Sapwood* white, heartwood yellow. *Stipules* small, usually falling early. *Petioles* 6–10 cm long, rhachis 15–25 cm long. *Leaves* paripinnate, 30–40 cm long. *Leaflets* opposite, 3–8 pairs, broadly ovate-elliptic, 7–18 by 4–8 cm, base cuneate to obtuse, apex acuminate to acute, with silky hairs when young, glabrous when mature, without glands, secondary veins numerous; petiolules 4–10 mm long. *Inflorescences* axillary, clustered racemes, 20–40 cm long, pendent, lax. *Bracts* 8–10 mm long, caducous. *Pedicels* 15–35 mm long, glabrous. *Flowers* bright yellow. *Sepals* 5, ovate-elliptic, 5–10 mm long, velutinous outside. *Petals* 5, broadly ovate, 20–35 by 10–15 mm, subequal, claw short. *Stamens* 10, unequal, 3 long ones with filaments 3–4 cm long, anthers 5 mm long opening by apical and basal slits; 4 shorter with filaments 6–10 mm long, anthers opening by basal pore; reduced stamens 3 with filaments 3–5 mm long and minute anthers. *Ovary* and style velutinous, stigma small. *Pods* pendulous, terete, tube-like, 20–60 cm long, 1.5–2 cm diam., indehiscent, glabrous, black. *Seeds* numerous, separated by spongy septa, elliptic, flattened, 8–9 by 4–6 mm, glossy, brown.

Distribution — Egypt, India, China, Indochina, Thailand, Malesia. In Laos: Louang Prabang (Namkhan, Xieng Ngeun, Ban Na Luang), Houa Phan (Houa Phan), Sayabouri (Pak Lay), Vientiane (Vientiane-Thakhek), Savannakhet (Muong Phalane), Saravane, and Champassak provinces. In Vietnam: Kon Tum, Gia Lai, and Dac Lac, and cultivated in Hà Nội, Nghệ An, Hồ Chí Minh, and Sông Bé.

Habitat & Ecology — In (sub)tropical climate, with an average annual temperature of 25 °C, in mixed deciduous forests, in dry deciduous Dipterocarp forests. Annual rainfall should be over 1227 mm. Tree, inclining towards light demanding, young trees slightly shade demanding and drought tolerant. Flowering: April to July; fruiting: May to August.

Uses — Timber used for construction, implements, and agricultural tool-making. Trunk bark is rich in tannin and used in dyeing. Pods and seeds used for medical purposes (laxative). Bark is traditionally used for treatment of wounds from snake bites and scorpion stings. Trees are planted for ornamental purposes.

Specimens examined: *Chamni 9944* (L); *Maxwell 97-60* (L); *Pedrond 172* (P); *Phengkhlai 37895* (L); *Poilane 2185* (L, P), 11406, 13299, 13346, 13788, 17975, 20153, 20207, 20784, 27975 (P); *Pottier 88, 514, 838* (P); *Saydara 034* (L); *Schmid s.n.* (P); *Thorel 2725* (P); *Vidal 932* (P).

5. DELONIX Raf.

Delonix regia (Bojer ex Hook.) Raf.

Basionym — *Poinciana regia* Bojer ex Hook.

Vernacular names — Fang hang nhoung, In si (L); Phượng, Diệp bông đỏ, Diệp tây (V).

Deciduous trees up to 15 m high. *Crown* umbrella-shaped. *Trunk* more or less buttressed. *Bark* grey, smooth. *Stipules* pinnately 4 or 5 lobed, c. 2 mm long, caducous. *Leaves* bipinnate, main rachis 50–60 cm long; pinnae opposite, 9–24 pairs, up to 10 cm long. *Leaflets* 10–40 pairs per pinna, opposite, subsessile or sessile, narrowly elliptic, 5–15 by 2–5 mm, base slightly oblique, apex rounded, mucronate, both surfaces finely puberulous, glabrescent. *Inflorescences* axillary, racemes, 10–15 cm long, glabrous. *Flowers* 5–10. *Pedicels* 5–8 cm long. Hypanthium shortly bell-shaped. *Sepals* 5, narrowly elliptic, 4–7 by 2–2.5 mm, apex acuminate, reddish inside. *Petals* 5, unequal, total length 3–7 cm long, blade orbicular, 3–4 cm wide, narrowed into an up to 3 cm long claw, one yellowish white and scarlet, the others only scarlet. *Stamens* 10, equal in length; filaments up to 4 cm long, red with white base; anthers elliptic c. 4 mm long. *Ovary* slightly velutinous; style filiform, glabrous, c. 2.5 cm long; stigma indistinct. *Pods* narrowly elliptic, flat, slightly curved, pendulous, 30–80 by 3.5–7 cm, woody-valved, blackish, beaked. *Seeds* 20–40, elliptic, compressed, 25 by 8 mm.

Distribution — Endemic to Madagascar. Widely cultivated in the tropics under the name ‘Flamboyant’ or ‘Flame Tree’. In Laos and Vietnam planted in all tropical parts.

Habitat & Ecology — Cultivated around villages, buildings, or in parks, along roadsides. Flowering: April to July; fruiting: May to August.

Uses — Planted for shade and ornamental purposes, seeds edible.

Specimens examined: *Clemens* 3300 (P); *Dau* 2341 (P); *Dournes s.n.* (P); *Hiep* 507, 872 (P); *Jarois* 79 (P); *Phuc Nhac* 1614 (P); *Poilane* 11729 (P), 11892 (L, P).

6. DIALIUM L.

Dialium cochinchinensis Pierre — Fig. 12

Vernacular names — Kheng (L); Xoay, Lá mét, Xàì mét (V).

Large deciduous trees up to 35 m high, bole 16–18 m high, 80–100 cm diameter, rarely 150 cm. *Trunk* straight, buttresses inconspicuous. *Bark* generally grey or whitish, slightly rugose, inner bark 6–8 mm thick, hard, brownish, exuding a little transparent resin, soon turning red. *Stipules* small, caducous. *Leaves* imparipinnate. *Petioles* 12–20 mm long, glabrous or pubescent; rachis 5–15 cm long, glabrous or glabrescent. *Leaflets* 5–9, alternate, broadly ovate to elliptic, 4–7 by 1.5–4.5 cm, base rounded to acute, slightly oblique, apex obtuse to tapered acuminate, secondary veins 6–10 pairs, anastomosing near the margin; petiolules 4–5 mm long, sparsely pubescent. *Inflorescences* axillary or terminal, panicles, 10–30 cm (or more) long, pubescent. *Pedicels* 2 mm long, pubescent. *Buds* ovoid, 3–4 mm long. *Sepals* 5, elliptic, connected at the base into a short tube, 4 mm long, grey pubescent outside, glabrous inside. *Petals* absent. *Stamens* 2, filaments very short, c. 1 mm long. *Ovary* silky, style glabrous, sessile with 2 ovules, stigma small. *Fruit* a pod, ovoid, slightly laterally compressed, 15–20 by 8–15 mm, finely pubescent to velvety. *Seed(s)* 1 or 2, elliptic, 9 by 6 by 3 mm, longitudinally striate, covered by whitish pulp.

Distribution — Myanmar, Thailand, Indochina, Malaysia. In Laos: Vientiane (Hatxifong, Ban Khuay Daeng, Sikhot Tabong Distr., Ban Ang Nhai) and Savannakhet (Ban




Fig. 12. *Dialium cochinchinensis* Pierre (Caesalpinaceae).

That, Champhon, Ban Phay) provinces. In Vietnam: Nghê An, Hà Tĩnh, Kon Tum, Đông Nai, Sông Bé, Hồ Chí Minh, and Ninh Thuận provinces.

Habitat & Ecology — Primary or secondary forest, mountain slopes, on sandy brown soil, closed or open canopy forest, in dense evergreen or mixed deciduous forests, or transitional forest between evergreen and open Dipterocarp forest, altitude up to 800 m, growing on well-drained basalt, along rivers, and stream banks. Growing together with species from the genera *Altingia*, *Dalbergia*, *Dipterocarpus alatus*, *Pterocarpus*, *Stereospermum*, *Terminalia*, and *Shorea obtusa*. Flowering: March to July; fruiting: June to November.

Uses — The wood is used for furniture, railway sleepers, heavy construction (staircases, doors and window frames), bridge, ship- and boat-building, vehicles, chopping board, and sport equipment, cart axles, agriculture implements, oil and sugar mills. Fruits are edible.

Specimens examined: *Chevalier* 39201 (P); *Harmand* 505 (P); *Phung Van Dieu* 38 (P); *Pierre* 814 (iso), 123, 244, 1026 (P), 1814 (L, P); *Poilane* 123, 141, 17866 (P), 6640 (L), 8460, 16365 (L, P); *Soejarto et al.* 10752, 11480 (L); *Soejarto & Sydara* 10848, 10850 (L); *Thorel* 369, s.n. (P); *Vinot* 16 (P).

7. PELTOPHORUM (Vogel) Benth.

Peltophorum dasyrhachis (Miq.) Kurz — Fig. 13

Synonyms — *Caesalpinia dasyrhachis* Miq.; *Peltophorum grande* Prain

Vernacular names — Sa kham, Sa phang, Sa fang (L); Hoàng linh, Lim xet, Lim vàng (V).

Large deciduous trees up to 30 m high, 70–90 cm diam., open crown. Young shoots or branchlets reddish brown, pubescent, later glabrous. *Bark* greyish brown, lenticels present on twigs, inner bark dark brown to reddish brown. *Stipules* remarkably antler-like, rather large. *Leaves* bipinnate, 15–32 cm long, pinnae 4–9 pairs, young leaves with dense reddish brown hairs. *Leaflets* 6–16 pairs per pinna, irregular elliptic,



Fig. 13. *Peltophorum dasyrhachis* (Miq.) Kurz. (Caesalpiaceae).

10–25 by 4–10 mm, base acute, obtuse or rounded, slightly oblique, apex rounded or slightly emarginated, above glossy dark green when mature, slightly brown hairy below. *Inflorescence* axillary, racemes, 15–30 cm long, with pubescent axis. *Bracts* persistent, linear, 7–12 cm long. *Pedicels* 20–40 mm long. *Buds* ovoid. *Sepals* broad-elliptic, 10–15 by 5–6 mm, brownish or ruddy pubescent outside, glabrous inside. *Petals* yellow, obovate, 15–25 by 10–12 mm, hairy towards the base of the upper side. *Stamens* 10, fertile, with filaments \pm as long as petals, 10–15 mm long; anthers 4–5 mm long. *Ovary* sessile, velutinous, with 4–8 ovules, style slender, stigma knob-like. *Pods* reddish brown, flattened, both ends tapered, 10–15 by 2–3.5 cm, margin thinly winged, 4–5 mm broad. *Seeds* 4–8, more or less transversally arranged in the pod, flat, 10–12 by 5 mm.

Distribution — Indochina, Thailand, Malaysia, Indonesia and furthermore widely cultivated. In Laos: Sayabouri (Pak Lay), Vientiane, Khammouane (Thakhek-Phong Tiou), and Saravane province (Pa Ta Lot). In Vietnam: Lang Son, Nghê An, Thua Thiên-Huê, Binh Dinh, Vinh Phu, Dac Lac, Khanh Hoa, Ninh Thuận, Đông Nai, Sông Bé, Tuyên Quang, Tây Ninh, and Kiên Giang provinces.

Habitat & Ecology — In tropical deciduous and evergreen forest, bordering agriculture fields, altitude up to 800 m. Light demanding when mature and shade tolerant when young. It is a pioneer species on shifting cultivation land or in secondary forests. Often mixed with *Alangium kurzii*, *Engelhardtia chrysolepis*, and *Ormosia balansae*. *Flowering*: February to April when the young leaves develop; *fruiting*: July to October.

Uses — Timber used for poles, house implements, vehicles, railway sleepers, and boat and ship building. Planted for ornamental purposes. Bark is used for medical purposes.

Note — K. Larsen, S.S. Larsen & J.E. Vidal, *Flore du Cambodge, du Laos et du Vietnam* 18 (1980) 60–61 combined *P. dasyrhachis* and *P. tonkinensis* and gave them both only varietal rank. Gagnepain, *Fl. Gen. Indochine* (1916) and Dung, *Vietnamese Forest Trees* (1996) 425–426, however, considered them as two separate species i.e. *P. dasyrhachis* Kurz and *P. tonkinensis* (Pierre) Gagnep. They differ in the stipules: in *P. dasyrhachis* the stipules have many branches, remarkably antler-like; in *P. tonkinensis* the stipules have no or sometimes only two branches.

Specimens examined: Bon 5438 (P); Cuong, Kien & Sinh 979 (L); Dussand 1 (P); Fleury 30142, 30201 (P); Hoi 165 (P); Honquan 799 (P); Lefèvre 312 (P); Maxwell 99-217 (L); Muller 1805 (L, P); Parrean 171 (P); Phung Van Dieu 72 (P); Pierre 219 (P), 1813 (L, P); Poilane 6255, 17312 (L, P), 10678, 13733 (L), 6772, 13746, 30515 (P); Schmid 705, s.n. (P); Service Forester 14, 17, 27, 172 (P); Soejarto & Southavong 10761 (L); Thorel s.n. (P); Vidal 1279, 5854 (P); Vinot 48 (P).

8. SENNA Mill.

Senna siamea (Lam.) Irwin & Barneby

Synonym — *Cassia siamea* Lam.

Vernacular names — Khi lek (L); Muồng đen (V).

Evergreen or semi-evergreen, medium-sized trees up to 20 m high, 30–45 cm diameter. *Bark* nearly smooth. Lenticels present on twigs. Young branches striate, finely

pubescent. *Stipules* minute, subulate, caducous. *Petioles* 2–3 cm long, rhachis 10–25 cm long. *Leaves* paripinnate. *Leaflets* 7–10 (rarely 15) pairs, ovate to elliptic 3–7 by 1–2 cm, base rounded, apex slightly emarginate with a short mucronate tip, upper surface glabrous, lower surface finely pubescent; petiolules short. *Inflorescences* large, terminal panicles. *Peduncle* 5–7 cm long. *Bracts* obovate, 5 mm long, apex long acute. *Bracteoles* absent. *Pedicels* 2–3 cm long, velutinous. *Sepals* 5, orbicular, thick, unequal, 2 outer small, 5 mm long, 3 inner up to 9 mm long, outside pubescent. *Petals* 5, yellow, broadly obovate or ovate, 1–2 cm long, claws short. *Stamens*: 2 with filaments c. 10 mm long and anthers 6–7 mm long, opening by apical pores; 4 or 5 with filaments 2–3 mm and anthers 5–6 mm long, opening the same way; 2 or 3 staminodes c. 3 mm long. *Ovary* densely velutinous, style glabrous, stigma indistinct. *Pods* flattened, glabrous or glabrescent, longitudinally waved with raised sutures, 15–30 by 1–1.8 cm. *Seeds* 20–30, ovate, broadly elliptic, flat, 10–25 by 5–6 mm, pale brown.

Distribution — South East Asia, India, Myanmar, China, Indochina, Thailand, Philippines. In Laos: Louang Namtha, Khammouane (Phong Tiou), and Champassak (Bassac, Sithandone, Khong) provinces. In Vietnam: Naturally distributed in Quang Binh, Kon Tum, Gia Lai, Khanh Hoa, Dac Lac, and Đông Nai provinces, and planted in many cities and provinces.

Habitat & Ecology — Thrives in subtropical climates. Various types of forest at lower altitudes with an annual average temperature between 21–24 °C. Neutral tree, inclining towards light demanding, adapted to calcareous, fertile alluvial soil, grows also on arid soil. Flowering: July to December; fruiting: January to April.

Uses — Timber used for indoor constructions and furniture. Flowers and young leaves edible. Can be planted as a shade tree along streets, or for cover in plantations, as ornamental in parks or roadside trees. The wood burns easily, producing much heat and is suitable for firewood. Locally used for medicinal purposes.

Specimens examined: *D'Alleizette s.n.* (L); *Belval* 61 (P); *Chevalier* 29544, 38659, 39011, 39414 (L), 38342 (P), 40057 (L, P); *Clemens* 3624 (P); *Dournes* 284 (P); *Fleury* 38059, 39010, 39011 (P); *Hiep* 797, 900 (P); *Krong-Pha* 925 (P); *Lamk* 186 (P); *Muong* 40189 (P); *Phung Van Dieu* 6 (P); *Pierre* 5984 (P); *Pételot* 1112 (P); *Poilane* 19, 45, 4690, 8413 (L, P), 9819, 18552 (L), 18551, 26226, 40096 (P); *Robinson* 1446 (P); *Thoauy* 2302 (P); *Thorel* 2720 (P); *Vidal* 1087, 1278 (P).

9. SINDORA Miq.

Sindora siamensis Teijsm. — Fig. 14

Synonyms — *Sindora cochinchinensis* Baill.; *S. wallichii* var. *siamensis* (Teijsm. ex Miq.) Baker

Vernacular names — Te nam (L); Gỏ mậ, Gụ mậ (V).

Deciduous or semi-evergreen trees up to 35 m high, 80–100 cm diam., cylindrical, much branched. *Trunk* 15–20 m high. *Crown* spreading umbrella-like. *Bark* dark brown with grey spots, slightly cracked and flaking when older, inner bark pinkish and rather fibrous. *Twigs* robust, compressed, lenticellate, yellow tomentose, glabrous. *Stipules* early falling, foliaceous, large, ovate, 13–18 mm long, base auricled, apex acute, reticulate veined and puberulous on both surfaces. *Petioles* 2–4 cm long; rhachis up to 10 cm long, both pubescent. *Leaves* paripinnate, 15–25 cm long. *Leaflets* 3 or 4 pairs, upper pair slightly larger than the others, broadly or very broadly elliptic, 6–15 by 4–8




Fig. 14. *Sindora siamensis* Teijsm. (Caesalpiniaceae).

cm, base broadly cuneate, acute, round to truncate, more or less unequal-sided, apex rounded or emarginated, both surfaces reticulate and slightly rough to the touch, upper surface minutely short-hairy, lower surface often more densely hairy especially on the midrib, lateral veins often straight to the margin. *Inflorescences* axillary or terminal, panicles, 10–35 cm long, with straight or zigzag axes, solitary, paired, or tripartite, densely brownish, pubescent. *Bracts* persistent during anthesis, foliaceous, large, ovate or heart-shaped, 3–7 mm long, outside densely puberulous, inside puberulous, pubescent on edge. *Bracteoles* long persistent, foliaceous, ovate, c. 5 by 2 mm, puberulous on both surfaces. *Pedicels* 2–4 cm long, densely dark brown pubescent. *Sepals* 4, imbricate in bud, elliptic to broadly elliptic, 7–10 by 4–6 mm, unequal-sized, outside puberulous, bearing several spinescent outgrowths near the apex and often showing numerous warts all over, inside glossy brown strigose. *Petal* 1, narrowly elliptic, 7–10 by 2–4 mm, when dry dull purple, claw involute, fleshy, with a central thick length zone; outside densely woolly hirsute, inside brown, glabrous. *Stamens* 10, the united basal parts of filaments very short; free filaments 15–20 mm long, slightly hairy at

the lower part; fertile anthers various in size. *Ovary* subsessile, densely hirsute along the suture, spines appear later in development, ovules 3–8, style recurved, glabrous, 8–10 mm long, stigma terminal, knob-shaped. *Pods* armed with numerous stout spines, flattened, irregularly ovate or broadly elliptic, 4–10 by 3–8 cm, beak curved, 5–7 mm long. *Seeds* placed on top of a large aril, orbicular or broadly elliptic, flattened, 15–25 by 14 by 17 mm; dry aril cushion-shaped, 10–15 mm.

Distribution — China, Thailand, Indochina, Peninsular Malaysia. In Laos: Vientiane (Ban Keun, Nam Ngum, Sikhot Tabong Distr., Ban Ang Nhai, Phu Phanang), Savannakhet (Champhon Distr., Ban Phay), Saravane, and Champassak (Khong Island, Khong Distr.) provinces. In Vietnam: Kon Tum, Gia Lai, Dac Lac, Lâm Đông, Đông Nai, Sông Bé, Ninh Thuận, Khanh Hoa, and Tây Ninh provinces. This species has been planted in Hà Nội Botanical Garden and Hồ Chí Minh city zoo.

Habitat & Ecology — Primary forest, remnants of forest with some bamboos, mixed evergreen forest, mixed deciduous forest, dry deciduous Dipterocarp forests.

Uses — Wood is used for construction posts, frame work, furniture, developing a beautiful appearance when used for a long time, the bark is used for dyeing fish nets. Can be planted as shade or ornamental trees due to its beautiful shape.

Specimens examined: Barry 57 (P); Chevalier 39125 (P); Dournes *s.n.* (P); Hiep 458 (P); Lecomte *s.n.* (P); Maxwell 98-459 (L); Muller 1790 (P); Pierre 127, 338, 1489, 6549 (P); Poilane 148, 6086, 9463, 9823, 13783 (P), 5612 (L, P); Service Forestier 5 (P); Soejarto & Southavong 10903, 11481 (L); Thorel 996 (P); Tixier *s.n.* (P); Vidal 5909, 5945 (P); Vinot 1 (P).

10. TAMARINDUS L.

Tamarindus indica L.

Synonyms — *Tamarindus occidentalis* Gaertn.; *T. officinalis* Hook.

Vernacular name — Kham (L); Me (V).

Trees up to 25 m high, 100 cm diam., with dense rounded crown composed of many branches and twigs. *Trunks* of old trees often twisted and grooved, often fluted but not buttressed. *Bark* greyish brown, somewhat fissured and scaly, inner bark green-white. *Stipules* minute, caducous. *Petioles* and rhachis 5–16 cm long, slightly pubescent to glabrous. *Leaves* paripinnate. *Leaflets* sessile or subsessile, 10–20 pairs, elliptic, 8–30 by 3–10 mm, base rounded, unequal, apex rounded or slightly mucronate, both surfaces glabrous. *Inflorescences* terminal or axillary racemes, 5–10 cm long, rarely up to 22 cm, axis puberulous. *Bracts* and bracteoles ovate-elliptic, 5 by 3 mm, caducous, lower surface pubescent, upper glabrous. *Pedicels* up to 14 mm long, puberulous, glabrescent. *Sepals* 4, zygomorphic, imbricate, narrowly elliptic, 8–12 by 4–5 mm, inside yellow, slightly puberulous at the base, outside reddish, glabrous. *Petals* 3¹, unequal, ovate-elliptic, 10–13 by 2–6 mm, yellowish orange or cream with red veins. *Stamens* less than 10, up to 15 mm long, filaments connate for about half their length into a pubescent tube; fertile anthers 3, longitudinally dehiscent; staminodes 4 or 5, tooth-like, alternating with fertile ones. *Ovary* stipitate, linear, c. 7 mm long, pubescent, ovules 8–14; style elongate, 7 mm long, pubescent; stigma terminal, subcapitate. *Pods* indehiscent, narrowly elliptic, 5–15 by 1–3 cm; slightly curved or straight, mesocarp pulpy, acid,

1) Often 2 lower, smaller petals are mentioned; however, they may be absent or overlooked.

pale brown. *Seeds* up to 10, obovate-orbicular, 11–17 by 10–12 mm, compressed, glossy, dark brown.

Distribution — Originated from tropical Africa. Widely cultivated all over the tropics, e.g. India, Thailand, Indochina, Indonesia. In Laos: Planted in all tropical parts of the country. In Vietnam: Hà Giang, Tuyên Quang, Kiên Giang, Quang Ninh, Bac Thai, Lai Châu, Sơn La, Hoà Bình, Hà Tây, Ninh Bình, Thanh Hoá, Ninh Thuận provinces and some provinces of South Vietnam.

Habitat & Ecology — Planted around buildings, gardens, coastal areas, and along roadsides in tropical regions, altitude below 500 m. Flowering: March to July; fruiting: April to August, in Vietnam: May to December.

Uses — Wood used for firewood, for making furniture or agricultural tools, chopping blocks, charcoal for smelting gold. The acid pulpy part of the pods is used for various preserves, jams, sweets, and beverages, soup cooking. Seeds are edible and also used as vermicide. For traditional medicine the bark is used against diarrhoea. Bathing with an infusion of the boiled leaves helps against skin diseases, such as scabies. It can be used as a shade tree along streets.

Specimens examined: *Balansa* 2141 (P); *Baudouin* 27 (P); *Belval* 84 (P); *Bon* 1802, 1838 (P); *Brousmich s.n.* (P); *Chevalier* 38045 (P); *Clemens* 3809 (P); *Dupuy* 146 (P); *Dussaud* 121 (P); *Evrard* 2530 (P); *Hiep* 155 (P); *Hiroshine* 30704 (L); *Lecomte* 1393 (P); *Maxwell* 89-617 (L); *Pierre s.n.* (P); *Pindo* 9 (P); *Poilane* 7956, 11473, 26729 (L, P), 11062, 27439 (P); *Spire* 928 (P); *Squires* 805 (P); *Thorel s.n.* (P); *Tri* 29 (P); *Waas* 708 (L).

CLUSIACEAE

Trees, rarely shrubs, dioecious or monoecious. *Crown* monopodial, at least when young. Inner bark with a usually sticky yellow, sometimes clear or white exudate, occasionally very slowly appearing. *Twigs* usually quadrangular at first. *Stipules* absent, but sometimes with tiny reduced needle-like leaves resembling stipules. *Petioles* short. *Leaves* simple, opposite, and decussate, commonly leathery, with oil glands, sometimes visible as translucent dots or stripes, secondary veins usually numerous. *Inflorescences* panicles, racemes, cymes, or flowers solitary, terminal, axillary or on the twigs below the leaves. *Flower* actinomorphic, bisexual or unisexual, often with bracteoles below the calyx. *Sepals* 2–5, free or basally connate. *Petals* 2–6, free, imbricate or contorted; stamens usually numerous, free or united in fascicles; ovary superior, 1–5(–12)-locular, ovule(s) 1 or 2 per locule, style 1, stigma often large, flat. *Fruit* a drupe, berry or capsule, calyx persistent. *Seeds* without endosperm.

Distribution — Throughout all regions of the world, mainly in the tropics: c. 40 genera, c. 1000 species. In Vietnam: c. 6 genera, c. 60 species. In Laos: the number of genera and species is unknown.

MESUA L.

Mesua ferrea L.

Synonyms — *Calophyllum nagassarium* Burm.f.; *Mesua nagana* Gardner; *M. nagassarium* (Burm.f.) Kosterm.

Vernacular names — Ka thang, May lek (L); Váp (V).

Trees up to 30 m high, up to 95 cm diameter. *Trunk* straight, cylindrical, often fluted at the base. *Bark* dull brown, adherent scaly, irregular fissured; inner bark brownish red to pink or red, slowly exuding a yellow sticky sap. *Young twigs* slender, grey. *Leaves* elliptic, 4–15 by 1–3.5 cm, acute at both ends, margin entire, lower surface glaucous white, shiny, glossy green above. *Young leaves* pink. *Petiole* 4–10 mm long. *Flowers* solitary or paired, axillary, white, up to 9 cm diam., very aromatic, sepals 4, petals 4, stamens numerous, filaments yellow, ovary ovoid, superior, 2-locular, ovules 2 per locule, style long, stigma peltate. *Fruit* dark orange or purple brown, ovoid with pointed tip, 2.5–3.5 cm diam., calyx persistent. *Seed(s)* 1–4, woody, shiny brown.

Distribution — India, Sri Lanka, Myanmar, Thailand, Indochina, Peninsular Malaysia, Singapore. In Laos: Louang Prabang and Champassak provinces. In Vietnam: Kiên Giang, Vinh Phu, Hoa Binh, Thanh Hoa, and many provinces of the South, such as Gia Lai, Kon Tum, Quang Nam, Da Nang, Đông Nai, Hồ Chí Minh, and Tây Ninh.

Habitat & Ecology — Found in dense evergreen forest, altitude below 800 m, growing well in sandy soil or sand, or deep and thick clay soil. Shade tolerant when young, light demanding when mature. Successfully planted in Đông Nai province, Vietnam. *Flowering*: March to June; *fruiting*: May to August.

Uses — Timber dark red, very hard and heavy, grain fine, glossy, resistant to termites and insects, it is considered the hardest wood in Vietnam and Laos (internationally called iron wood). Used for making bridge poles, railway sleepers, and in house construction. Because of its beautiful shape and fragrant flowers, this species is usually planted around pagodas and temples in Vietnam.

Specimens examined: D'Alleizette 326 (P); Brillet 7 (P); Cadays 38252 (P); Chevalier 31298, 31312, 39128 (P); Evrard 2655 (P); Fleury 32118 (P); Forestier 30075 (P); Hiep 648, 1071 (P); Hồ s.n. (P); Hooker s.n. (P); Martin 491 (P); Pételot 4817, s.n. (P); Pierre 1 (P), 1486 (L, P); Poilane 6653, 26476 (P); Thorel 3397 (P); Vidal 4520 (P).

COMBRETACEAE

Trees, rarely shrubs or woody climbers. *Leaves* simple, opposite, subopposite, spirally arranged or rarely alternate, margin entire or wavy (*Lumnitzera*), glabrous or hairy, often conspicuously scaly and often with domatia, stipules absent. *Inflorescences* terminal or axillary, spikes, racemes, panicles or globose heads. *Flowers* usually bisexual, or unisexual and some male flowers in the same inflorescences, actinomorphic or rarely zygomorphic, 4- or 5-merous, receptacle (calyx tube) usually in two distinct parts, the lower surrounding and adnate to the ovary and an upper part varying from short to long tube terminating in the calyx-lobes. *Calyx-lobes* 4 or 5, rarely more or almost absent. *Petals* 4 or 5 or absent (*Anogeissus*, *Calycopteris*, *Terminalia*), white, yellow, orange, red, or purple, glabrous or hairy, occasionally scaly. *Stamens* usually twice as many as petals, inserted inside the upper receptacle, often in two series, anthers attached at the back. *Disc* intra-staminal usually present, sometimes absent or very small. *Ovary* completely inferior, unilocular with usually 2 (occasionally up to 6) pendulous ovules, style simple, usually free. *Fruit* a 2–5-winged, ridged or angled, sessile or stipitate, usually indehiscent drupe. *Seed* 1, pericarp usually thin and papery, sometimes leathery, rarely fleshy.

Distribution — Throughout tropics and subtropics: 18 genera, 450 species. In Laos: c. 5 genera, c. 20 species. In Vietnam: c. 6 genera, c. 25 species.

KEY TO THE GENERA

- 1a. Leaves opposite or subopposite, without glands at petiole. Inflorescences axillary and terminal, globose heads. Fruits densely packed into a spherical head, a capsule. **1. Anogeissus**
- b. Leaves opposite, subopposite, or spirally arranged, often with a pair of cup-shaped glands at the top of the petiole. Inflorescences an axillary and terminal panicle. Fruits dry and winged or a drupe, usually densely arranged along the peduncle **2. Terminalia**

1. ANOGEISSUS

Anogeissus acuminata (Roxb. ex DC.) Guill. ex Perr. — Fig. 15

Synonyms — *Anogeissus harmandii* Pierre; *Conocarpus acuminata* Roxb. ex DC.; *C. hirtus* Buch.-Ham. ex Wall.

Vernacular names — Ben mon (L); Chò nhai, Râm (V), Xa rừi, Sên (South Vietnam).

Large deciduous trees up to 40 m high, 100 cm diam., crown narrow, trunk long, straight, buttresses absent. *Twigs* slender, pendulous. *Bark* dark grey, slightly longitudinally fissured; inner bark whitish yellow. *Stipules* absent. *Petiole* 2–6 mm long, slender. *Leaves* subopposite, elliptic, 3–8 by 1.5–4 cm, acuminate or obtuse at base. Young leaves silvery with long silky hairs, venation pinnate, veins 5–9 pairs. *Inflorescences* axillary and/or terminal, globose heads, 1.3–1.8 cm diameter. *Bracteoles* caducous. *Flowers* yellowish green, bisexual, actinomorphic, 4–5 mm diameter. *Peduncle* 7–13 mm long. *Sepals* united into a cup with 5 small lobes, hairy. *Petals* absent. *Stamens* 10 in two whorls, much longer than calyx cup. *Disc* velutinous. *Ovary* 2-celled. *Fruits* densely packed into a spherical head, 1.4–1.7 cm diam., each fruit 4–6 by 3–5 mm, with 2 wings and persistent calyx tube at the top.

Distribution — India, China, Indochina, Thailand. In Laos: Vientiane, Louang Prabang, and Savannakhet provinces. In Vietnam: in almost all provinces, mainly in Hoa Binh, Ninh Binh, Thanh Hoa, Nghê An, Quang Tri, Đông Nai, Lâm Đông, and Ninh Thuận.

Habitat & Ecology — Found in tropical lowland open forest or semi-deciduous forests, altitude below 700 m. This species is growing well in deep, humus-rich, loamy soils, along streams or river banks. Usually mixed with *Pometia pinnata* and *Terminalia myriocarpa* in the north or with *Lagerstroemia calyculata*, *Pterocarpus pendatus*, and *Xylia xylocarpa* in the south. Natural regeneration is good in open places in the forest. Flowering: May to July; fruiting: July to October or until March.

Uses — Sapwood and heartwood distinctive, sapwood yellow and heartwood greyish red. Poorly resistant to water or humidity, used for light construction, furniture, and musical instruments. This species can also be planted as an ornamental tree along roads, river banks or in parks.

Note — Two specimens (*Soejarto et al. 10541, 11500*, L) are aberrant as the tube of the calyx is nearly glabrous, as described for *Anogeissus tonkinensis* Gagnep. The tubes of the other specimens are densely hairy. Whether these forms are constant and need formal recognition as a variety cannot yet be determined.



Fig. 15. *Anogeissus acuminata* (Roxb. ex DC.) Guill. ex Perr. (Combretaceae).

Specimens examined: *Balansa* 2394 (P); *Chevalier* 35684 (P); *Dubroudieu* 39166 (P); *Fleury* 30914 (P); *Forestier* 35677 (P); *Harmand* 2905 (P); *Hiep* 127 (P); *Magnen* 37 (P); *Poilane* 6, 9796 (L), 1323, 5839, 21681 (L, P), 1786, 2536, 10059, 21636 (P); *Soejarto et al.* 10541, 11500 (L).

2. TERMINALIA

Terminalia myriocarpa Van Heurck & Müll.Arg.

Synonyms — *Myrobalanus myriocarpa* Kuntze; *Pentaptera saja* Buch.-Ham.; *Terminalia saja* Steud.

Vernacular names — Kheo nua, Sam ta (L); Chò xanh (V).

Large, semi-deciduous trees up to 40 m high, up to 2 m diam., buttresses large, up to 4–5 m high. *Twigs* stout, horizontally spreading. *Young branchlets* tomentellous, sometimes rapidly glabrescent. *Bark* shallowly regularly cracked and flaky, inner

bark yellowish brown. *Stipules* absent. *Petiole* often glabrous, thick, 3–9 mm long, with a pair of cup-shaped glands at the top. *Leaves* opposite or subopposite, elliptic, 14–20 by 4–8 cm, rounded or subcordate at base, margin entire or serrate, acuminate or pointed at apex, venation pinnate, veins 15–30 pairs. *Inflorescences* axillary and terminal panicles, up to 35 cm long, peduncle 2–5 cm long, slender. *Flowers* bisexual, pinkish. *Bracts* hairy. *Calyx* lobes pubescent, triangular. *Petals* absent. *Stamens* 7–18, usually 10, free, filament slender, glabrous. *Disc* usually hairy. *Ovary* 2-celled, style cylindrical. *Fruits* numerous, arranged densely along the peduncle, 2-winged, 2–3 by 10–13 mm. *Seed* 1.

Distribution — Myanmar, China, Indochina, Thailand, Malaysia. In Laos: Louang Prabang and Vientiane provinces. In Vietnam: Lai Châu, Sơn La, Ninh Bình, Thanh Hóa, Nghệ An, and Hà Tĩnh provinces.

Habitat & Ecology — Evergreen primary and seasonal deciduous forests, altitude below 700 m. Usually mixed with *Castanopsis indica*, *Chukrasia tabularis*, *Dracontomelum dao*, *Parashorea chinensis*, *Pometia pinnata*, and *Toona sureni*. This species grows on well-drained clay or sandstone soil, along rivers or in valleys. Flowering: August to October; fruiting: November to February of the next year.

Uses — Sapwood yellowish brown, distinguishable from the brown heartwood. Grain straight, easy to nail, glue, and peel off. Used for making nice veneer and plywood in construction, musical instruments, gymnastic equipments, and for ship planking.

Specimens examined: *Balansa* 3663 (L, P), 4445 (P); *Bullock* 813 (L); *Domat* 38128 (L, P); *Dussaud* 61 (L, P); *Kerr* 21229 (P); *Poilane* 20015 (L, P); *Simons* 51 (L); *Thomson* 50, 53 (L); *Tiep s.n.* (P).

DATISCACEAE

Deciduous trees, usually with large buttresses, shrubs or herbs. *Stipules* absent. *Leaves* spirally arranged, simple or lobed, veins prominent, palmate. *Inflorescences* spikes or panicles. *Flowers* unisexual; regular, parts not overlapping. *Bracts* present. *Male flowers*: sepals connate in a tube, lobes 4–9; petals present or absent, free; stamens 4 or more, inserted on the sepals, filaments often long; sometimes with a pistillode. *Female flowers*: sepals tubular, lobes 4–8; petals absent or present; staminodes absent. *Ovary* inferior, 1-celled, carpels 3, ovules many on the wall, styles 3–8, simple or bifid, stigma variable. *Fruit* a leathery capsule, opening at apex with slits, or splitting laterally. *Seeds* numerous, very small.

Distribution — Asia and West Central America: 3 genera, 4 species. In Laos: only *Tetrameles nudiflora*.

TETrameLES R.Br.

Tetrameles nudiflora R.Br.

Synonym — *Tetrameles grahamiana* Wight

Vernacular names — Phoung, Sa phoung (L); Thung (V).

Very large, deciduous trees up to 50 m high, 100–130 cm diameter. *Trunk* straight, very tall, buttresses large, up to 7 m high. *Bark* thin, pale silver-grey, smooth and shiny

with large circular warts when younger, pock-marked when old, inner bark thick and soft, pale yellow-brown, fibrous. *Twigs* stout and spreading with large, prominent leaf scars. *Stipules* absent. Completely leafless from December to March. *Petioles* slender, 6–10 cm long. *Leaves* alternate, clustered near the end of twigs, broadly ovate or almost circular, 12–20 by 10–15 cm, base rounded or heart-shaped, margin irregularly and bluntly toothed and entire near the base, apex shortly acuminate, often slightly lobed, secondary veins 5, prominent, palmate. *Young leaves* silky hairy, mature leaves pale green, thin, minutely hairy below, distinctly raised above. Male and female flowers on different trees. *Male panicles* 8–18 cm long, much branched. *Female spikes* 10–27 cm long, mostly unbranched or with few branches. *Male flowers*: sepals 4, 1.5–2 mm long; petals absent, if present 1–4; stamens 4, about 3 mm long, opposite the sepals, surrounding a depressed disc. *Female flowers*: calyx bell-shaped, 4-toothed at the apex, sometimes with 1 or 2 smaller teeth, 3–5 mm long, petals absent, styles 4, opposite calyx teeth, 2 mm long. *Fruits* a capsule, 0.5 by 0.4 cm, pale brown, ovoid or globose, finely ridged, outer layer very thin and papery with 4 openings and remains of calyx at the top. *Seeds* many, elliptic, very small.

Distribution — India, Sri Lanka, Myanmar, South China, Thailand, Indochina, Malesia, North Australia. In Laos: Sayabouri province and Vientiane Municipality (Sa Phan Meuk). In Vietnam: Cao Bang, Ninh Binh, Hoa Binh, Thanh Hoa, Nghê An provinces, and Botanical Garden in Hồ Chí Minh city.

Habitat & Ecology — Primary, secondary, deciduous forest, altitude up to 800 m, along rivers or streams, on flat alluvial areas. The trees prefer deep, humus-rich, grey soil, sometimes growing on rocks. Flowering: January to April; fruiting: May to July.

Uses — For timber, furniture, plywood, ceiling boards, cheap planking, packing boxes, tea-chests, and matchboxes, and planted for ornamental purposes in parks.

Specimens examined: *Chevalier* 36730 (P); *Hiep* 433, 903 (P); *Thorel* 3256 (P); *Vidal* 1190 (P).

DILLENACEAE

Evergreen or deciduous trees, without latex. *Stipules* absent. *Petioles* sometimes winged. *Leaves* simple, alternate or spirally arranged, usually clustered near the ends of twigs, secondary veins straight, parallel towards the margin, margin toothed. *Inflorescences* terminal or axillary, cymes or racemes, sometimes reduced to a single flower. *Flowers* bisexual, large, regular, yellow or white. *Sepals* free, overlapping, fleshy, becoming larger and completely covering the fruits. *Petals* free, overlapping in bud, opening early in the morning but quickly fallen off. *Stamens* many in two whorls; carpels clustered on a central cone. *Fruits* fleshy, covered by persistent, overlapping sepals, later splitting open.

Distribution — In the Old and New World: 10 genera, c. 350 species. In Laos: only *Dillenia* with 5 species. In Vietnam: 2 genera, 6 species.

DILLENIA L.

Dillenia ovata Wall. ex Hook.f. & Thomson

Vernacular names — May san, San nhai (L); Sô (V).

Deciduous trees up to 30 m high, up to 100 cm diam., with rather knotted trunk and low-attached crown. *Bark* reddish or brown, flaky, inner bark reddish brown. *Twigs* slender, softly hairy. *Petioles* 2–4.5 cm long. *Leaves* ovate or elliptic, 10–20 by 7–13 cm, base rounded to acute, often unequal, margin finely toothed, apex rounded to obtuse, secondary veins 16–25 pairs, venation scalariform, densely velvety hairy beneath. *Flowers* solitary, terminal, very large, up to 16 cm diam., petals bright yellow with slight odour. *Pedicel* 0.5–1.7 cm long. *Sepals* ovate, 25–30 by 15–25 mm. *Petals* ovate, 7 by 5.5 cm. *Stamens* in 2 distinct groups: outer ones curved, 16–20 mm long; inner ones reflexed at apex, 25 mm long. *Carpels* 10–12, 6–7 by 2–2.5 mm, style 20 mm long, ovules many. *Fruits* indehiscent, pale yellow, globular, 4–6 cm diameter. *Seeds* blackish brown.

Distribution — Indochina, Thailand, Peninsular Malaysia, Indonesia (Sumatra, Bangka). In Laos: Savannakhet province. In Vietnam: Khanh Hoa, Quang Tri, Ninh Thuân, Đông Nai, Dac Lac, Quang Nam, Tây Ninh, and Bà Rịa-Vung Tàu provinces.

Habitat & Ecology — In open areas, mixed evergreen, deciduous, or dry Dipterocarp forest, or savannah, altitude up to 1500 m. Flowering: January to April; fruiting: May to September.

Uses — Trees are planted as ornamentals. The wood is suitable for beams, planks, and furniture. The fruits are eaten as vegetable or in jellies. The bark is used against diarrhoea.

Specimens examined: *Chevalier* 30935, 35826, 35827, 36757, 39251, 39633 (P); *Dau* 85 (P); *Fleury* 30084 (P); *Hiep* 667 (P); *Muller* 994, 1711, 1971 (P); *Pierre* 44, 2038, 2040 (P); *Poilane* 1287, 5817, 6694, 11822, 12073, 24687, 30065, 32373, 40890 (P); *Squires* 775 (P); *Vinot* 29 (P).

DIPTEROCARPACEAE

Medium to large trees, usually with resin, buttressed, and often (if mature trees) with flaky or fissured bark. Some or most parts with a tomentum of fascicled hairs, or sometimes single or peltate hairs. *Stipules* present, paired, large or small, persistent or caducous, leaving small to amplexicaul scars. *Leaves* simple, alternate, margin entire or sinuate, often with domatia, venation pinnate, tertiary veins scalariform or reticulate. *Inflorescences* panicles or racemes, rarely cymose, terminal or axillary. *Bracts* or bracteoles paired, small or large, persistent or caducous. *Flower* bisexual, actinomorphic, 5-merous. *Calyx* 2–5 sepals, persistent, usually greatly enlarging into wing-like lobes in fruit, sepals either free to base, imbricate in bud, or fused at base, forming a cup or tube enclosing the fruit, adnate to or free from it. *Corolla* 5-merous, contorted, base connate or free, usually hairy. *Stamens* usually numerous, filaments free or connate, anther 2-celled. *Ovary* superior or inferior, 3- rarely 2-locular, ovules 2 or 3 in each locule. *Fruit* indehiscent, 1-seeded, nearly always winged, wings 2–5. *Nut* enclosed between the grown calyx lobes (ovary superior) or under the calyx lobes (ovary inferior).

Distribution — Throughout the tropical regions of the world, mainly in South East Asia: c. 75 genera, 850 species. In Laos: 6 genera, 27 species. In Vietnam: c. 6 genera, c. 55 species.

KEY TO THE GENERA

- 1a. Petiole swollen at top 2
 b. Petiole not swollen at top 4
 2a. Bark lamellated. Leaves in bud not folded, flat, intramarginal vein present
 **1. Anisoptera**
 b. Bark not lamellated. Leaves folded in bud, usually distinctly undulate between
 each vein, intramarginal vein absent 3
 3a. Fruit with 2 large and 3 small or 5 much reduced wings. Stipules large, stipule scar
 amplexicaul **2. Dipterocarpus**
 b. Fruit with 5 equal or nearly equal wings. Stipules small, stipule scar short, not
 amplexicaul **4. Parashorea**
 4a. Fruit with 3 large and 2 small wings **5. Shorea**
 b. Fruit with 2 large and 3 small wings or wings absent 5
 5a. Stilt roots absent. Tertiary venation reticulate, sometimes with glands at the junction
 of secondary veins **6. Vatica**
 b. Stilt roots present. Tertiary venation scalariform, glands absent **3. Hopea**

1. ANISOPTERA Korth.

Anisoptera costata Korth. — Fig. 16

Synonyms — *Anisoptera cochinchinensis* Pierre; *A. oblonga* Dyer; *A. robusta* Pierre

Vernacular names — Bak, Mai bak (L); Vên vên, Vên vên trắng, Vên vên xanh (V).

Large evergreen trees up to 40 m high, up to 1.7 m diam., bole straight, buttresses cylindrical, few, up to 4 m tall, thick, rounded, straight, continuing up the bole as ribs up to 10 m. *Bark* greyish brown, glabrous when young, deeply fissured when old, outer bark thick, inner bark up to 3 cm thick, lamellated, cream-yellow. Sapwood cream-white, heartwood pale yellow with silica. *Twigs* at first flat or angular, densely yellow stellately hairy, becoming minutely striated or smooth. *Stipules* c. 8 mm long, c. 3 mm wide, hastate, acute, early caducous. *Petioles* swollen, 2–4 cm long, hairy, thickened at top. *Leaves* elliptic to obovate, 6–20 by 3–11 cm, apex with a short obtuse point, base rounded or nearly cordate, glabrous on upper surface, stellate caducous tomentose on lower surface, venation pinnate, secondary veins 15–20 pairs, intramarginal vein present. *Inflorescences* a raceme, axillary or terminal, 10–15 cm long, stellate hairy. *Sepals* 5, tomentose, lobes triangular. *Petals* 5, obtuse, narrow elliptic, cream-coloured. *Stamens* 25–35, anthers oblong, tapering apically. *Ovary* cylindrical, 2-locular, ovules 2 in each locule, style short, pubescent at base. *Fruit* subglobose, brown, 1–1.5 cm diam., style persistent at top, two fruit-wings large, 10–16 by 1–1.5 cm, and the three others shorter, 1.5–2.5 by 0.2–0.5 cm.

Distribution — Indochina, Thailand, Malesia. In Laos: Vientiane, Bolikhamsai, Savannakhet, and Champassak provinces. In Vietnam: Kon Tum, Gia Lai, Dac Lac, Bien Hoa, Bà Rịa-Vung Tàu, Ninh Thuận, Quang Nam-Dà Nang, Long An, Tây Ninh, Đông Nai, Hồ Chí Minh, and Kiên Giang provinces.

Habitat & Ecology — In moist or slightly dry forests, sometimes gregariously growing in pure stands. Usually mixed with *Dipterocarpus* spp. and *Shorea* species. Flowering: November to March; fruiting: February to May.




Fig. 16. *Anisoptera costata* Korth. (Dipterocarpaceae).

Uses — Wood whitish yellow with indistinctive sapwood and heartwood, used for making interior furniture, flooring, ship planking, veneer, and plywood. Resin from wood soft at first, then become hard, grey, strongly smelling.

Specimens examined: *D'Alleizette s.n.* (L); *Aubreville 3* (L); *Bodeuneuve 36800* (P); *Chevalier 35742, 35813, 36580, 39176* (P); *Fleury 29935* (P); *Forestier 36855* (P); *Hoi 234* (P); *Kerr 9543, 10042, 17029* (L), 20736 (P); *Lefèvre 325* (P); *Manein 32* (P); *Parreau 174* (P); *Pierre 149, 1549, 1565, 1566, 1571* (P); *Poilane 830, 12057, 12067* (L, P), 7516, 11724, 11881 (P); *Put 642* (L); *Sigaldy 44* (P); *Thorel 727, 2696* (P); *Tremessaignes 36734* (P); *Vidal 1281* (P), 4419 (L, P).

2. DIPTEROCARPUS C.F. Gaertn.

KEY TO THE SPECIES

- 1a. Leaves ovate to elliptic, 14–25 cm long. Stipules greyish yellow. Fruit globose, with 5 ridges up to 8 mm broad, 2 larger wings 10–14 cm long **1. *D. alatus***
 b. Leaves broadly elliptical, 16–40 cm long. Stipules red. Fruit subglobose, without ridges, 2 larger wings 15–20 cm long **2. *D. retusus***

1. *Dipterocarpus alatus* Roxb. & G. Don — Fig. 17

Synonyms — *Dipterocarpus gonopterus* Turcz.; *D. incanus* Roxb.; *D. lemeslei* Vesque

Vernacular names — Nhang, Nhang khao (L); Dầu rái, Dầu con rái, Dầu nước (V).

Large trees up to 45 m high, up to 1.5 m diameter. *Trunk* cylindrical, straight. *Crown* umbrella-like. *Bark* whitish grey, inner bark yellowish brown, resinous. *Leaves* ovate to elliptic, 14–25 by 6–15 cm, folded in bud, usually distinctly undulate between each vein, base cuneate to rounded, margin entire or undulate, apex acute or acuminate, secondary veins 11–20 pairs, prominent beneath, tertiary venation scalariform, sparsely pubescent above, densely pubescent beneath. *Petioles* swollen at top, 2.5–4 cm long, densely hairy. *Stipules* greyish yellow pubescent, stipule scar amplexicaul.

Inflorescences axillary, racemes up to 12 cm long. *Flowers* bisexual, pale pink; sepals 5; petals 5; stamens about 30. *Ovary* superior, ovoid, densely hairy, 3-locular. *Fruit* globose, with 5 ridges 8 mm broad, 2 larger wings 10–14 by 2–3 cm, three smaller ones 1–1.4 by 1–1.3 cm, wings red when young.

Distribution — India and South East Asia. In Laos: Vientiane, Savannakhet, Saravane, and Champassak provinces. In Vietnam: from Quang Nam-Dà Nang province southwards, most common in Đông Nai, Long An, and Bà Rịa-Vung Tàu provinces.

Habitat & Ecology — Tropical evergreen or monsoon forest, altitude below 800 m, mainly 200–500 m, often growing gregariously in small pure stands along streams and river banks. Usually mixed with *Dalbergia cochinchinensis*, *Dipterocarpus turbinatus*, *Pterocarpus macrocarpus*, and *Sindora siamensis*. Light demanding tree, but shade tolerant when young. Natural regeneration is good, especially along rivers or on moist, flat land. Flowering: November, December; fruiting: April, May.




Fig. 17. *Dipterocarpus alatus* Roxb. & G. Don (Dipterocarpaceae).

Uses — A very common timber tree in South Vietnam. Wood rather hard with fine grains, easy to saw and polish. Used in light construction, for furniture and household implement. Its resin is used in the paint industry. This species is a very common shade tree in Hồ Chí Minh city and in many towns of the southern provinces. Also planted in Hà Nội capital from the beginning of this century.

Specimens examined: *Abriae* 35773 (P); *D'Alleizette* s.n. (L); *Boudeneuve* 36797 (P); *Chevalier* 30589, 36783, 36847, 37284, 39150 (P), 31351 (L, P); *Evard* 786 (P); *Fleury* 29936 (P); *Forestier* 35757 (P); *Lefèvre* 594 (P); *Pierre* 1693 (L, P), 1717, 5047 (P); *Poilane* 555 (P), 9520 (L, P), 11566 (L); *Robinson* 1350 (L), 1351 (P); *Schmid* s.n. (P); *Thorel* 962 (P); *Vinot* 36 (P).

2. *Dipterocarpus retusus* Blume — Fig. 18

Synonyms — *Dipterocarpus macrocarpus* Vesque; *D. pubescens* Koord. & Valetton; *D. tonkinensis* A. Chev.

Vernacular names — Mai nhang dong, Nhang dong, Pak munung (L); Chò nâu, Chò nển (V).

Large deciduous trees up to 40 m high, 1 m diameter. *Trunk* straight, cylindrical. *Bark* of trunk and branches densely lenticellate. *Twigs* covered with dense, caducous hairs. *Leaves* broadly oval 16–40 by 7–25 cm, coriaceous, folded in bud, usually distinctly undulate between each vein, base rounded or cordate, margin entire or undulate, apex obtuse, leaf blade glabrous on both surfaces, midrib densely hairy, secondary veins 15–20 pairs, prominent beneath, densely covered with stellate hairs, tertiary venation scalariform. *Petioles* swollen at top, 2.5–4 cm long, pubescent when young, glabrous and blackish when mature. *Stipules* large, red, 8–12 cm long, stipule scar amplexicaul. *Inflorescences* axillary, panicles to 10 cm long. *Flowers* bisexual, bud 3 by 1 cm. *Calyx* tube subglobose, 2.5 by 2 cm. *Sepals* 5. *Petals* 5. *Stamens* 30, exceeding style at anthesis, filament short, anthers long, linear, tapering. *Ovary* ovoid, densely pubescent, 3-locular. *Fruits* subglobose, 2–3 cm diam., smooth, without ridges, 2 large wings 15–20 by 2–3 cm, with prominent veins, other 3 wings much reduced, auricular, about 5–8 mm long, wings red when young.

Distribution — India, Myanmar, Indochina, Thailand, Malaysia, Indonesia. In Laos: Vientiane province. In Vietnam: in the north, Tuyên Quang, Vinh Phu, Hà Tây, Bắc Thái, and Hà Bắc provinces. Nowadays this species is planted in many provinces in the north of Vietnam.




Fig. 18. *Dipterocarpus retusus* Blume (Dipterocarpaceae).

Habitat & Ecology — Tropical evergreen or monsoon forest, altitude 100–1000 m, mainly 300–700 m, on loam, deep, fertile, and well-drained soils. Usually mixed with *Magnolia conifera*, *Michelia mediocris*, *Parashorea chinensis*, and *Terminalia myriocarpa*. Natural regeneration is good in secondary forest. Flowering: January, February; fruiting: April to July.

Uses — The wood is soft, susceptible to termite and insect attack. Used for construction and making furniture. Can be planted as an ornamental and shade tree along streets or in parks.

Specimens examined: *Brillet s.n.* (P); *Chevalier* 4028, 37386, 37387, 37521, 41028 (P), 37546 (L), 38004, 41029 (L, P); *Fleury* 37546 (P); *Forestier* 39699 (P); *Gilbert* 38039, 39716 (P); *Kerr* 18753 (L), 21242 (L, P); *Pételot* 1026 (P); *Pierre* 8408 (P); *Van Beusekom* 195 (L); *Vidal s.n.* (P).

3. HOPEA Roxb.

KEY TO THE SPECIES

- 1a. Bark greyish brown. Twigs glabrous. Leaves glabrous on both surfaces. Fruit 1–1.5 cm diam., two larger wings 6–10 cm long **1. *H. chinensis***
- b. Bark blackish brown. Twigs covered with the grey stellate hairs. Lower leaf surface with domatia at the axils of secondary veins. Fruit 0.6–0.8 cm diam., two larger wings 4–7 cm long **2. *H. odorata***

1. *Hopea chinensis* (Merr.) Hand.-Mazz. — Fig. 19

Synonyms — *Hopea hongayensis* Tardieu; *Shorea chinensis* Merr.

Vernacular names — Khen si (L); Sao hòn gai, Táu mặt quì (V).

Large evergreen trees up to 40 m high, 60 cm diam., stilt roots usually present. *Bark* greyish brown, when old peeling off in large pieces, leaving concentric rings on the trunk. *Twigs* glabrous, blackish brown when dry. *Leaves* narrowly ovate, 9–21 by 3–6 cm, glossy-green above, paler beneath, glabrous on both surfaces, base rounded, margin entire, apex 1–2.5 cm long, pointed, midrib slightly evident above, and clearly evident beneath, secondary veins 7–10 pairs, curved near the margin. *Petioles* 0.6–1 cm long. *Inflorescences* axillary, panicles, 8–10 cm long, consisting of many branches, with 3 or 4 flowers. *Flowers* bisexual, pedicel very short. *Sepals* 5, imbricate, triangular. *Petals* 5, contorted, 7–9 mm long, glabrous inside, pubescent outside. *Stamens* 15, filaments flat, 1 mm long, often connected with the petals, anthers 2 mm long, connective 1–1.5 mm long. *Ovary* superior, ovoid, with stylopode, 3-locular, style short, stigma obtuse. *Fruits* ovoid, glabrous, 1–1.5 by 0.6–1 cm, two larger wings, 6–10 by 1.4–1.7 cm, with 9–13 parallel veins, 3 small wings, 6–8 by 4–6 mm, wings pink when young.

Distribution — South China, Laos, Vietnam. In Laos: Louang Prabang, Xieng Khouang, Vientiane, Bolikhamxai, and Khammouane provinces. In Vietnam: Hà Bac, Quang Ninh, Lang Son, Tuyên Quang, Ninh Binh, and Nghệ An provinces.

Habitat & Ecology — Tropical evergreen forest, altitude below 950 m. Usually mixed with *Hopea mollissima*, *Vatica fleuryana*, and *Vatica odorata*. Shade demanding when young, then gradually becoming light demanding when older. Natural regeneration very good. Flowering: June to October; fruiting: February, March.




Fig. 19. *Hopea chinensis* (Merr.) Hand.-Mazz. (Dipterocarpaceae).

Uses — Timber hard, heavy, and resistant to termites and insects, used in house pillar making and in construction.

Specimens examined: *Bonnel s.n.* (P); *Casabianca* 39628 (P); *Tsang* 27053, 27260, 30154, 30465 (P).

2. *Hopea odorata* Roxb.

Synonym — *Doona odorata* (Roxb.) Burck

Vernacular names — Khen, Khen hua (L); Sao, Sao đen, Sao cát, Sao nghệ, Sao xanh (V).

Large evergreen trees up to 40 m high, 80 cm diameter. Bole straight, cylindrical up to 25 m high, stilt roots usually present. *Bark* surface scaly, blackish brown, outer bark rather thick, inner bark dull yellow. *Twigs* and petioles covered with grey stellate hairs, later glabrous. *Leaves* ovate or elliptic, 8–14 by 3–6 cm, base broadly cuneate, unequal, margin entire, apex acute, secondary veins 8–12 pairs, evident beneath, lower leaf surface with domatia at the axils of secondary veins, tertiary veins scalariform. *Petioles* 1–1.8 cm long. *Stipules* small, early caducous. *Inflorescences* axillary or terminal panicles, up to 17 cm long, hairy, consisting of many branches, with 3–9 flowers. *Flowers* bisexual. *Sepals* 5, hairy on both sides, outer 2 lanceolate, inner 3 broadly ovate. *Petals* pale cream-yellow, hairy outside. *Stamens* 15, filaments slender, anthers narrowly ellipsoid. *Ovary* superior, ovoid, hairy. *Fruits* ovoid, 6–8 mm diam., 2 larger wings 4–7 by 1–1.6 cm, 3 shorter ones 3–4 by 3–4 mm.

Distribution — Myanmar, India, Indochina, Thailand. In Laos: Xieng Khouang, Vientiane, Khammouane, Savannakhet, Champassak, and Attopeu provinces. In Vietnam: most provinces of southern Vietnam, from Gia Lai, Kom Tum provinces southwards. Recently this species is planted in many provinces in North and South Vietnam.

Habitat & Ecology — Tropical evergreen forest, altitude below 600 m. This species demands wet and deep soil. Natural regeneration is good under thin forest cover. Flowering: February, March; fruiting: April to July.

Uses — Wood is a good and valuable timber in Vietnam and Laos. The timber is hard, heavy, and resistant to termites and insects, used for making furniture, floors, and railway sleepers. Trunk contains much resin and gums. This species is planted as a shade tree in many cities in Vietnam.

Specimens examined: *D'Alleizette s.n.* (L); *Bejaud 22999* (L); *Chevalier 31751, 32032, s.n.* (L); *Collins 1400* (L); *Kerr 15788, 17934, 19905* (L); *Nimanthong 67* (L); *Poilane s.n.* (L); *Put 608* (L); *Vinol 19* (L).

4. PARASHOREA Kurz

Parashorea chinensis H.C. Wang — Fig. 20

Synonyms — *Shorea chinensis* (H.C. Wang) H. Zhu; *S. wangtianshuea* Y.K. Yang & J.K. Wu

Vernacular names — Chò chì (V), Mạ hao, Mạ Khay (Mường, V).

Large trees up to 50 m high, 100 cm diam. (sometimes up to 200 cm). *Trunk* cylindrical. *Bark* grey with slight longitudinal fissures, inner bark slightly aromatic, containing aromatic resins. *Leaves* elliptic to ovate, 7–20 by 2.6–8 cm, base rounded, margin entire, apex attenuate or shortly caudate, glabrous and shining on the upper side except on the veins, stellate hairy and with dot-like glands on the lower surface, folded in bud, usually distinctly undulate between each vein, secondary veins 13–20 pairs. *Petiole* 1–1.5 cm long, with stellate hairs. *Stipules* ovate, pointed at the top, cordate at the base, 2 embracing the twigs, with stellate hairs, stipule scar short. *Inflorescences* terminal or axillary panicles, 7–17 cm long, rhachis cylindrical, densely soft stellately hairy. *Bracts* ovate-oblong, 10–15 by 5–6 mm, usually 2 embracing the flower. *Sepals* imbricate, finely velutinous on both sides. *Petals* ovate, yellowish, with whitish indument outside. *Stamens* 12–15, filaments 2–4.5 mm long, the inner whorl much longer, anthers 1.6 mm long, apiculate, inner ones shorter, anther prolongation 1–1.2 mm long, subulate. *Ovary* superior, ovoid, 3–5 by 2–2.5 mm, densely velutinous, 3-locular, ovules 2 in each locule, style glabrous. *Fruits* ovoid to ellipsoid, densely covered by grey scales, calyx lobes glabrous, a little bit unequal in length, embracing the fruit, 3 larger ones elliptical-oblong, 6–10 by 1.2–2 cm, 5–7 veins, the two smaller ones 3.5–7 by 0.6–1 cm.

Distribution — Endemic in South China and North Vietnam. In Vietnam: most provinces of North Vietnam from Lai Châu, Lào Cai to Quang Binh provinces, mainly in Hà Giang, Tuyên Quang, Kien Gian, Ninh Binh, Thanh Hoa, Nghệ An, and Hà Tĩnh provinces.

Habitat & Ecology — In Vietnam in valleys, along rivers, in evergreen forests, altitude 100–1000 m, but more often 300–700 m. Usually mixed with *Aglaia gigantea*, *Dracontomelum dao*, *Pometia pinnata*, *Saraca dives*, *Terminalia myriocarpa*, and *Vatica odorata*. Flowering: April to June; fruiting: July to September.




Fig. 20. *Parashorea chinensis* H.C. Wang (Dipterocarpaceae).

Uses — The wood is hard, heavy, with a fine grain, yellow to pink, resistant against termite attack. The excellent wood is used for construction and making furniture. This species can be planted as shade tree along streets because of its beautiful shape.

Specimens examined: *Cuong 275* (L); *Dung s.n.* (P); *Favrot s.n.* (P); *Schmid s.n.* (P).

5. SHOREA Roxb. ex C.F. Gaertn.

Shorea siamensis Miq. — Fig. 21

Synonyms — *Pentacme siamensis* (Miq.) Kurz; *Shorea bracteata* Pierre ex Laness.; *S. mekongensis* Pierre ex Laness.

Vernacular names — Hang (Vientiane, L), Pahu (Louang Prabang, L); Cẩm liên, Cà chắc xanh (V).

Large deciduous trees up to 30 m high, 80 cm diam. or more. *Trunk* straight. *Bark* grey, deeply fissured, 1.5 cm thick, inner bark reddish brown with yellowish brown resin. *Leaves* ovate or oval, 10–22 by 7–16 cm, base cordate, margin entire, apex obtuse or

acuminate, young leaves reddish brown with stellate hairs, mature leaves usually dull green, glabrous above, tomentose or glabrous beneath, secondary veins 10–16 pairs, raised beneath, tertiary venation scalariform. *Petioles* swollen, 2–4 cm long, slender and slightly flattened. *Stipules* 1–2 cm long, narrowly ovate, falling early. *Inflorescences* a terminal or axillary raceme, up to 20 cm long. *Flowers* bisexual. *Sepals* 5, red, narrowly ovate, hairy outside. *Petals* 5, with recurved tips, fused at base and falling together as a rosette. *Stamens* 15 in two whorls, 10 in outer and 5 in inner whorl, filament short, anthers glabrous with long narrow tips, 5–6 mm long. *Ovary* superior, smooth, 3-locular, ovules 2 in each locule, style glabrous, 5–6 mm long, with single stigma. *Fruit* ovoid, 3 larger wings 7–10 by 1–1.5 cm, 2 shorter ones 5–7 by 0.8–1 cm.

Distribution — Myanmar, Indochina, Thailand. In Laos: Louang Prabang, Xieng Khouang, Vientiane, Khammouane, Savannakhet, Saravane, and Attapeu provinces. In Vietnam: most provinces of southern Vietnam, especially in Quang Tri, Gia Lai, Kom Tum, Quang Ngai, Binh Dinh, Phu Khanh, Dac Lac, Lâm Đông, Ba Ria, Bien Hoa, Sai Gon, Tây Ninh, Phan Rang, and Phuoc Long provinces. Recently this species has been planted in many provinces in North and South Vietnam.

Habitat & Ecology — Common in dry deciduous Dipterocarp forest in pure stands or mixed with *Shorea obtusa* and *Terminalia* species. A xerophytic tree, mainly grow-



Fig. 21. *Shorea siamensis* Miq. (Dipterocarpaceae).

ing on shallow poor, sandy, and stony soils. Shedding leaves in late February to early March. Flowering: February to April, before new leaves appear; fruiting: April, May.

Uses — Sapwood and heartwood distinctive. Sapwood red, heartwood brownish red, very heavy and solid. Used for long term construction and valuable furniture.

Specimens examined: *D'Alleizette* 717, *s.n.* (L); *Chevalier* 30030, 30953, 35769, 36578, 36586, 36655, 39147, 39148 (P), 35781 (L); *Dournes* 185 (P); *Dussaud* 41 (P); *Forestier* 35771, 36857, 40918 (P); *Harmand* 1336 (P); *Hiep* 369 (P); *Millet* 38405, 38406 (P); *Pierre* 649 (L), 796 (L, P); *Poilane* 1346, 8928, 9143, 9613 (P), 13740 (L, P); *Schmid* 912 (P); *Thorel* 3096 (P); *Tixier* 15 (P); *Vidal* 714, 814, 1285 (P); *Vinot* 35789 (P).

6. VATICAL.

Vatica odorata (Griff.) Symington — Fig. 22

Synonyms — *Sunaptea odorata* Griff.; *Vatica fleuryana* Tardieu; *V. tonkinensis* A. Chev.

Vernacular names — Chik dong, Si daeng (L); Táu (V).

Large trees up to 40 m high, up to 1 m diameter. Bole straight, cylindrical. *Bark* grey, inner bark white, very fibrous. *Young twigs*, leaf buds and petioles densely persistently pale yellowish brown to fulvous tomentose. *Buds* ovoid, 2 by 1.5 mm. *Leaves* elliptic to ovate, 4–16 by 2–5.5 cm, thinly coriaceous, base obtuse or cuneate, margin entire, apex acuminate, secondary veins 8–15 pairs, prominent beneath, tertiary venation reticulate. *Petioles* 4–12 mm long. *Stipules* 3–8 mm long, caducous. *Inflorescences* terminal or axillary panicles, up to 10 cm long, hairy. *Flower bud* to 8 mm long. *Calyx* densely shortly pale grey-brown tomentose, sepals 5, tomentose on both sides. *Petals* 5, hairy to glabrous, twisted when young. *Stamens* 12, connective obtuse. *Ovary* superior, hairy, 3-locular, 1 ovule per locule, style long, stigma divided into 3 triangular lobes. *Fruits* globose, tomentose, style persistent, two larger wings 4–6.5 by 1–2 cm, with 5 parallel veins, 3 shorter ones 1–1.5 by 0.3–0.4 cm.

Distribution — Myanmar, Indochina, Thailand, Malesia. In Laos: Vientiane, Savannakhet, Saravane, and Champassak provinces. In Vietnam: Tuyên Quang, Hà Giang, Lang Son, Bac Thai, Kiên Giang, and Quang Ninh provinces.

Habitat & Ecology — Primary forests, altitude 300–900 m, slow growing, suitable for reforestation on (sandy) clay and sloping hill sides. Usually mixed with *Castanopsis* spp., *Cinamomum* spp., *Machilus odoratissima*, *Madhuca pasquieri*, and *Garcinia cowa*. Light demanding, natural regeneration is very prolific under a forest cover of 70–90%. Flowering: May, June; fruiting: September, October.

Uses — Wood hard and heavy. Sapwood greyish white, heartwood brown. Timber valuable, resistant to termites and insects, much used in construction, making sleepers, bridges, ferry boats, and planks.

Note — Smitinand, Vidal & Hô in Lecomte, Flore du Cambodge, du Laos et du Vietnam 25 (1990) 3 treated *Vatica tonkinensis* as *Vatica mangachapoi* Blanco subsp. *obtusifolia* (Elmer) Ashton, and *Vatica fleuryana* Tardieu as a synonym of *Vatica diospyroides* Symington. At this stage we are not able to make a decision with solid arguments, as we are unable to study the types and related material. This is the reason that we take a conservative view and treat these names as synonyms of *V. odorata*.




Fig. 22. *Vatica odorata* (Griff.) Symington (Dipterocarpaceae).

Specimens examined: *D'Alleizette s.n.* (L); *Chevalier* 31720, 31990, 36775, 39051, *s.n.* (L), 36780, 36785, 39195 (P), 38914 (L, P); *Geesink et al.* 7303 (L); *Helfer* 142, 716, 717 (L); *Maxwell* 76-193, 94-457, 94-674 (L); *Pierre* 173, 1577, 1578, 1583, 1584 (P), 1579 (L, P), 1580 (L); *Poilane* 11988, 13765 (P); *Prumette* 36833 (P); *Vidal* 1735, 2726, 6001 (P).

EBENACEAE

Trees or shrubs, latex absent. *Bark* usually black, hard and brittle, with smooth, fissured, or cracked surface. *Branches* usually pseudowhorled. *Leaves* simple, alternate, drying yellowish, dark brown or black dots on one or both sides, margin entire. *Stipules* absent. *Inflorescences* cymose (with a basic cymose unit of 3 or 5 or 7 flowers, sometimes reduced to one flower, sometimes branches carrying multiple units). *Flowers* unisexual, regular, 3–5(–8)-merous. *Sepals* mostly united at base, sometimes free, the lobes imbricate or valvate, persistent in fruit. *Petals* basally united into a tube. *Male flower*: stamens (3–10)–20(–100), commonly inserted at base of corolla tube, sometimes on the receptacle, often in two whorls, filaments free or united in pairs, triads, or larger fascicles, or even into a single column, anthers basifixed, usually longer than the filaments, 2-locular, dehiscing by longitudinal slits. Pistillode present or absent. *Female flowers*: ovary superior, multilocular, 2–8 carpels as indicated by the number of styles or stigmas lobes, ovules pendulous, anatropous, bitegmic, staminodes reduced to a single whorl of sterile epipetalous lobes or absent. *Fruit* a berry, pericarp fibrous or fleshy, calyx persistent in fruit. *Seed(s)* 1–16, arranged in a single whorl around central axis, embryo straight or slightly curved, endosperm present.

Distribution — Worldwide: 2 genera, c. 500 species. Except for a few species of *Euclea* (in Africa) the bulk of the family belongs to *Diospyros*, which is almost completely tropical and subtropical, a few species occur in temperate climate. In Laos: only *Diospyros* with c. 15 species. In Vietnam: only *Diospyros* with c. 62 species.

DIOSPYROS L.

Diospyros mun A. Chev. & Lecomte

Vernacular names — Nang dam (Northern, L), Lang dam (Vientiane, Savannakhet, L); Mun, Mùng, Mẹt (V).

Medium trees up to 18 m high, 40 cm diameter. Bole cylindrical, early branched. *Bark* black with shallow, longitudinally fissures. *Crown* dense. *Leaves* ovate, 7–10 by 3–5 cm, base acute, apex acuminate, light green, venation pinnate, often with black glands below. *Petiole* slender, smooth, 0.8–1.2 cm long. *Flowers* unisexual. *Male flowers* in cymes in the axils of leaves, each bearing 3–5 yellow flowers, 4-merous, stamens 16, arranged into 2 whorls, filaments inserted on petals. *Female flowers* solitary, calyx cup-shaped, 4-merous, styles 3, bifurcate at the top. *Fruit* subglobose 1–2 cm diam., glossy-green, becoming black when mature, sepals 4, persistent. *Seeds* 6–8 arranged in a single whorl around the central axis.

Distribution — Indochina. In Laos: Vientiane and Savannakhet. In Vietnam: in the northern provinces: Hà Giang, Tuyên Quang, Lang Son, Hoa Binh, Ninh Binh, Thanh Hoa, Nghệ An, Hà Tĩnh, Quang Binh, and Khanh Hoa (Cam Ranh Distr.) provinces.

Habitat & Ecology — On limestone mountains, altitude below 800 m. In the south on yellow ferrallitic soil. Light demanding and drought tolerant tree, growing on poor soil. Slow growing, regenerations by seeds or suckers near the stump. Flowering: June to September; fruiting: July to November.

Uses — Very precious timber, heartwood black, hard and heavy, becoming glossy black when dry. Used for flower pots, statues, and especially for making chopsticks.

Fresh seeds contain black dye for dyeing silk, cloth, and other materials. This species is rated as vulnerable in the Vietnam Red Data Book.

Specimens examined: *D'Alleizette s.n.* (L); *Chevalier 34, 35887* (P); *Fleury 30036* (P); *Poilane 57, 133, 134, 8779* (P).

EUPHORBIACEAE

Herbs, shrubs, small, medium, or large trees, rarely climbers. *Bark* usually smooth, inner bark occasionally with white latex or watery sap. *Leaves* simple, mostly alternate, rarely opposite, sometimes palmate or lobed (*Bischofia* and *Hevea*). *Stipules* usually present, but often small and caducous; petiole often long, slender, swollen at the top and sometimes with glands near insertion with blade, venation of leaves pinnate or palmate. *Flowers* usually tiny, green, yellow, or white, in cluster, spikes, racemes or panicles, terminal or axillary, unisexual, dioecious or monoecious, structure of flowers extremely variable; sepals 2–5, overlapping or not; petals usually absent, or as many as sepals; disc often present; stamens 1 to many, free or variously fused; male flowers often with pistillode; ovary superior, 2- or 3-locular, 1 or 2 per locule, styles 2 or 3, sometimes joined at base. *Fruit* typically a smooth, 3-lobed, woody capsule, splitting, often explosively into 3 or bivalved parts, or not lobed, or fleshy to leathery, not splitting and then often with a stone. *Seeds* always few, endosperm oily.

Distribution — Throughout the world except the coldest regions: c. 300 genera, 8100 species. This family is the sixth biggest family of flowering plants (after Orchidaceae, Compositae, Leguminosae, Gramineae, and Rubiaceae). In Laos and Vietnam: c. 62 genera, c. 416 species.

KEY TO THE GENERA

- 1a. Leaves simple 2
- b. Leaves trifoliolate **3. Bischofia**
- 2a. Hairs simple, slash of the bark without sap. Petiole usually kneed at both ends. Fruit a berry **2. Baccaurea**
- b. Hairs stellate, slash of the bark with sap. Petiole not kneed at the ends, fruit a drupe 3
- 3a. Sap clear. Leaves usually bearing 2 crap-eye-shaped glands at the lower side of the base of the lamina. Bracts present. Petals absent. Fruit a small drupe, 1–2 cm diameter **4. Endospermum**
- b. Sap pinkish. Leaves bearing 2 glands at the upper side of the base of the lamina. Bracts absent. Petals present. Fruit a large drupe, 4–6 cm diameter **1. Aleurites**

1. ALEURITES J.R. Forst. & G. Forst.

Aleurites moluccana (L.) Willd. — Fig. 23

Synonyms — *Aleurites ambinux* Pers.; *A. lobata* Blanco; *Jatropha moluccana* L.

Vernacular names — Nam man (L); Lai (V).

Trees up to 30 m high, up to 70 cm diameter. *Trunk* straight and cylindrical. *Bark* smooth, greenish grey, slash with watery, clear sap, inner bark pink. *Branchlets* angular, verticillate, covered with fulvous stellate, tomentose hairs. *Leaves* simple, alternate, crowded at the top of branchlets, 10–20 by 5–17 cm, broadly elliptic, or subrounded, margin often undulate or dentate, apex acute. Leaves of saplings or of lower branches 3–5-lobed, larger and with heart-shaped base. *Petioles* 6–12 cm long, bearing 2 globose, pink glands at the apex. *Inflorescences* panicles, terminal, branches widely spreading with slender peduncle. *Male flowers*: pedicels 10–12 mm long. *Calyx* tube with 2 or 3 irregular teeth, 2.5–3.5 mm long, apex obtuse, stellate tomentose outside. *Petals* 5, free, 5–7 mm long, pale yellowish white, narrow elliptic, with a hair tuft at the base.




Fig. 23. *Aleurites moluccana* (L.) Willd. (Euphorbiaceae).

Stamens 15–20, filament rolled in bud, anthers elliptic. *Female flowers*: pedicels bul-
late, 4–6 mm long. *Calyx* tube with 2 or 3 irregular teeth, 4–5 mm long, apex obtuse,
stellate tomentose outside. *Petals* 5, free, 8–10 mm long, pale yellowish white, oblong.
Ovary ovoid, stellate tomentose, 2-locular, 2 ovules per locule, styles 2, deeply bilobed.
Fruit globose, drupe, greenish yellow, 4–6 cm diam., with rufous stellate hairs. *Seeds*
2 in each locule, ovate, seed coat black, wrinkled, very hard, rich oil in endosperm.

Distribution — Originally from Malesia, but now also found in China, Thailand,
India. In Laos: Vientiane province. Planted or naturally growing in many provinces
from North to South Vietnam: in Cao Bang, Lang Son, Bac Thai, Hà Bac, Vinh Phu,
Quang Ninh, Hoa Binh, Thanh Hoa, Nghệ An, Đông Nai, and Tien Gian.

Habitat & Ecology — Rain forest, monsoon forest or tall advanced secondary forest,
on mountain slopes, very abundant on ridges, sometimes in sandy soil among granite
boulders, loam soil or on former shifting cultivation lands, altitude below 300 m. Light
demanding and fast-growing tree, good regeneration by seed. Either flowering in June,
July and then fruiting in August, September or flowering from February to March and
then fruiting from December to February.

Uses — Wood white and soft with fine grain, susceptible to termite attack, used for
general furniture. This species is cultivated for the oil obtained from its seed, which
is liquid, fast drying and used in paint-mixing, soap production and as lamp oil. The
fruit is edible, and the seed is mixed with curry in cooking and used as spice.

Specimens examined: *Balansa* 3287 (P); *Bon* 604, 2587, 2922 (P); *Chevalier* 29755 (P); *Clemens*
3858 (P); *Eberhardt* 3270, 4295, 5063 (P); *Fleury* 37903 (P); *Forestier* 14 (P); *King* 258 (L); *Mocret*
375 (P); *Pierre* 5291 (P); *Thorel* 671 (P); *Tsang* 27417 (P).

2. BACCAUREA Lour.

Baccaurea ramiflora Lour.

Synonyms — *Baccaurea cauliflora* Lour.; *B. oxycarpa* Gagnep.; *B. sapida* (Roxb.) Müll.Arg.;
Pierardia sapida Roxb.

Vernacular names — Dâu da đất, Du quả nhọn (V).

Trees up to 15 m high, up to 40 cm diameter. *Bark* orange-brown, inner bark often
reddish brown. *Branches* (sub)glabrous. *Leaves* simple, alternate, blade elliptic to
obovate 7–25.5 by 3–8.8 cm, papery, base attenuate to cuneate, margin entire, apex
cuspidate, upper surface glabrous except for midrib, lower surface glabrous, veins 4–9
per side. *Petioles* 1–5 cm long, kneed at both ends. *Stipules* 2.5–6 mm, pubescent,
caducous. *Inflorescences* axillary to cauliflorous, racemes, densely covered with red-
fulvous pubescence, bracts triangular, 1.5–4 mm long. *Male inflorescences* up to 15.5
cm long, usually 3 flowers per bract. *Male flowers* 1.1–4 mm diam., yellow; pedicel
0.8–2.6 mm long; sepals 4 or 5, different in shape, 1–2.4 mm long, stamens 5–9, free,
filaments 0.5–0.9 mm long, straight. *Female inflorescences* just below the leaves or
really cauliflorous, up to 16 cm long, 1–3 flowers per bract. *Female flowers* 3–8 mm
diam., yellow; pedicel 1.5–3 mm long; sepals 4 or 5, elliptic, 3.5–5 mm long; ovary
(2-) or 3- or (4-)locular; ovules 2 per locule, stigmas sessile, not lobed. *Fruits* globose
to ovoid berries, 19–32 by 14–25 mm, red to orange to pink to purplish outside, cream
inside, (sub)glabrous outside. *Seeds* 9–15 by 9–11 mm; arillode white.

Distribution — India (Andaman and Nicobar Islands), Myanmar, China (Yunnan, Hainan), Indochina, Thailand, Peninsular Malaysia. In Laos: Louang Prabang, Vientiane, Savannakhet, and Xieng Khouang provinces. In Vietnam: almost all provinces.

Habitat & Ecology — Primary or secondary rain forest, altitude below 1000 m, often cultivated on soil, sand, granite, moist and well-drained, along stream banks, in valleys, or on hill sides. Usually mixed with *Deutzianthus tonkinensis*, *Dracontomelum dao*, *Phoebe* spp., and *Pometia pinnata*. Flowering: March, April; fruiting: August, September.

Uses — Wood used for making furniture and agricultural implements. Arillode edible, sweet to sour.

Specimens examined: *D'Alleizette* 6431 (L); *Balansa* 3275 (L, P), 3276, 4757 (P); *Chevalier* 31902, 37416 (P); *Clemens* 4354 (P); *Contest* 300 (P); *Dong* 150 (P); *Eberhardt* 2684, 2702, 2767, 3043, 3917, 4580 (P); *Fleury* 30118, 32043, 37435 (P); *Kerr* 2493, 21280 (L); *Nidegene* 2081 (P); *Pételot* 4689, 4699 (P); *Pierre* 53, 145, 186, 395, 6276 (P); *Poilane* 1109, 1271, 10523, 11175, 11696, 12275, 20496 (L, P), 11964, 12041, 13661, 20094 (P); *Soejarto et al.* 10269 (L), 10567, 10659 (P); *Thorel* 770, *s.n.* (P); *Tsang* 29136 (L, P); *Vidal* 1428, 2009 (P);

3. BISCHOFIA Blume

Bischofia javanica Blume — Fig. 24

Synonym — *Bischofia trifoliata* Hook.

Vernacular names — Khom fat (Louang Prabang, L), Foug fat (Xieng Khouang, L); Nhội (V).

Trees up to 40 m high, up to 1.5 m diameter. *Trunk* straight, stout, and cylindrical. *Bark* pale brown or reddish brown, smooth, becoming darker and scaly with ages, inner bark brownish pink with red sap. *Stipules* papery, caducous. *Leaves* trifoliolate, alternate. *Petioles* 7–15 cm long; petiolules 0.5–1.2 cm long, end one 2–4 cm long; leaflets elliptic or ovate, 8–15 by 5–9 cm, cuneate at base, margin dentate, apex cuspidate, mature leaves leathery, dark green, smooth, old leaves red. *Inflorescences* panicles, axillary, dioecious, greenish yellow. *Male inflorescences* 7–20 cm long. *Bracts* lanceolate, 1–4 mm long, falling early. *Pedicel* 2–5 mm long. *Sepals* 5. *Petals* absent. *Stamens* 5, free, opposite sepals, filament short. *Disc* absent. *Female inflorescences* looser and less branched than the male, 15–30 cm long. *Pedicel* 4–6 mm long. *Sepals* 5, slightly fused at base. *Petals* absent. *Ovary* smooth, 3-locular, 2 ovules per locule, style very short with 3 long and spreading to recurved stigmas. *Fruits* globose, 0.5–1 cm diam., hanging in large clusters. *Seed(s)* 1 or 2 per locule, oblong, shiny, 4–5 mm long.

Distribution — India, South China, Thailand, Indochina, and throughout the Malasian areas (but rare in Peninsular Malaysia and large parts of Borneo). In Laos: Louang Prabang and Xieng Khouang provinces. In Vietnam: almost all provinces.

Habitat & Ecology — Dense primary or secondary evergreen forest, altitude below 1000 m. Growing on a wide range of soil type, such as well-drained clay soil, old basalt or alluvium soil of ravines and river bank. Natural regeneration is good. During the first year, scattered light is suitable for seedlings and young trees. Flowering: March, April; fruiting: May to December.

Uses — Timber heavy, sapwood brownish yellow, distinct from the reddish brown heartwood. Used for general construction, making furniture, doors, bridges, railway




Fig. 24. *Bischofia javanica* Blume (Euphorbiaceae).

sleepers, mining props, and agricultural implements. The bark contains tannin, used for toughening nets and ropes. This species is very successfully planted along the roads and parks of many provinces in Vietnam.

Specimens examined: *Balansa* 1501, 3966, 3967, 3968 (P); *Bon* 358, 431, 450, 875, 1424 (P); *Chevalier* 29759, 32191 (P); *Fleury* 30198, 37991 (P); *Harmand* 1885 (P); *Hiep et al.* 138 (P); *Hohenacker* 1573 (L); *Jaumatre* 2714 (P); *Master s.n.* (P); *Pierre s.n.* (P); *Poilane* 1229, 1596, 2159, 2343, 4834, 6683, 10735, 11265, 91265 (P); *Soejarto et al.* 10350b (P); *Thomson s.n.* (P); *Tsang* 27117 (P), 29206, 29933, 30319 (L, P); *Vidal* 775b, 1498 (P).

4. ENDOSPERMUM Benth.

Endospermum chinense Benth.

Vernacular name — Vạng trứng (V).

Trees up to 35 m high, up to 1 m diameter. *Trunk* straight, cylindrical. *Bark* yellowish grey with transversal rings around the trunk, inner bark white with watery sap. *Twigs* and *petioles* stellately tomentose. *Stipules* small, triangular, hairy. *Leaves* simple, alternate, young leaves cordate, 10–25 cm long, old leaves smaller, ovoid or nearly round, 8–15 by 5–13 cm, base of lower surface with 2 crab-eye-shaped glands, margin entire, apex rounded, upper surface green, lower surface yellowish green, venation palmate, 3–5 secondary veins per side. *Petioles* 3–10 cm long, hairy. *Flowers* unisexual, dioecious. *Male inflorescences* panicles, up to 21 cm long, with branches up to 4 cm long. *Calyx* 4-lobed, pubescent outside, glabrous inside. *Petals* absent. *Stamens* 10, spirally arranged on an androphore, filaments free, anthers 4-celled. *Female inflorescences* racemose, up to 18 cm long. *Sepals* 4 or 5. *Petals* absent. *Ovary* globose, 2- or 3-locular, 1 ovule per locule, style very short, stigmas 2- or 3-lobed. *Fruit* a globose drupe, 1–1.5 cm diam., stigmas persistent.

Distribution — India, Myanmar, China (Hainan), Hong Kong, Vietnam, Thailand. In Vietnam: Bac Thai, Yên Bái, Vinh Phú, Hoa Bình, Quang Ninh, Hà Tây, Ninh Bình, Thanh Hoa, and Quang Tri provinces. In Laos: not yet recorded.

Habitat & Ecology — Primary and secondary forest, often along streams and on water logged or on dry sandy soil, altitude 50–500 m. Usually mixed with *Canarium* spp. and *Cryptocaria lenticellata*. A light-demanding and fast-growing tree, deciduous during winter time. Natural regeneration is good under 50–60% forest cover. Flowering: May, June; fruiting: November, December.

Uses — Wood yellow-white, soft and light. Used for making matches, pencils, table tennis rackets, plywood, and general furniture.

Specimens examined: *D'Alleizette* 6531, s.n. (L); *Chevalier* 32260 (P); *Fleury* 30137 (P); *Forestier* s.n. (P); *Poilane* 141, 1296 (P).

FAGACEAE

Usually medium or big trees, rarely shrubs. Buttresses or stilt roots sometimes present. *Crown* deep, dense, domed, often with massive, more or less spreading branches, often grey or silvery from below. *Bark* usually grey, smooth with prominent lenticels often in lines, scaly or fissured, inner bark usually cream and brown, with much tannin, darkening on exposure, hard, firm, coarsely fibrous textured. Cambium often ridged and penetrated by hardwood rays. Indumentum consisting of simple, stellate or fascicled hairs. *Stipules* present, caducous or rather long persistent. *Leaves* simple, alternate, spirally arranged, distichous (a few species of *Castanopsis*), or crowded near the top of

the twigs (species of *Quercus*), margin entire, denticulate, serrulate, crenate or lobed, venation pinnate. *Inflorescences* unbranched or branched, male and female separated or mixed. *Male inflorescences* erect or pendulous in the axils. *Male flowers* solitary or in dichasial clusters of 3–7(–30) along the rhachis, sessile. *Sepals* 4–6(–9), scale-like, connate or distinct. *Stamens* (4–)6–12(–15), filaments filiform, anthers reniform or ovoid, longitudinally dehiscent. *Female inflorescences* erect in the axils. *Female flowers* sessile, solitary or in dichasial clusters of 3–15 along the rhachis. *Sepals* 6, scale-like. *Petals* absent. *Staminodes* 6–12 or absent. *Ovary* inferior, 3–6(–9)-locular, 2 ovules per locule, styles as many as ovary locules, stigmas capitate (*Quercus* and *Trigonobalanus*) or punctiform (*Castanopsis* and *Lithocarpus*). *Cupules* solitary or in dichasial clusters along the rhachis, saucer- to cup-shaped or completely enclosing the fruit, at maturity indehiscent or splitting into 2–8 (or more) segments, external wall muriculate, scaly, spiny, with concentric, spiral lamellae or rarely almost smooth. *Fruits* a 1-seeded nut, 1–4 nuts grouped within a cup (cupule). *Seeds* without endosperm, embryo and cotyledons large.

Distribution — In temperate and tropical regions but mostly in the northern hemisphere: c. 8 genera, more than 900 species. In Laos and Vietnam: c. 8 genera, c. 211 species.

KEY TO THE GENERA

- 1a. Deciduous trees. Leaf margin dentate. Stipules up to 1 cm long. Ovary 6–9-loculed. Capsules with 2 or 3 nuts, entirely covered with hard spines **1. *Castanea***
- b. Evergreen trees. Leaf margin entire or dentate. Stipules smaller than 4 mm long. Ovary 3–6-loculed. Capsules with 1 nut, entirely or partly covered with spines **2. *Castanopsis***

1. *CASTANEA* Tourn. ex L.

***Castanea mollissima* Blume**

Synonyms — *Castanea duclouxii* Dode; *C. fargesii* Dode; *C. hupehensis* Dode

Vernacular names — Dẻ trứng khảnh, Dẻ tàu, Dẻ trổng (V).

Trees up to 25 m high, up to 1 m diameter. *Trunk* much branched, branches brown fulvous, with white scales. *Twigs* grey, tomentose, slender and densely arranged when old. *Bark* rough. *Leaf buds* small, ovate, tomentose. *Stipules* narrow ovate, c. 1 mm long, rather coriaceous persistent or caducous. *Leaves* simple, alternate, distichous, ovate or oval, rough, dry, thick, 14–20 by 5–7 cm, base rounded, truncate or obliquely cordate, margin sparsely and irregularly serrate, apex acute to acuminate, sparsely white tomentose beneath when young, lateral veins 12–16 pairs, obviously evident beneath. *Petioles* glabrous or slightly tomentose, 0.7–2 cm long. *Male inflorescence* 8–20 cm long. *Perianth* scale-like. *Stamens* usually 10–12, slender. *Female flowers* densely arranged. *Styles* 7–9 mm long, erect or spreading, much tomentose. *Cupule* globose, 4.5–6 cm diam., entirely covered with acute and hard spines, over 1 cm long and stellately hairy, 2 or 3 nuts in each cupule. *Seeds* 2–3 cm diam., reddish brown, glabrous at the base, hairy at the top.

Distribution — China and Vietnam. In Vietnam: Lào Cai, Cao Bang (Trung Khach), Lang Son, Lai Châu and many other provinces. In Laos: not yet recorded, but likely present.

Habitat & Ecology — Altitude 500–2000 m. Light demanding, growing on waste land, stony, acid, or limestone soils. Natural regeneration is good. It can be planted under secondary forest cover. Flowering: February, March; fruiting: October to December.

Uses — Wood rather valuable, brownish with straight veins, gross and heavy. Used for construction, boat building, making agricultural tools and pit props. Seeds edible, contains much starch.

Specimens examined: *Farges 820* (P); *Kiang 1021* (P); *Rock 4754, 6436* (P).

2. CASTANOPSIS Spach

KEY TO THE SPECIES

- 1a. Leaves obovate, petiole 1–2 cm long. Cupule enclosing about half of the nut, scales arranged into many irregular bands **1. *C. cerebrina***
 b. Leaves elliptic, petiole 0.4–1.3 cm long. Cupule enclosing the nut totally with long and hard spines **2. *C. indica***

1. *Castanopsis cerebrina* (Hickel & A. Camus) Barnett

Synonym — *Pasania cerebrina* Hickel & A. Camus

Vernacular names — Dẻ bốp, Sỏi phẳng, Sỏi đen (V).

Trees up to 25 m high, up to 90 cm diameter. *Trunk* straight, young branches covered with dense reddish brown, simple and stellate hairs. *Leaves* distichous, obovate, 12–20 by 4–8 cm, base cuneate, margin acutely and regularly serrate for 3/4 of the length towards the top, apex acuminate, green above, reddish brown tomentose beneath, if mature almost glabrous, venation pinnate, lateral veins 10–20 pairs, curved at the margin, prominent on lower surface. *Petiole* 1–2 cm long, slightly tomentose. *Male catkins* 4–9 cm long, in branched panicles, usually terminal. *Perianth* puberulous. *Anthers* small, filament slender. *Pistillode* small, styles 3, divergent. *Female inflorescences* 5–11 cm long. *Cupule* cup-shaped, 1.8–2.5 cm long, 1.6–2.2 cm diam., enclosing about 1/2 the seed, densely hairy inside, slightly tomentose outside, scales acute, flat, arranged into many irregular bands, pedicel 4–5 mm long, solitary. *Nut* narrowly conical, 2–2.5 by 1.4–1.8 cm, tomentose, apex sometimes 3-angled.

Distribution — China (Southeast Yunnan), Indochina, North Thailand. In Laos: Louang Namtha province. In Vietnam: Hà Tĩnh, Nghệ An, Thanh Hoa, Yên Bái, Tuyên Quang, Quang Ninh, Hà Bắc, and Hà Tây provinces.

Habitat & Ecology — On low hills or on the forest edge, altitude below 400 m. Light demanding tree. Natural regeneration is good, germination rate very high, also in the nursery. Flowering: February to April, October, November; fruiting: March to June, October to December.

Uses — Wood hard, can be used for construction and making agricultural tools.

Specimens examined: *Balansa 2380* (P); *Fleury 30117* (P); *Forestier s.n.* (P); *Pételot 1551, 4831* (P); *Poilane 26264, 26315* (P); *Toan 32A* (L).

2. *Castanopsis indica* (Roxb. ex Lindl.) A.DC.

Synonyms — *Castanea indica* Roxb. ex Lindl.; *Castanopsis sinensis* A. Chev.; *C. subacuminata* Hayata

Vernacular names — Ko ket, Ko lakai, Ko luay, Ko tang (L); Dê gai Ân Đô, Cà ôi (V).

Trees up to 30 m high, up to 1 m diameter. Bole straight. *Crown* broadly rounded. *Bark* brownish grey with dark grey traces, wrinkled, thick, grooved, inner bark much fibrous. Bud scales, young branchlets, petiole, leaf blades abaxially and rhachis of inflorescences densely covered with reddish brown hairs. *Leaves* elliptic 9–20 by 3–9 cm, base obtuse, margin acutely and regularly serrate for 3/4 of the length towards the top, apex acute or acuminate, green above, reddish brown tomentose beneath, venation pinnate, lateral veins 14–20 pairs, curved at the margin, prominent on lower surface, tertiary veins scalariform, young leaves densely pubescent above and on nerves beneath. *Petiole* 0.4–1.2 cm long, fulvous hairy. *Male inflorescences* in branched spikes, 10–18 cm long. *Flowers* with aromatic smell, nearly rounded. *Perianth* of 6 segments, scale-like. *Stamens* 8–14, filament slender. *Female inflorescences* in spikes, 12–20 cm long. *Flowers* often solitary on the rhachis, longer than the leaf, bearing abortive stamens. *Ovary* inferior, 3-locular, styles 3, long, spreading. *Cupule* enclosed nut totally, 1.6–2.2 cm diam., covered with spines. *Nut* 1 per locule, broadly conical, 0.6–1.4 cm diam., shiny brown, dehiscent when mature.

Distribution — India, Nepal, Bangladesh, Myanmar, China, Indochina, Thailand. In Laos: Phong Sali, Louang Prabang, and Xieng Khouang provinces. In Vietnam: in almost all provinces in the north, mainly in Vinh Phu, Hà Bac, and Lang Son provinces.

Habitat & Ecology — Primary and secondary forests, but only occurring scattered, not forming clearly dominant stands. Usually mixed with *Engelhardtia* spp., *Ixonanthes cochinchinensis*, *Lithocarpus* spp., *Madhuca pasquieri*, and *Quercus* spp. Light demanding tree, suitable on sandy soils with medium humidity. Natural regeneration is good, coppice regeneration is possible. Flowering: January to March, October to December; fruiting: January, February, June to August.

Uses — Wood is of good quality, pinkish brown, hard, and resistant to termites and insects. Used for construction and making furniture. The bark is rich in tannin and the fruits are edible.

Specimens examined: *D'Alleizette* s.n. (L); *Balansa* 2388 (L); *Blauche* 4382 (P); *Bon* 5048 (P); *Brune* 3865 (P); *Chevalier* 1669, 1732, 35719 (P); *Cuong* 363 (L); *Dussau* 79 (P); *Eberhardt* s.n. (L); *Fleury* 30100 (P); *Hiep* 4229 (L); *Pételot* 4740, 4741 (P); *Poilane* 1913, 2194, 2394, 2425, 2621, 4805, 5311, 10554, 10687, 10815, 11071, 12127, 12153, 12197, 12641, 13427, 16244, 17954, 18880, 19027, 20008, 20124, 22829, 22831, 24203, 24727, 24840, 24876, 24888, 24912, 24914, 25075, 27154, 30063 (P), 13027, 20298, 20504 (L), 19926, 24283, 24875, 27124 (L, P); *Soejarto* 10395, 11034 (L); *Toan* 32 (L); *Tsang* 27132, 27408, 27459, 30733, 30754 (P); *Vidal* 5697, 16966, 74413 (P).

IRVINGIACEAE

Evergreen trees with conspicuous annular leaf scars on young branches. *Stipules* sword-like, enclosing the terminal buds, caducous. *Petiole* grooved above. *Leaves* simple, coriaceous or subcoriaceous, glabrous or pubescent, margin entire. *Inflorescences* axillary or terminal panicles. *Bracts* small, early caducous. *Flowers* 5-merous, bisexual.

Sepals connate at base, imbricate in bud. *Petals* exceeding sepals, imbricate in bud. *Stamens* 10, inserted beneath the disc. *Ovary* superior, 2-locular, conical or somewhat flattened, sessile on the torus-like disc, stigma inconspicuous, ovules solitary. *Fruit* a large 1-seeded drupe.

Distribution — Worldwide: 1 genus, 8 species. In Laos and Vietnam: 1 species. Often included under Simaroubaceae.

IRVINGIA Hook.f.

Irvingia malayana Oliv. ex A. Benn. — Fig. 25

Synonyms — *Irvingia harmandiana* Pierre ex Laness.; *I. oliveri* Pierre

Vernacular names — Mai bok, Bok (L); Konia (V).

Evergreen trees up to 35 m high with dense spreading crown and massive, buttressed trunk at base, up to 200 cm diameter. *Bark* pale grey-brown, smooth becoming irregularly cracked and flaky when old, inner bark pale orange. *Branchlets* with conspicuous



Fig. 25. *Irvingia malayana* Oliv. ex A. Benn. (Irvingiaceae).

annular leaf-scars. *Stipules* sword-like, 1.5–3 cm long, enclosing the terminal buds, early caducous. *Petiole* 1–2 cm long, glabrous, grooved above. *Leaves* elliptic, ovate, 7–20 by 2.5–9 cm, base acute, obtuse, slightly oblique, margin entire, apex acute, mature leaves completely glabrous, dark green and shiny above, usually pale grey-green and glabrescent below, secondary veins 8–10 pairs, venation reticulate, conspicuous on both surfaces. *Inflorescences* axillary, or terminal panicles or raceme. *Bracts* ovate with acute apex. *Flowers* small, 6 mm long, greenish white or yellowish. *Petals* free, about 3 times as long as sepals. *Stamens* 10, free, attached outside and below the central disc. *Ovary* superior, 2-locular, style 1, stigma inconspicuous. *Fruit* ovoid or ellipsoid, 6 by 4 cm, fleshy, green when young and yellow when mature, pendulous on long stalk. *Seed* 1 with white, fatty cotyledons.

Distribution — India, Myanmar, Indochina, Peninsular Malaysia, Indonesia (Sumatra, Borneo). In Laos: All tropical parts of the country. In Vietnam: Quang Nam-Dà Nang and Lâm Đông provinces.

Habitat & Ecology — Light-demanding tree when young, frequent in dry deciduous Dipterocarp forest, dry evergreen forest, tropical rain forest, altitude below 300 m. In Laos: flowering: January to March; fruiting: February to April. In Vietnam: flowering: May, June; fruiting: September to November.

Uses — Used for general construction, excellent as charcoal. Ripe fruits sweet and edible. Seeds contain white or yellow oil, used for soap making and lighting, and are edible either cooked or raw.

Note — The genus *Irvingia* is treated as a member of Simaroubaceae in Nootboom, Flora of Malesiana, Series 1, Vol. 6 (1962) 193.

Specimens examined: *Bon* 3288 (P); *Evrard* 2302 (P); *Harmand* 653 (P); *Hiep* 666 (P); *Nootboom* 71286 (L).

LAURACEAE

Trees or shrubs, rarely parasitic leafless climbers (*Cassytha*); trees usually without buttresses. *Bark* usually smooth, rarely fissured or scaly, often covered with large lenticels, inner bark usually thick, granular, mottled or laminated, often with strong aromatic smell, yellow, orange-brown, pinkish or reddish. Terminal bud naked or covered with bud scales, sometimes resembling small leaves (*Actinodaphne*). *Stipules* absent. *Leaves* simple, spirally arranged, alternate, rarely opposite, subopposite or whorled, often glaucous below, usually with numerous minute oil dots, often aromatic when crushed, secondary veins pinnate, sometimes tripliveined (*Cinnamomum*). *Inflorescences* in racemes or panicles or in clusters arising from the trunk or branches, terminal or axillary. *Flowers* small, regular, greenish white or yellow, fragrant or with rancid smell, bisexual or unisexual, perianth with 3 or 6 lobes in 2 whorls. *Stamens* 9–12 alternate, usually attached to the throat of the tube, usually 3 whorls, rarely more (*Cinnadenia*), anthers 2- or 4-celled, cells opening by valves, usually from base to apex, outer two whorls of anthers as a rule introrse, 3rd whorl extrorse, in *Litsea* all whorls extrorse, ovary superior. *Fruit* small to large, berry or drupe, usually with persistent perianth at base, except *Litsea*. *Seed* 1 with thin testa, cotyledons large, flat.

Distribution — Throughout the world: c. 35 genera, 2500 species. In Laos and Vietnam: c. 19 genera, 245 species.

CINNAMOMUM Schaeff.

KEY TO THE SPECIES

- 1a. Bark smooth or lenticellate. Leaves opposite or subopposite, tripliveined at base 2
 b. Bark fissured. Leaves alternate, venation pinnate 3
 2a. Twigs quadrangular, dark green. Leaves leathery, very fragrant when crushed **1. C. aromaticum**
 b. Twigs terete, pale green. Leaves papery, only slightly fragrant when crushed **4. C. iners**
 3a. Leaves tripliveined, lateral veins 2 or 3 pairs, domatia present in the axils of secondary veins and midrib. Petiole 2–5 cm long. Perianth cup 0.4–0.6 cm high **2. C. camphora**
 b. Leaves with 4 or 5 pairs of lateral veins, domatia absent. Petiole 2–3 cm long. Perianth cup 1–1.6 cm high **3. C. glaucescens**

1. Cinnamomum aromaticum Nees in Wall.

Synonym — *Cinnamomum cassia* Blume

Vernacular names — Sa chuang, Say chuang (L); Quế (V).

Medium trees up to 20 m high, up to 70 cm diameter. *Trunk* straight, cylindrical. *Bark* greyish brown, smooth or lenticellate. All parts of the tree with aromatic smell. *Young twigs* quadrangular, green, densely blackish brown tomentose, indumenta not persistent. *Petioles* 1.2–2.5 cm long. *Leaves* subopposite, elliptic, 10–24 by 4–7 cm, leathery, shining green, very fragrant when crushed, base cuneate, apex blunt or pointed, tripliveined, veins raised beneath. *Inflorescences* in terminal and axillary panicles, up to 16 cm long, hairy. *Flowers* white, perianth densely silky hairy, lobes 6, in 2 whorls. *Stamens* 9, in 3 whorls. *Ovary* ovoid, glabrous, style glabrous. *Fruit* 1–1.3 by 0.7–0.9 cm, seated on the lobed perianth cup, cup 0.5–0.8 cm high.

Distribution — India, China, Taiwan, Indochina, Indonesia. In Laos: Xieng Khouang and Vientiane provinces. In Vietnam: Quang Ninh, Yên Bái, Lào Cai, Thanh Hoa, Nghệ An, and Quang Nam provinces. Nowadays this species is widely cultivated in many provinces, such as Son La, Lang Son, Hà Tây, Vinh Phu, Hoa Binh, Hà Tĩnh, and Khanh Hoa.

Habitat & Ecology — Naturally distributed in primary and secondary forest, altitude below 800 m. Light-demanding tree, slightly shade-tolerant when young, preferring deep and acid soil. Flowering: May to August; fruiting: October to December.

Uses — Wood used for making furniture, sawing boards, implements. Essential oil, extracted from all parts of the tree, used in industry and medicine, for digestive complaints such as flatulence, colic, dyspepsia, diarrhoea, and nausea. It can also be used for colds, influenza, fevers, diabetes, athlete's foot, arthritis, and rheumatism.

Specimens examined: Chevalier 35977 (P), 37713 (L, P), 37782, 38575, 38782 (P); Fean 37717 (P); Fleury 37918 (L, P); Forestier 35976, 38555 (P); Hiep et al. 1329 (P); Pételot 6942 (P); Poilane 3557, 7055 (P), 31713, 31761 (L, P), 31722, 32367 (P); Quesnel 37711 (P).

2. *Cinnamomum camphora* (L.) T. Nees & C.H. Eberm.

Synonym — *Laurus camphora* L.

Vernacular names — Chuang (L); Long nảo, Re (V).

Trees up to 30 m high, up to 80 cm or more diameter. *Trunk* usually short, stout. *Bark* greyish brown, deeply fissured, inner bark with aromatic smell. *Twigs* shining, green. *Buds* ovoid, pubescent. *Leaves* alternate, broadly ovate to elliptic, 4–9 by 2.5–5 cm, base oblique or obtuse, apex acuminate, glabrous above, glaucous beneath, 3 main veins, 2 or 3 pairs of lateral veins, raised beneath, with small domatia, fresh leaves aromatic when crushed. *Petioles* 2–5 cm long. *Inflorescences* in terminal and axillary panicles, up to 8 cm long, whitish hairy. *Flowers* white. *Perianth* densely silky hairy, lobes 6. *Stamens* 9, in 3 whorls, 1st and 2nd whorl of stamens eglandular, 3rd whorl with 2 ovate subsessile glands at the base; staminodes 3. *Ovary* ovoid, glabrous, style glabrous. *Fruit* globose, 0.6–1 by 0.5–0.7 cm, green when young and blackish violet when mature, seated on a lobed 0.4–0.6 cm high perianth cup.

Distribution — Originated in China. Planted in India, Indochina, Thailand, Malaysia, Indonesia. In Laos and Vietnam: planted in almost all provinces.

Habitat & Ecology — Naturally in primary and secondary forest, altitude below 1500 m. Light demanding tree, slightly shade tolerant when young, preferring deep, moist and humus rich soil. Tolerates cold well, but poorly resistant to frost. Flowering: February to April; fruiting: September to December.

Uses — Timber greyish or pinkish grey, relatively hard and heavy, resistant to insects and termites, aromatic. Used for construction and making furniture. Essential oil, extracted from all parts of the tree, used in industry and medicine, to treat parasitic infections, diarrhoea, inflammation, itchiness, nervous-related ailments, chest and muscle pains, and toothaches. This species is very successfully planted along the roads and parks of many provinces in Vietnam.

Specimens examined: *Bon* 1826, 5688 (P); *Chevalier* 7, 29753, 29762, 29763, 29955, 36594 (P); *Dufort* 18 (P); *Forestier s.n.* (P); *Pierre* 5160 (P).

3. *Cinnamomum glaucescens* (Nees) Hand.-Mazz.

Synonym — *Cinnamomum balansae* Lecomte

Vernacular names — Chuang (L); Vũ hương, Gù hương (V).

Trees up to 45 m high, up to 80 cm or more diameter. *Trunk* straight. *Branchlets* glabrous, turning pale black. *Bud* densely scaled. *Bark* brown, longitudinally fissured, inner bark with aromatic smell, 1–2 cm thick. *Leaves* alternate, broad ovate to elliptic, subcoriaceous, 5–14 by 3–6 cm, base obtuse or acuminate, glabrous, apex acuminate, venation pinnate, 4 or 5 pairs of veins, raised beneath, midrib and secondary veins tomentose above. *Petioles* 1–3 cm long. *Inflorescences* in axillary panicles, up to 10 cm long, hairy. *Flowers* bisexual, white. *Perianth*, lobes 6, hairy outside. *Stamens* 9, in 3 whorls; staminodes 3, triangular. *Ovary* ovoid, glabrous, style short, stigma disciform. *Fruit* globose, 0.8–1.5 by 0.6–1 cm, seated on a lobed perianth cup, cup 1–1.6 cm high.

Distribution — China, Vietnam, Laos, Thailand. In Laos: Attopeu province. In Vietnam: Vinh Phu, Hoa Binh, Ninh Binh, Thanh Hoa, Nghệ An, Quang Tri, Đông Nai, and Binh Dinh provinces.

Habitat & Ecology — Primary forest, altitude below 700 m. Light-demanding tree, slightly shade tolerant when young, preferring deep, moist and well-drained soil. Usually mixed with *Dracontomelum dao*, *Parashorea chinensis*, and *Pometia pinnata*. Natural regeneration is very poor. Flowering: February to April; fruiting: September to November.

Uses — Wood heavy, resistant to insects and termites, aromatic, easy to work. Used for construction, making high quality furniture, and sawing boards. Essential oil, extracted from all parts of the tree, used in medicine, soap and lubricant producing.

Specimens examined: *Fleury* 30173, 37993 (L); *Hennipman* 3626 (L); *Poilane* 1073, 15939, 18790, 18868, 19940 (P), 1108, 1272, 1368, 24505 (L, P), 19848 (L).

4. *Cinnamomum iners* Reinw. ex Blume

Synonyms — *Cinnamomum nitidum* Hook.; *C. paraneuron* Miq.

Vernacular names — Chuang, Si khai ton (L); Re hương, Re xanh (V).

Medium sized trees up to 24 m high, up to 60 cm diameter. *Bark* greyish brown, smooth or lenticellate, inner bark pinkish, with strong fragrant smell. Sapwood whitish. *Twigs* slender, pale green, slightly pubescent, then glabrescent. *Leaves* opposite or subopposite, young leaves pink, elliptic, 8–23 by 3–8 cm, base cuneate, margin entire, apex blunt or pointed, venation tripliveined, lower surface often glaucous, midrib and nerves raised beneath. *Petioles* 1–1.5 cm long. *Inflorescences* in terminal and axillary panicles, up to 20 cm long, hairy. *Flowers* creamy, with unpleasant smell. *Perianth* densely silky hairy, lobes 6. *Stamens* 9, in 3 whorls; staminodes 3, triangular. *Ovary* ovoid, glabrous, style glabrous. *Fruit* globose, 1–1.5 by 0.8–1 cm, seated on a lobed perianth cup, cup small, 0.4–0.6 cm high, perianth lobes persistent in fruit.

Distribution — India, Myanmar, Indochina, Thailand, Malaysia, Indonesia, Philippines. In Laos: Vientiane province. In Vietnam: Hoa Binh, Quang Ninh, Vinh Phu, Thanh Hoa, Ninh Binh, Đông Nai, Kom Tum, and Sai Gon provinces.

Habitat & Ecology — Common in primary and secondary forest, often in moist forest, altitude up to 1200 m. Natural regeneration is good under a forest cover of 20–40%. Flowering: February to April; fruiting: July, August.

Uses — Wood used for construction and making furniture. Bark aromatic, often used for incense production, or together with leaves in tea, and also as a substitute for Cinnamon. The oil from the leaves is used for flavouring sweets and confectionery. This species is sometimes planted as a shade tree.

Specimens examined: *Chevalier* 35933, 35956, 36680, 36681 (P); *Cuong* 4504 (P); *Fleury* 29918 (P); *Harmand* 185 (P); *Poilane* 7077, 29392 (P); *Thorel* 185 (P); *Tixier s.n.* (P); *Vidal* 4026 (L, P), 5060 (L).

LOGANIACEAE

Trees, shrubs, climbers, or herbs. Hairs simple. *Stipules* present or absent. *Leaves* simple, entire, opposite or whorled, connate at the base by a transverse ridge. Leaf blade ovate, base cuneate, apex acuminate, margin entire, secondary veins pinnate, rarely

tri-pinnate or -palmate. *Inflorescences* terminal or axillary, solitary or in cymes. *Flowers* bisexual, 5- to many, rarely 4-merous, actinomorphic. *Calyx* united or free. *Corolla* united, twisted in bud. *Stamens* inserted on the corolla tube or in the throat, anthers attached at the base. *Ovary* superior, usually with 2 locules, rarely 1 or 4, placentas axile or parietal, locules with 1 to many ovules, style single, head not swollen. *Fruits* berries, drupes, or capsules. *Seeds* usually numerous, often irregular.

Distribution — Worldwide: c. 20 genera, 400 species. In Laos: 4 genera, c. 12 species. In Vietnam: 5 genera, c. 26 species.

FAGRAEA Thunb.

Fagraea fragrans Roxb.

Synonyms — *Fagraea cochinchinensis* A. Chev.; *F. peregrine* Blume

Vernacular names — Man pa (L); Trai nam bô, Trai (V).

Evergreen trees up to 8–30 m high, up to 150 cm diam. with small buttress at the base. *Bark* yellow-grey or brown, deeply longitudinal fissured, yellowish white sap, twigs greyish brown, terete, glabrous, nodose. *Stipules* 1–2 mm long. *Petiole* 0.7–2.5 cm long, slender, terete, without auricles. *Leaves* elliptic, 7–12 by 1.5–5 cm, coriaceous, base acute or cuneate, margin entire, apex acute, acuminate or caudate, both surfaces glabrous, midrib slightly raised above, prominently keeled or ridged below, secondary veins 5–16 pairs, barely visible or invisible above, visible below. *Inflorescences* axillary or terminal corymbs, 4–13 cm long, 3 to many flowers, peduncle 1.5–7 cm long. *Flowers* very fragrant. *Pedicels* 0.3–2 cm long. *Sepals* 5, connate into a bell-shaped tube, 1.7–3 mm long, lobes 0.7–2 mm long, apex rounded, spreading in fruit. *Petals* 5, 12–20 mm long, creamy white turning dull-yellow, tube 6–12 mm long, narrowly funnel-shaped, lobes 5.5–8 mm long. *Stamens* 5, filaments 1.4–2.2 cm long, anthers elliptic, 1.5–2 mm long. *Ovary* glabrous, style longer than the stamens, 17–24 mm long, stigma 0.7–1 mm long, capitate or obconical, occasionally faintly bilobed. *Fruit* globose or subglobose, drupe, 4.5–10 mm diam., green turning orange or red, tip rounded or pointed. *Seeds* roughly angular, 0.5–2 mm long.

Distribution — India, Myanmar, Indochina, Thailand, Andaman Island, Malesia. In Laos: Savannakhet (Ban That), Vientiane Municipality, and Vientiane (pers. obs.) provinces. In Vietnam: Hà Bac, Quang Binh, Quang Tri, Quang Nam-Dà Nang, Dac Lac, Đông Nai, Lâm Đông, Kiên Giang, and Tây Ninh provinces.

Habitat & Ecology — Evergreen forest, mixed deciduous forest, on sandy soil, along roadsides, streams, or rivers, along paddy fields, altitude below 800 m. Often planted as an ornamental or for timber. Flowering: April to July; fruiting: July to November.

Uses — Timber used for house pillars, in construction, and furniture-making. Heartwood used as fuel. Decoctions of the bark are used as a febrifuge and decoctions from the leaves in the treatment of severe diarrhoea. Trees planted for shading.

Specimens examined: *D'Alleizette* 4904, 6909 (L); *Pierre* 1832 (L, P), 3697, 3698, 3699 (L); *Poilane* 16384 (L, P), 18549 (P); *Vidal* 1752 (P).

LYTHRACEAE

Trees, shrubs, or herbs. *Stipules* absent. *Leaves* opposite or whorled, simple, margin entire. *Inflorescences* terminal or axillary panicles. *Flowers* bisexual, regular. *Calyx* bell- or funnel-shaped, smooth or ridged, with 3–6 lobes. *Petals* free, 2–9, or absent, inserted near the mouth of the calyx tube. *Stamens* twice as many as petals or numerous, inserted on the calyx tube. *Ovary* with 1–6 chambers, half inferior, ovules numerous, style 1, stigma a small head. *Fruits* small or large capsules, seated on or surrounded by the persistent calyx. *Seeds* numerous.

Distribution — Throughout the world except the coldest parts: 25 genera, 600 species. In Laos: 1 genus, c. 10 species. In Vietnam: 5 genera, 23 species.

LAGERSTROEMIA L.

Lagerstroemia calyculata Kurz

Synonym — *Lagerstroemia angustifolia* Pierre ex Laness.

Vernacular names — Peuay dok khao, Peuay peuak na (L); Bằng lằng, Sang lằ (V).

Large deciduous trees 20–35 m high, 40–80 cm diameter. *Bark* pale creamy-grey, smooth, flaking in thin, rounded plates, inner bark yellowish, fibrous, 6–10 mm thick. *Twigs* slender, yellow, stellately hairy. *Stipules* absent. *Leaves* ovate or elliptic, 7–18 by 2–5 cm, base obtuse, unequal, apex acute, dark brown above and pale yellow beneath when dry, young leaves and twigs densely covered with yellowish star-shaped hairs which easily rub off, lateral veins 10–13 pairs, larger tertiary veins scalariform, smaller venation reticulate. *Petioles* 3–5 mm long, tomentose. *Inflorescences* in terminal panicles, densely yellow-tomentose, 12–20 cm long. *Bracts* and bracteoles early caducous. *Flowers* white, in groups of 6–8. *Calyx* bell-shaped, covered with many stellate hairs, tube 5 mm long, with 6 teeth, densely yellow-hairy outside and inside of teeth apex, without grooves or ridges, *Petals* 6, orbiculate. *Stamens* numerous, subequal, anther orbiculate. *Ovary* 5- or 6-locular, style simple or divided. *Fruit* a capsule, ovoid, 6–12 mm long, dark brown and glossy, dehiscent into 6 pieces, calyx persistent. *Seeds* c. 8 mm long.

Distribution — Indochina and Thailand. In Laos: Xieng Khouang, Savannakhet, Saravane, and Attopeu provinces. In Vietnam: Son La, Thanh Hoa, Nghê An, Binh Thuân, Tây Ninh, and Đông Nai provinces.

Habitat & Ecology — Deciduous forests, usually associated with *Anisoptera* spp., *Shorea* spp., *Terminalia* spp., or in Bamboo thickets. Natural and coppice-regeneration are good, although the trees are rather slow growing. Flowering: May to July; fruiting: August to April.

Uses — Sapwood and heartwood distinctive, sapwood white, heartwood greyish yellow or brown, wood hard and heavy. Used for timber, interior furniture-making, wood-boards.

Specimens examined: *Kerr* 71286 (L); *Maxwell* 89-775 (L); *Poilane* 3368, 6633, 10798, 15438, 16258, 21523 (P).

MAGNOLIACEAE

Trees or shrubs, trees usually with buttresses. *Bark* usually smooth, rarely fissured or scaly, inner bark usually thick, with a very spicy smell. *Leaves* simple, alternate, spirally arranged, commonly with very conspicuous reticulations on both surfaces. *Stipules* large, enclosing the young bud, soon falling and leaving ring-like scars on the twigs. *Flowers* solitary, bisexual, large, terminal or sometimes axillary. *Tepals* mostly 9 or more, free, imbricate, variously coloured. *Stamens* numerous, free, spirally arranged, filament short, anthers linear, 2-locular. *Carpels* numerous, free, spirally arranged on an elongated axis, ovules 2 or more in each carpel, ovaries superior. *Fruit* apocarpous, sometimes syncarpous, fruiting carpels mostly dry and dehiscent, rarely fleshy or indehiscent, each with 1 or more seeds, with red or creamy white fleshy aril, embryo small.

Distribution — Throughout the temperate and tropical regions: c. 12 genera, 200 species. In Laos and Vietnam: c. 7 genera, 39 species.

MICHELIA L.

KEY TO THE SPECIES

- 1a. Inner bark greyish yellow, with aromatic smell. Twigs pale black. Leaves glossy above, silvery ferruginous villose beneath. Petiole slightly swollen at base. Fruit with many follicles, 1–5 seeds in each follicle **1. M. balansae**
- b. Inner bark cream, bad smelling. Twigs pale green. Leaves dull above, pale green beneath. Petiole not swollen at base. Fruit with 3–7 separated follicles, 3–5 seeds in each follicle **2. M. mediocris**

1. *Michelia balansae* Dandy

Synonym — *Michelia baviensis* Finet & Gagnep.

Vernacular names — Giỏi, Giỏi bà, Giỏi lông (V).

Trees up to 20 m high, up to 60 cm diameter. *Trunk* straight. *Bark* greyish brown, glabrous, inner bark greyish yellow, with aromatic smell. *Twigs* pale black, ferruginous tomentose, later glabrous, lenticels sparse, white. *Leaves* simple, alternate, obovate, 10–20 by 4–10 cm, coriaceous, base cuneate, apex with a short point, glabrous, glossy above, silvery ferruginous villose beneath, venation pinnate, lateral veins 10–15 pairs, obvious, tomentose when young. *Petioles* 1.5–3 cm long, slightly swollen at base, tomentose. *Flower bud* enclosed by 3 caducous bracts. *Flowers* axillary, aromatic. *Pedicels* 3–3.5 cm long, ferruginous tomentose. *Tepals* 9, in 3 whorls, white. *Stamens* and carpels numerous, free, spirally arranged on a tomentose, cylindrical receptacle. *Fruit* compound, formed by many separate follicles, follicles narrow ovate, 30–45 by 15–18 mm, dehiscing by 2 valves when ripe, 1–5 seeds in each follicle. *Seeds* red when mature.

Distribution — China and Vietnam. In Vietnam found in almost all provinces of North and Central Vietnam. In Laos: not yet recorded, but most probably present.

Habitat & Ecology — Primary and secondary tropical and subtropical forests, altitude below 1000 m. Slow-growing tree, shade tolerant, prefers humid, deep, and fertile soil. Usually mixed with *Canarium album*, *Cinnamomum* spp., *Lithocarpus* spp., *Machilus* spp., and other *Michelia* species. Natural regeneration is good under thin forest covered. Flowering: April, May; fruiting: September, October.

Uses — Sapwood and heartwood are distinctive. Sapwood is brownish yellow, heartwood is brown. The timber is very hard, texture fine, aromatic, grain fine, resistant to termites and insects. Used for construction, making furniture and art articles. Seeds used as spices.

Specimens examined: *Balansa* 3886 (P); *Brillet s.n.* (P); *Chevalier* 40966, 41042 (P); *Domat* 38191 (P); *Eberhardt* 4460 (P); *Fleury* 30103 (P); *Forestier* 40937 (P); *Pételot* 3548, 5661, 6713, 8648 (P); *Poilane* 1534, 18893, 27556 (P); *Tsang* 26866 (P).

2. *Michelia mediocris* Dandy

Synonyms — *Magnolia mediocris* (Dandy) Figlar; *Michelia tonkinensis* A. Chev.

Vernacular names — Giỏi, Giỏi xanh, Giỏi tanh (V).

Trees up to 30 m high, up to 1 m diameter. *Trunk* straight, cylindrical. *Bark* greyish brown, glabrous, inner bark cream, with bad smell. Sapwood whitish. *Twigs* pale green. *Leaves* alternate, elliptic, 6–20 by 3–8 cm, coriaceous, base rounded or cuneate, apex with a short point or acuminate, both surfaces glabrous, green above, pale green beneath, venation pinnate, lateral veins 7–15 pairs, conspicuous. *Petioles* 1–2 cm long. *Flower buds* enclosed by 3 caducous bracts. *Flowers* terminal, axillary or opposite leaves. *Pedicel* 3–12 mm long, tomentose. *Tepals* 9 or 10, white. *Stamens* and carpels numerous, free, spirally arranged on a tomentose, cylindrical receptacle. *Fruit* compound, 3–7 separated follicles, 10 cm long when mature, 3–5 seeds in each follicle, follicles oblong ovate, 12–20 by 10–13 mm, lenticels many, dehiscing when ripe. *Seeds* red when mature.

Distribution — China, Vietnam, Cambodia. In Vietnam: in almost all provinces of North and Central Vietnam, mainly in Lào Cai, Hoa Binh, Thanh Hoa, Nghệ An, Hà Tĩnh, Kom Tum, and Gia Lai. In Laos: not yet recorded, but most probably present.

Habitat & Ecology — Primary and secondary tropical and subtropical forests, altitude up to 1000 m. Light demanding and rather fast-growing tree. Usually mixed with *Aglaia gigantea*, *Canarium album*, *Cinnamomum* spp., *Engelhardtia chrysolepis*, *Lithocarpus* spp., *Machilus* spp., *Madhuca pasquierii*, and other *Michelia* species. Natural regeneration is good under thin forest cover. There are two seasons. Main flowering: February, March; fruiting: September, October; minor flowering: July, August; fruiting: March, April.

Uses — Wood with distinct sapwood and heartwood. Sapwood is yellowish beige, heartwood is brownish yellow. Timber very hard, texture fine, easy to work, used for construction, boats, and making furniture. Seeds used as spices and as medicine to treat fever and stomachache.

Specimens examined: *Balansa* 3195 (P); *Hsien* 1078 (P); *Poilane* 297, 6497, 6531, 18054, 18181, 18229, 22018, 23172 (P).

MELIACEAE

Trees or shrubs, rarely with latex or sap. *Bark* smooth, cracked, fissured, scaly, or flaky, inner bark pinkish, white, or yellowish. *Stipules* absent. *Leaves* spirally arranged, alternate, rarely opposite, mostly compound, pinnate, sometimes bi- or tripinnate, rarely simple, leaflets usually opposite or subopposite. *Inflorescences* panicles, racemes, or spikes, sometimes reduced to fascicles or flowers solitary, axillary. *Flowers* bisexual or unisexual. *Calyx* cup-shaped with 3–5 lobes. *Petals* 3–7, free, narrow and curved backwards, green, white, pink, cream, or yellow. *Stamens* usually partly or completely united into tube or globose, 3–10 in 1 or rarely 2 whorls. *Disc* around ovary. *Ovary* superior, 1–6-locular, with 1 to many ovules per locule. *Fruit* a capsule, berry, or drupe. *Seed* winged and then attached to a woody columella, or covered by a sarcotesta or an aril, or without both, usually without endosperm.

Distribution — Throughout the tropical and subtropical regions of the world: c. 51 genera, 575 species. In Laos and Vietnam: c. 15 genera, c. 55 species.

KEY TO THE GENERA

- 1a. Slash of the bark without white latex. Stellate hairs or peltate scales absent. Stamens united into a tube 2
- b. Slash of the bark with white latex. Stellate hairs and/or peltate scales present. Stamens united into a globose head **1. Aglaia** (*spectabilis*)
- 2a. Leaves with pseudogemmula (bud at the end of the leaf). Fruit with 3 or 4 seeds, covered with an orange or red aril **2. Chisocheton**
- b. Leaves without pseudogemmula. Fruit with many winged seeds . . . **3. Chukrasia**

1. AGLAIA Lour.

Aglaia spectabilis (Miq.) S.S. Jain & Bennet — Fig. 26

Synonyms — *Aglaia gigantea* Pellegr.; *Amoora gigantea* Pierre; *A. spectabilis* Miq.

Vernacular names — Nook kok (L); Gôi nếp, Gôi (V).

Large trees up to 40 m high, 150 cm diameter. Usually with large buttresses. *Bark* greyish brown, flaking in squarish scales, sometimes with large orange lenticels, inner bark pinkish, reddish orange, or brown, latex white. *Twigs* covered with reddish brown or brown stellate hairs or scales. *Leaves* imparipinnate in dense spirals, up to 100 cm long. *Petioles* 9–25 cm long, flattened on the adaxial side, petiole, rachis, and petiolules covered with pale brown or reddish brown scales. *Leaflets* 5–9 pairs, opposite, ovate, 8–21 by 3–7 cm, coriaceous, upper surface usually rugulose, lower surface with pale brown or reddish brown stellate hairs or scales, apex acuminate, base rounded, lateral veins 7–16 pairs. *Inflorescences* panicles, up to 50 cm long. *Calyx* usually stellately hairy outside, 3- or 4-lobed. *Petals* 3, with or without hairs or scales outside. *Stamens* united, slightly shorter than corolla, with 5–10 (usually 6) anthers. *Disc* present. *Ovary* ovoid, 3- or 4-locular, 1 ovule per locule, style absent. *Fruit* a globose capsule 4–7 by 2–4.5 cm, red or yellow, densely covered with reddish brown or yellowish brown




Fig. 26. *Aglaia spectabilis* (Miq.) S.S.Jain & Bennet (Meliaceae).

hairs, a little bit splitting open when mature, pericarp usually with white latex. *Seeds* 3 or 4, black, covered with an orange-red or red aril.

Distribution — India, Myanmar, China, Indochina, Thailand, Malaysia, Philippines, New Guinea, Australia. In Laos: Sayabouri and Champassak provinces. In Vietnam: in many provinces in the north, mainly in Ninh Binh, Thanh Hoa, and Nghê An provinces.

Habitat & Ecology — Primary and secondary forests, altitude below 800 m. Growing on deep, sandy clay, and well-drained soils. Usually mixed with *Cryodaphnopsis tonkinensis*, *Dracotomelon dao*, *Dysoxylon cochinchinensis*, *Erythrophloeum fordii*, *Parashorea chinensis*, and *Pometia pinnata*. Fast-growing tree. Natural regeneration is good. Flowering: April, May; fruiting: September to November.

Uses — Sapwood pinkish, heartwood reddish brown. Used for making furniture, wall panelling, plywood, boxes, suitable for the production of pulp. This species can also be used as ornamental tree.

Specimens examined: *Charoenchai* 794 (L); *Delambre s.n.* (P); *Fleury* 29938, 29940, 30148, 37820 (P); *Harmand* 545, 882 (P); *Maxwell* 76-132, 76-175 (L); *Pierre* 55 (L), 1444 (P); *Poilane* 6322, 10739 (P).

2. CHISOCHETON Blume

Chisocheton cumingianus (C.DC.) Harms

Synonyms — *Chisocheton balansae* C.DC.; *C. chinensis* Merr.; *C. cochinchinensis* Pierre; *C. paniculatus* Hiern; *C. siamensis* Craib

Vernacular names — Kouang (L); Quyếch, Quyếch tía, Gội nước (V).

Trees up to 25 m high, 80 cm diameter. Usually with large buttresses. *Bark* greyish brown, inner bark chestnut brown, sapwood straw. *Twigs* covered with blackish brown tomentum. *Leaves* paripinnate or imparipinnate with pseudogemmula (apical bud) at tip, alternate, up to 100 cm long, petiole 5–14 cm long. *Leaflets* 5–10 pairs, opposite or subopposite, ovate to elliptic, 7–25 by 3–7 cm, papery to coriaceous, base nearly rounded and slightly oblique, margin entire, apex shortly cuspidate, dark green when mature. *Petiollules* 4–12 mm long. *Inflorescences* panicles, up to 50 cm long, axillary. *Flowers* bisexual. *Sepals* united, 4-lobed. *Petals* 4, united at lower part into a cylindrical tube, imbricate, yellow to white. *Stamens* tube cylindrical, as long as corolla, with 6 anthers. *Disc* present. *Ovary* ovoid, 4-locular, 1 ovule per locule, style hairy. *Fruit* a capsule, globose to pyriform, red or orange-red, 3–6 by 2.5–6 cm, splitting into 3–5 sections, but usually 4, pericarp usually with white latex. *Seeds* 3 or 4, black, covered with an orange aril.

Distribution — India, China, South East Asia. In Laos: Sayaboury province. In Vietnam: Yên Bái, Lào Cai, Tuyên Quang, Hà Giang, Vinh Phú, Hoa Bình, Hà Tây, Ninh Bình, Thanh Hoa, Nghệ An, Hà Tĩnh, Minh Hải, and Kiên Giang provinces.

Habitat & Ecology — Wet and very light conditions in secondary forests. Usually mixed with *Aglaiia gigantea*, *Albizia lucida*, *Caryodaphnopsis tonkinensis*, *Dalbergia balansae*, and *Pterocarpus tonkinensis*. Fast-growing tree. Natural regeneration is good. Flowering: January, February; fruiting: April, May.

Uses — Wood white and soft. Used for making furniture, wall panelling, plywood, boxes, and is suitable for the production of pulp. This species can also be used as ornamental tree.

Specimens examined — *Coode* 5454 (L); *Pierre* 1397 (L); *Soejarto et al.* 10209, 10254 (L); *Sonett* 1224 (P); *Sulit* 5662 (L); *Tanalt s.n.* (P).

3. CHUKRASIA A. Juss.

Chukrasia tabularis A. Juss.

Synonyms — *Chickrassia tabularis* (A. Juss.) Wight & Arn.; *Dysoxylum esquirolii* H. Lévl.; *Swietenia chickrassa* Roxb.

Vernacular names — Nhom, Nhom hin, Nhom khao (L); Lát, Lát hoa (V).

Deciduous trees up to 40 m high, up to 120 cm diam., buttresses present. *Bark* dark brown, fissured vertically and scaling or cracking into rectangular blocks, inner bark reddish brown or pinkish. Sapwood straw, heartwood yellow to reddish brown. *Twigs* grey. *Leaves* bipinnate with incised or lobed leaflets when young, 30–80 cm long. *Petioles* 4–9 cm long, swollen at base. *Leaflets* 7–13 pairs, alternate or subopposite, ovate to elliptic, 6–15 by 3–6 cm, base oblique, margin entire (when mature), apex tapering, venation pinnate, secondary veins 10–15 pairs, domatia present, petiolules 2–6 mm long. *Inflorescences* panicles, axillary, sometimes terminal. *Bracts* narrowly triangular, often caducous. *Flowers* bisexual. *Calyx* 5-lobed, densely brown hairy. *Petals* 5, free, slightly curved, spreading, hairy outside, glabrous inside. *Stamen tube* cylindrical, slight narrower towards top, glabrous, with 10 anthers attached to the margin. *Ovary* tomentose. *Fruit* a capsule, ovoid, woody, dark brown, 2–4.5 by 2–3.5 cm, opening by 3–5 valves from apex. *Seeds* many, 1–1.3 cm long, winged.

Distribution — Sri Lanka, India, Nepal, China, Japan, Indochina, Thailand, Malaysia, Indonesia (Borneo). In Laos: Houa Phan (Sam Nua Distr.) and Xieng Khouang provinces. In Vietnam: in many provinces, such as Lang Son, Son La, Lào Cai, Yên Bái, Lai Châu, Tuyên Quang, Bình Ninh, Hoa Binh, Thanh Hoa, Nghệ An, Hà Tĩnh, and Quang Binh.

Habitat & Ecology — Usually on limestone mountains, altitude below 800 m. Usually mixed with *Dracontomelum dao*, *Garcinia* spp., and *Peltophorum tonkinensis*. Slow growing and light demanding tree, but tolerant to shade when young. Natural regeneration is good. Flowering: June, July; fruiting: November to January.

Uses — Wood hard and heavy, grain fine. Used for construction, high grade cabinet work, decorative panelling, doors, windows, light flooring and making ships, boats, and furniture.

Specimens examined: *Brillet* 9, 12 (P); *Fleury* 30143 (P); *Harmand* 731, 907, 931 (P); *Kerr* 21292 (L, P); *Kostermans* 23460 (L); *Maxwell* 75-388, 91-923, 92-506 (L); *Pierre* 877, 6487, s.n. (P); *Poilane* 11526, 17818 (L); *Ridsdale* 16 (L).

MIMOSACEAE

Trees, shrubs or lianas, very rarely herbs (*Mimosa*, *Neptunia*); branches unarmed or armed with stipular thorns (rarely axillary thorns) or scattered prickles on the internodes. *Stipules* rarely absent, usually caducous; in some genera reduced to spines. *Leaves* alternate, pinnate or bipinnate, often with glands on the rachis. *Leaflets* many pairs, some with sleep movements, or touch-sensitive. *Flowers* regular (actinomorphic), crowded into globose heads or cylindric spikes, rarely racemose, some inflorescences monoecious. *Sepal lobes* 5, usually valvate. *Petals* 5, equal, valvate in bud, usually united above the base. *Stamens* few to numerous, free, joined at the base, or united,

forming a tube; anthers attached at the back. *Ovary* solitary or several, free, superior, stigma small, tubular, terminal. *Pods* dehiscent or indehiscent, straight, curved, or spirally twisted, sometimes breaking into 1-seeded segments. *Seeds* usually in two rows, ovate or orbicular, often compressed.

Distribution — Mainly in the tropical and subtropical regions of the world: c. 60 genera, c. 3000 species. In Vietnam: 13 genera, 62 species. In Laos: 12 genera, 34 species.

KEY TO THE GENERA

- 1a. Leaves with one pair of pinnae 2
- b. Leaves usually with more than one pair of pinnae 3
- 2a. Branchlets armed with spinescent stipules. Leaflets 1 pair per pinna, sessile **5. Pithecellobium** (*P. dulce*)
- b. Branchlets unarmed and without spinescent stipules. Leaflets more than 2 pairs per pinna, petiolulate **7. Xylia** (*X. xylocarpa*)
- 3a. Leaves with extrafloral nectaries on petioles, rachises, or pinnae. Leaflets opposite. Seeds brown or black when mature 4
- b. Leaves without extrafloral nectaries on petioles, rachises, or pinnae. Leaflets alternate. Seeds red when mature **1. Adenantha**
- 4a. Inflorescences pendulous or not, not consisting of clavate head 5
- b. Inflorescences consisting of pedunculate pendulous clavate heads **4. Parkia** (*P. sumatrana*)
- 5a. Flowers usually heteromorphic (excepted *Albizia procera*). Pods brownish or blackish outside when mature. Seeds with pleurogram 6
- b. Flowers uniform. Pods reddish or orange yellowish outside when mature. Seeds without pleurogram **3. Archidendron**
- 6a. Central flowers with 5 perianth segments. Pods dehiscent **2. Albizia**
- b. Central flowers with 7 or 8 perianth segments. Pods indehiscent . . . **6. Samanea**

1. ADENANTHERA L.

Adenantha microsperma Teijsm. & Binn. — Fig. 27

Synonym — *Adenantha pavonina* L. var. *microsperma* (Teijsm. & Binn.) I.C. Nielsen

Vernacular names — Lam ta khouay, Lam ta khway (L); Muồng ràng ràng (V).

Deciduous trees up to 30 m high, up to 90 cm diameter. *Crown* broad, rounded and large, branches spreading. *Bark* dark brown or greyish, flaking, inner bark pale cream. *Stipules* filiform, less than 0.5 mm long, puberulous, caducous. *Leaves* bipinnate, imparipinnate, 12–21 cm long, without extrafloral nectaries, pinnae opposite, 3–5(–6) pairs. *Leaflets* 5–9 per pinna, alternate, irregular ovate-elliptic or obovate-elliptic, 1.5–3.5 by 1–2 cm, base cuneate to rounded, apex rounded to truncate, often emarginate or mucronulate, dark grey-green above, pale grey-green below, both surfaces puberulous to glabrous. *Inflorescences* racemes, incl. peduncle 6–20(–25) cm long. *Bracts* filiform, 0.75 mm long, caducous. *Pedicels* 1.5–3 mm long, usually puberulous to sericeous (rarely glabrous). *Flowers* creamy-yellow, turning orange with age. *Calyx* 5-lobed,




Fig. 27. *Adenanthera microsperma* Teijsm. & Binn. (Mimosaceae).

shallowly cup-shaped, 0.5–0.75 mm long, puberulous to sericeous (rarely glabrous). *Petals* 5, narrowly ovate or elliptic, 2.5–3.5 mm long, apex acute, usually glabrous (rarely puberulous towards the apex). *Stamens* 10, free, 3–4.5 mm long; anthers 0.5 mm long, glabrous, with a gland at the tip. *Ovary* 1.6–2.2 mm long, densely sericeous, style 1–1.5 mm long. *Pods* brown to dark brown, strap-shaped, very thin, twisted in a tight coil, dehiscent, 15–20 by 0.8–1.2 cm. *Seeds* orbicular to elliptic, 5–8 by 4–7 by 4 mm, bright red, smooth, glossy.

Distribution — Myanmar, South China, Thailand, Indochina, Malesia. In Laos: Louang Prabang, Sayabouri, and Vientiane (Ban That Louang) provinces. In Vietnam: Bac Thai, Kiên Giang, Quang Ninh, Vinh Phu, Quang Nam-Dà Nang, Gia Lai, Kon Tum, Đông Nai, Hồ Chí Minh, Hà Nội, and Minh Hai provinces.

Habitat & Ecology — Dry evergreen and deciduous forest, forest margins, along streams and rivers, in gaps, altitude up to 600 m. Flowering and fruiting throughout the year.

Uses — Timber used for construction and for making implements. The tree is planted as a decorative and shade providing tree.

Specimens examined: *D'Alleizette* 1985 (L); *Balansa* 5, 1302, 2177, 4726, 4736 (L, P), 4839 (P); *Belval* 19 (P); *Bon* 2328, 2589, 4787 (P); *Carrick* 1602 (L); *Chevalier* 36759, 37605, 37749, 37881 (P); *Clemens* 4131 (P); *Donmat* 38159 (P); *Evrard* 2303 (P); *Fleury* 37614 (P); *Hiep* 2 (P), 539 (L, P); *Maxwell* 87-348 (L); *Pételot* 4935 (P); *Pierre* 730, 1816, 6552 (P); *Poilane* 42, 8334, 10113, 18152, 21347, 28943 (P), 9820 (L, P), 27132 (L); *Service forestier* 1 (P); *Soejarto* 5869 (L); *Talmy* 271 (P); *Tsang* 29859 (L, P), 27445, 29119, 29984 (P), 30449 (L); *Vidal* 829 (P).

2. ALBIZIA Durazz.

Shrubs, trees, usually unarmed, or lianas, or lianas armed with recurved hooks from the base of the leaf-scars. *Stipules* present, subular to linear or auriculate, often early caducous, rarely spinescent. *Leaves* bipinnate, usually more than one pair of pinnae, not sensitive to the touch, with glandular rachis and pinnae. *Leaflets* opposite, sessile or stalked. *Inflorescences* composed of pedunculate glomerules or corymbs which are either axillary or aggregated into terminal or axillary panicles; floral bracts small, linear-oblong, or absent. *Flowers* pentamerous, usually dimorphic (uniform for 5 species, incl. *Albizia procera*), the marginal flowers in each flowerhead bisexual, the central flower male and enlarged with a conspicuous nectary. *Sepals* connate, 5-toothed, valvate. *Petals* connate, valvate, 5-lobed (rarely 4). *Stamens* numerous, filaments united into a tube at the base; anthers eglandular, glabrous. *Ovary* solitary, sessile or stipitate. *Pods* chartaceous to coriaceous, straight, sometimes curved, flat, dehiscent; endocarp not separated from exocarp, brownish or blackish outside. *Seeds* dark brown or black, circular or elliptic, ± flattened, with a hard testa, wingless, with a pleurogram, aril absent; endosperm absent; cotyledons large, radicle curved.

KEY TO THE SPECIES

- 1a. Leaflets up to 6 pairs. Glomerules solitary from leaf-axils. Sepals more than 3 mm long. Ovary sessile **1. A. lebbeck**
- b. Leaflets 7 or more pairs. Glomerules in axillary or terminal panicles. Sepals less than 3 mm long. Ovary stalked 2
- 2a. Petiolules up to 1 mm long. Sepals 1–1.5 mm long. Petals up to 6.5 mm long. Ovary up to 1 mm long, hairy **2. A. odoratissima**
- b. Petiolules more than 1 mm long. Sepals 2.5–3 mm long. Petals up to 12.5 mm long. Ovary 0.3–0.5 mm long, glabrous **3. A. procera**

1. *Albizia lebbeck* (L.) Benth. — Fig. 28

Synonyms — *Acacia lebbeck* (L.) Willd.; *Mimosa lebbeck* L.

Vernacular names — Khan hong, Ka se (L); Lim vang, Mang lai (V).

Deciduous trees up to 30 m high, up to 90 cm diam., bole up to 15 m tall. *Crown* spreading, branches large, twisted. *Bark* dark brown, densely but shallowly cracked. *Stipules* inconspicuous. *Leaves* bipinnate, paripinnate, 5–10 cm long, with a gland c. 1.5 cm above the base of the petiole and often additional ones at the junction of some of the pinnae; pinnae opposite, 2–4(–5) pairs, up to 9 cm long. *Leaflets* opposite, 3–8 pairs per pinna, chartaceous, irregular trapezoid to rhomboid or obovate, 1.5–5.5 by

0.9–3.5 cm, base irregular broadly cuneate to truncate or obliquely cordate, apex rounded to truncate, lower surface tomentose to glabrous, upper surface glabrous to puberulous. *Inflorescences* consisting of 2 or more pedunculate corymbs in the distal leaf-axils or corymbs arranged in a short raceme at the end of the branches, peduncle 10 cm long, tomentose to sparsely puberulous, corymbs with 15–40 pedicellate, dimorphic flowers. Marginal *flowers* pentamerous, pedicel 4(–7) mm long. *Calyx* greenish yellow, tube-like, funnel-shaped, 3.5–5 mm long, puberulous, teeth narrowly triangular, 0.75–1 mm long. *Corolla* 5, greenish yellow, funnel-shaped, 7.5–11 mm long, tube glabrous, lobes broadly ovate, 2.5–4 mm long, puberulous at the apex. *Stamens* white to yellow at the base, pale green or yellowish green towards the top, staminal tube shorter than the corolla tube, 4–5 mm long. *Ovary* sessile, 3–4 mm long, glabrous. *Pods* pale straw-coloured to brownish, strap-like, gradually narrowing towards both ends, flat, 15–35 by 3–5 cm, glabrous, dehiscent along both sutures. *Seeds* flat, elliptic, 10 by 6–7 by 1–1.5 mm, areole c. 5 by 2 mm, with pleurogram parallel to the margins, open towards hilum.

Distribution — Tropical and subtropical Asia and Africa, but introduced and cultivated all over the tropics. In Laos: Louang Prabang and Vientiane provinces. In Vietnam: Hà Bac, Quang Nam-Dà Nang, Gia Lai, Long An, Hà Nội, Hồ Chi Minh, and An Giang provinces.

Habitat & Ecology — Often common in deciduous forest, mixed evergreen forest, open forest, altitude up to 750 m. Flowering: March to July; fruiting: April to August.

Uses — Used for general construction, furniture, bark pounded used for soap, planted as a shade tree because of the rather short trunk and wide crown.

Specimens examined: *D'Alleizette* 951 (L); *Bon* 5668 (P); *Chevalier* 29562, 39415, 39722 (P); *Clemens* 3266 (P); *Dussaud* 66 (P); *Evrard* 1667 (P); *Maxwell* 75-347, 96-428, 99-190 (L); *Pierre* 1860 (P); *Poilane* 5840, 8397 (L, P), 9638, 39999, 40048 (P); *Pételot* s.n. (P).




Fig. 28. *Albizia lebeck* (L.) Benth. (Mimosaceae).

2. *Albizia odoratissima* (L.f.) Benth. — Fig. 29

Synonyms — *Acacia odoratissima* (L.f.) Willd.; *Mimosa odoratissima* L.f.

Vernacular names — Khang hung, Du salen, Len (Louang Prabang prov.) (L); Xưá (Kien Gian), Hop hoàn thôm (V).

Deciduous trees up to 40 m high, up to 60 cm diam., branchlets terete. *Bark* dark grey, slightly cracked, peeling in irregular small flakes, inner bark reddish. *Stipules* filiform, c. 2.5 mm long, early caducous. *Leaves* bipinnate, paripinnate, 7–20 cm long; glands 1–2 cm above the base of the petiole and between the junctions of the 2 distal pairs of pinnae; lower gland elliptic, cushion-shaped, 2 mm diam.; upper gland circular, 1 mm



diam.; pinnae opposite, 3–5 pairs, 7–14 cm long. *Leaflets* opposite, 10–16 pairs per pinna, petiolulate (up to 1 mm long), chartaceous, broadly elliptic, distal pair obovate, 1.1–3.5 by 0.6–1.2 cm, base irregular cuneate to truncate, apex obtuse, rounded, often mucronate, both surfaces glabrous to tomentose, main vein removed 1/3–1/5 of the width of the leaflet from the upper margin. *Inflorescences* terminal panicles, 8–20 cm long, puberulous to tomentose, composed of pedunculate corymbs, 1–2 cm long, bearing heads of 10–15, sessile, dimorphic flowers. *Calyx* tube-like, funnel-shaped, 1–1.5 mm long, teeth triangular, less than 0.25 mm long, apex acute, puberulous to tomentose. *Corolla* funnel-shaped, 4.5–6.5 mm long, lobes ovate-elliptic, 2–2.5 mm long, apex acute, puberulous to tomentose. *Stamens* numerous, longer than the petals, filaments slender, filiform. *Ovary* puberulous to tomentose, stipe c. 1 mm long. *Pods* flat, pale brown, irregularly rectangular, 10–22 by 2–3.5 cm, glabrous to puberulous, dehiscent. *Seeds* 8–12, ovate, flat, 9 by 6 by 1.5 mm, areole 5 by 2.5 mm, pleurogram parallel to the margins.

Distribution — India, Myanmar, China, Thailand, Indochina. In Laos: Phong Sali, Luouang Namtha (Muong Sing), Louang Prabang (Pac Ou, Phou Soun, Phou Khouane), and Xieng Khouang (Muong Soui-Muong You) provinces. In Vietnam: Lai Châu, Kiên Giang, Dac Lac, Vinh Phu, Lâm Đông, Đông Nai, and An Giang provinces.

Fig. 29. *Albizia odoratissima* (L.f.) Benth. (Mimosaceae).

Habitat & Ecology — Open areas, dry deciduous forest, secondary forest, in fire-damaged forest, in hill mixed evergreen forest, hardwood forest, altitude 400–1500 m. Flowering: March to June; fruiting: May to December.

Uses — Wood used in construction, house-building, especially for high quality exterior joineries, gymnastic instruments, wood carving, and writing materials.

Specimens examined: *Averyanov et al.* VH 1322 (P); *Chevalier* 37287, 37696, 38080, 40286 (P); *Dussaud* 66 (P); *Fukuoka* T-63707 (L); *Maxwell* 96-510, 97-396 (L); *Pierre* 5965 (P); *Poilane* 3995, 26275 (L, P), 18167 (L), 20126, 20214, 20216, 23479, 26023, 26917 (P); *Schmid s.n.* (P); *Thorel* 99478 (P); *Vidal* 2532, 2605 (P).

3. *Albizia procera* (Roxb.) Benth. — Fig. 30

Synonyms — *Acacia procera* (Roxb.) Willd.; *Mimosa procera* Roxb.

Vernacular names — Thon (L); Mù cua, Muồng xanh (V).

Deciduous or semi-deciduous trees up to 30 m high, up to 60 cm diam., bole straight or crooked, up to 9 m. Bark greyish, glabrous, inner bark grey-brown. Branchlets terete, glabrous. Stipules filiform, c. 0.5 mm, or early caducous. Leaves bipinnate, paripinnate, 10–30 cm long, gland 1–1.5 cm above the base of the petiole, ± narrowly elliptic, sessile, flat and disc-like or with raised margins and concave, 4–10 mm long; pinnae opposite or subopposite, 2–5 pairs, 12–20 cm long, glabrous, with elliptic glands, c. 1 mm diam., below the junctions of the 1–3 distal pairs of petiolules. Petiolule c. 2 mm. Leaflets opposite, 5–11 pairs per pinna, petiolulate (> 1 mm long), chartaceous, ovate to elliptic or ± rhomboid, 2–4.5(–6) by 1–2.5(–3.2) cm, base irregular cuneate to truncate, apex rounded or subtruncate, often emarginate, mucronate, both surfaces sparsely appressed puberulous, main vein slightly excentric, often diagonal. Inflorescences composed of pedunculate glomerules aggregated into terminal or axillary panicles; peduncles 2–5 in fascicles, 0.8–3 cm, glomerules with 15–30 sessile, uniform, bisexual, 5-merous flowers. Calyx light green, tube-like funnel-shaped, glabrous, 2.5–3 mm long, teeth triangular, 0.75–1.2 mm long, apex acute. Corolla greenish white, tube-like funnel-shaped, 5–6.5 mm long, lobes elliptic, 2–2.5 mm long, apex acute, puberulous.




Fig. 30. *Albizia procera* (Roxb.) Benth. (Mimosaceae).

Stamens greenish white, numerous, staminal tube longer than the corolla-tube. *Ovary* glabrous, c. 1.5 mm long, stipe 0.3–0.5 mm long. *Pods* flat, dark brown or red brown, strap-like to rectangular, 11–20 by 2–2.5 cm, very short stalk, dehiscent, glabrous, veins inconspicuous, with distinct marks over the seeds. *Seeds* 8–12, obovate-elliptic, flattened, 7–8.5 by 4.5–6.5 by 1.5 mm, areole 4.5 by 3 mm, pleurogram nearly parallel to the margin, open towards the hilum.

Distribution — In India, Myanmar, South China, Taiwan, Indochina, Thailand, Malesia. In Laos: Louang Prabang, Vientiane, Savannakhet, and Champassak (Sithandone) provinces. In Vietnam: Quang Nam-Dà Nang, Quang Tri, Đông Nai, Sông Bé, and Tây Ninh provinces.

Habitat & Ecology — Tropical rain forest, dry open forest and savannah, mixed deciduous forest, altitude up to c. 1550 m. Flowering: June, July; fruiting: August to December.

Uses — Wood hard and heavy, used in construction, for domestic utensils, vehicles, agricultural tools, for moulding, furniture, cabinets; also for making charcoal.

Specimens examined: *D'Alleizette* 118 (P); *Balansa* 2179 (P); *Bon* 11 (P); *Chevalier* 29771, 36827 (P); *Fleury* 37767 (P); *Hardial* 574 (L); *Larsen* 3346 (L); *Pierre* 5968 (P); *Poilane* 11407, 27907 (L, P), 758, 7541, 11530, 13645, 18755 (P); *Schmid* 708 (P); *Thorel* s.n. (P); *Vidal* 1360, 4230 (P).

3. ARCHIDENDRON F. Muell.

Archidendron clypearia (Jack) I.C. Nielsen var. **clypearia** — Fig. 31

Synonyms — *Inga clypearia* Jack; *Pithecellobium clypearia* (Jack) Benth.

Vernacular names — Ben bay, Ben fay (Xieng Khouang) (L); Phân mã, Cút ngựa (V).

Shrubs or small trees 10–15 m high. *Trunk* straight and cylindrical with longitudinal grooves. *Bark* smooth to closely fissured, lenticellate, brown to pale grey, inner bark greenish yellow to purplish red. *Branchlets* angular to winged by decurrent ridges from the leaf-bases, puberulous to tomentose. *Stipules* sometimes present. *Leaves* bipinnate, paripinnate, 15–50 cm long, gland 1.5–2 cm above the base of the petiole, up to c. 5 mm long, flat or urceolate, elliptic or circular, upper glands below the junction of the pinnae, smaller, nearly circular, with slightly raised margins; pinnae 3–10 pairs, 2–15 cm long, glands just below the junctions of the petiolules, c. 0.5 mm diameter. *Leaflets* 3–10 pairs per pinna, opposite, petiolulate, ovate to trapezoid, 0.7–8 by 0.2–4 cm, base irregular cuneate, apex acuminate and often mucronate, upper surface faintly puberulous, lower surface faintly puberulous to tomentose, main vein diagonal, lateral veins prominent on lower surface. *Inflorescences* umbels or corymbs, with c. 10 pedicellate flowers, peduncle 1–2.5 cm long. *Flowers* uniform, pedicel 1–3(–4) mm long. *Calyx* and corolla connate, valvate. *Calyx* cup- to funnel-shaped, 1–3 mm long, teeth triangular to deltoid, 0.1–1 mm long, faintly puberulous to sericeous. Corolla funnel- to bell-shaped, 4–8 mm long, lobes ovate, 2–3 mm long, apex acute, puberulous to sericeous. *Stamens* many, staminal tube united with corolla tube at base. *Ovary* c. 1.5 mm long, puberulous to sericeous, stipe 1–1.5 mm long. *Pods* up to 20 by 1 cm, coriaceous to woody, compressed, twisted in a spiral, somewhat sinuate between the seeds, outside yellowish orange, inside reddish when mature, puberulous to tomentose, dehiscing first along the ventral suture. *Seeds* without pleurogram, endosperm, and aril,




Fig. 31. *Archidendron clypearia* (Jack) I.C. Nielsen var. *clypearia* (Mimosaceae).

dangling from the funicle after dehiscence, elliptic to globose, 7–10 by 6–9 by 6–7 mm, not occupying the cavity of the pod, with a bluish black testa.

Distribution — Sri Lanka, India, Bangladesh, Myanmar, South China, Indochina, Thailand, Malesia. In Laos: Louang Namtha, Xieng Khouang, Saravane, Champassak, and Attopeu provinces. In Vietnam: almost all provinces.

Habitat & Ecology — Primary and secondary rain forest, swamp forest, evergreen forest on clayey or grey silt, laterite, sand, and limestone soils, dry deciduous Dipterocarp forest and upper mixed deciduous forest, altitude up to 1700 m. Flowering and fruiting throughout the year, but mostly flowering from March to May and fruiting from April to July.

Uses — The timber is used for light construction, interior joinery, furniture and cabinet work, knife handles, weapon sheaths, boxes, coffins, and fuel.

Specimens examined: *Balansa* 2138, 2174, 2175 (P); *Bon* 1601, 3170, 3350 (P); *Chevalier* 30164 (L), 30685, 37430, 37476, 38729, 40390 (P); *Clemens* 3622 (P); *Contest* 342 (P); *Eberhardt* 2567, 2662, 2683, 4946, 5012 (P); *Evrard* 797 (L, P); *Fleury* 601, 30164, 38021 (P); *Jacquet* 634 (P); *Jean* 37036 (L, P); *Harmand* 1409, 5971, 5972 (P); *Ninh et al.* 9810 (L); *Pételot* 8853 (P); *Pierre* 5971 (P); *Poilane* 848, 1390, 2038, 5801, 5918, 13250, 15673, 18323, 22575, 23618, 23999, 25202, 26267, 26317, 28506, 29788, 30186, 31515, 32598 (P), 994, 10002, 11191, 11291, 18854 (L, P), 7226 (L); *Thorel* 601 (L); *Tixier s.n.* (P); *Tsang* 29201, 29903 (L, P); *Vidal* 2755 (P); *Vinot* 37 (P).

4. PARKIA R.Br.

***Parkia sumatrana* Miq.** — Fig. 32

Synonyms — *Parkia dongnaiensis* Pierre; *P. insignis* Kurz; *P. macrocarpa* Miq.; *P. streptocarpa* Hance

Vernacular names — Hua lon (Savannakhet), Som poy luang (Louang Prabang) (L); Cay thoi, Thui (Bien Hoa) (V).

Trees up to 35 m high, c. 40 cm diameter. *Branchlets* puberulous. *Bark* black, lenticellate. *Leaves* bipinnate, paripinnate, petioles (3.5–)5–10 cm long, gland 1–1.5 cm above the base of the petiole, elliptic, 2–7 mm long, 1 (or 2) inconspicuous glands below the

junctions of the terminal pair of pinnae, rhachis 13–30 cm long, pinnae 4–12 pairs, 10–15 cm long, opposite or subopposite. *Leaflets* 12–27 pairs per pinna, opposite, elliptic, 10–28 by 3–10 mm, base weakly auriculate on proximal side, apex rounded or slightly retuse, often slightly emarginate, midrib slightly oblique towards apex, lateral veins prominent. *Inflorescence* consisting of pedunculate pendulous clavate heads, 1.5–3 cm wide. *Flowers* white, pedicellate, bracts 6–13 mm long. *Sterile flowers*: calyx tube narrowly funnel-shaped, c. 7 mm long, with linear lobes, c. 5.5 mm long, apex acute; staminal tube nearly 6–7 mm long, anthers c. 0.8 by 0.5 mm. *Bisexual flowers*: calyx tube 11 mm long; corolla tube 12 mm long, lobes linear, c. 8 mm long, apex acute; staminal tube 7–8 mm long, anthers 0.5–2 mm long. *Ovary* 4 mm long, appressedly puberulous. *Pods* 35–56 by 1.5–2 cm (incl. stalk), straight or slightly twisted, swollen over the seeds. *Seeds* obliquely longitudinal, 10–20 by 11 mm, testa thin and black.

Distribution — South Myanmar, Indochina, Thailand, Peninsular Malaysia, Indonesia (Sumatra, Borneo, Sulawesi). In Laos: Sayabouri, Louang Prabang, Vientiane, Savannakhet, and Attopeu provinces. In Vietnam: Quang Tri, Vinh Phu, Quang Nam-Dà Nang, Khanh Hoa, Kiên Giang, Hồ Chí Minh, and Đông Nai provinces.




Fig. 32. *Parkia sumatrana* Miq. (Mimosaceae).

Habitat & Ecology — Scattered in dry evergreen forest, often along streams, hilly areas, on sandy, stony, or clayey soils, altitude 100–600(–900) m. Flowering: February or March.

Uses — Wood used for construction purposes. Fruit pulp, young leaves, and seeds edible. Bark has been used for bathing, the powdered bark has been used as local medicine, and as vermicide or against leeches.

Specimens examined: *Chevalier* 36760 (P); *Harmand* 1236, 2336 (P); *Pierre* 1425, 1435, 5817 (P); *Poilane* 1583, 3231, 6772, 11784, 11835, 13638 (P); *Service forestier* 30076 (P); *Thorel s.n.* (P); *Vidal* 1734, 2593 (P).

5. PITHECELLOBIUM Mart.

Pithecellobium dulce (Roxb.) Benth. — Fig. 33

Synonym — *Mimosa dulcis* Roxb.

Vernacular names — Kham thet, Khaam theed (L); Keo tây, Me keo (V).

Small to large trees up to 25 m high, 45–60 cm diameter. *Trunk* usually straight, cylindrical, and much branched. *Bark* grey or greenish, splitted 3–4 mm deep, nodose, inner bark whitish, 2–7 mm thick. *Branchlets* numerous, slender, brownish, sparsely

puberulous, glabrescent, armed with spinescent stipules, 1–1.5 cm long. *Leaves* bipinnate, petiole 0.3–5 cm long, with one gland at the junction of the pinnae, circular, with raised margins, 0.25 mm diam.; pinnae 1 pair, 0.2–1 cm long, with one gland at the junction of the leaflets, with small terminal stipular spines. *Leaflets* sessile, 1 pair per pinnae, irregular elliptic to obovate-elliptic, 0.7–5 by 0.3–2.3 cm, chartaceous, base irregular obtuse, apex obtuse-emarginate, both surfaces glabrous, lateral veins reticulate, hardly visible above, raised beneath. *Inflorescence* composed of pedunculate glomerules aggregated into terminal, densely puberulous panicles or racemes, peduncles 0.5–2 cm long, bearing a glomerule of 15–20 flowers. *Flowers* green. *Calyx* cup-shaped, 1–1.5 mm long, tomentose, teeth triangular, 0.3–0.4 mm long. *Corolla* funnel-shaped, tomentose, 3–4.5 mm long, lobes ovate, apex acute, 1 mm long. *Stamens* white, up to 9 mm, tube equalling the corolla-tube. *Ovary* 1–3 mm long, puberulous,




Fig. 33. *Pithecellobium dulce* (Roxb.) Benth. (Mimosaceae).

stipitate 0.1–0.2 mm long. *Pods* slightly flattened, 10–12(–15) by 1–1.6 cm, outside dark brown, inside reddish to pinkish, glabrous or slightly puberulous, dehiscent along both sutures, swollen over the seeds. *Seeds* black, glossy, obovate-oblong, often asymmetric in outline, flattened, 9–12 by 7–8 by 1–2 mm, funicle gradually thickened into a fleshy white or pink aril, pleurogram up to 7.5 by 3 mm.

Distribution — Native of Central America (Mexico), now cultivated and naturalized in most of the tropics, especially in areas with a seasonal climate as India, Myanmar, Indochina, Thailand, Malaysia. In Laos: Louang Prabang and Vientiane provinces. In Vietnam: Hà Bac, Hà Nội, Hồ Chí Minh, and Quang Nam-Dà Nang provinces.

Habitat & Ecology — Low and medium altitudes in both wet and dry areas under full sunlight, well-drained soil is best, usually clay soil, altitude 0–500 m. Flowering and fruiting: March to November.

Uses — Bark used for tanning and dyeing fishnets. The root bark is used to cure dysentery. The leaves when applied as a plaster can allay pain of venereal sores and relieve convulsions and taken with salt can cure indigestion, but can also induce abortion. The sweet aril is edible and used in lemonades. Timber soft but heavy used for general construction as posts or fuel. Used for hedges after pruning. Cattle and goats feed on the fallen fruits. Seed oil is also edible.

Specimens examined: *Balansa* 4823 (P); *Chevalier* 29530, 29612 (P); *Davidse* 9026 (L); *Dupuy* 265 (P); *Evrard* 2482 (P); *Kostermans* 24833, 25244 (L); *Malhotra* 26697 (L); *Perianayagam* 26623 (L); *Pételot s.n.* (P); *Pierre* 265 (P); *Pottier* 696 (P); *Spire* 891, *s.n.* (P); *Tatemi Shimizu et al.* 19470, 21666 (L); *Thorel s.n.* (P); *Zimmerman* 18 (L).

6. SAMANEA Merr.

Samanea saman (Jacq.) Merr. — Fig. 34

Synonyms — *Albizia saman* (Jacq.) F. Muell.; *Enterolobium saman* (Jacq.) Prain; *Inga saman* (Jacq.) Willd.; *Mimosa saman* Jacq.; *Pithecellobium saman* (Jacq.) Benth.

Vernacular names — Sam sa (L); Me tây (V).

Deciduous or semi-deciduous trees up to 30 m high, c. 100 cm diam., with a large crown spreading to 33 m in a complete canopy. *Branchlets* puberulous to tomentose. *Stipules* present. *Leaves*: rhachis 15–40 cm long, with gland(s) just below the junction of the basal pair of pinnae and distally at all other pairs of pinnae, circular, concave, c. 0.5 mm diam.; pinnae 3–9 pairs, c. 11 cm long, with gland(s) at the junctions of the leaflets. *Leaflets* opposite, the lowest pair ovate or elliptic, middle pairs rhomboid, terminal pair obliquely obovate, 1.5–2.5 by 3–5 cm, 2–10 pairs per pinnae, base half rounded and half truncate, apex rounded or obtuse, often emarginate or mucronate, main vein diagonal, lateral veins densely reticulate, raised, upper surface glabrous, lower surface densely short-pubescent. *Inflorescence* peduncles densely shortly yellowish pubescent, 2–5 together in the distal leaf-axils, 5–10 cm long, bearing a terminal corymb. *Flowers* usually heteromorphic, marginal flowers c. 3 cm long, pedicellate, central flowers with 7 or 8 perianth segments. *Calyx* funnel-shaped, 5–7 mm long, tomentose or woolly, teeth 5, broadly triangular, acute, 0.5–1 mm long. *Corolla* red or yellowish red, funnel-shaped, 10–12 mm long, distal part tomentose or woolly, lobes triangular-ovate, c. 2 mm long. *Stamens* white at base, purple toward the top, 20–35 mm long, tube shorter




Fig. 34. *Samanea saman* (Jacq.) Merr. (Mimosaceae).

than the corolla-tube. *Ovary* sessile, glabrous. Central flower sessile. *Calyx* 8–9 mm long, broadly tubular, tomentose-woolly, teeth 7–8, 0.5–1 mm long, triangular, acute. *Corolla* c. 12 cm long, tubular, inside glabrous, outside tomentose-woolly. Staminal tube longer than the corolla. *Ovary* sessile, glabrous. *Pod* strap-like, 15–20 by 1.5–2.3 cm long, straight, indehiscent, outside black when mature, inside transversely septate. *Seeds* with pleurogram, elliptic, strongly biconvex, c. 8 by 5 by 4 mm, brown, areole elliptic, c. 7 by 3 mm.

Distribution — Native of northern tropical South America, now planted and appearing spontaneous all over the tropics. In Laos and Vietnam: occurring in all parts with tropical climate.

Habitat & Ecology — Sandy, coastal areas, along roadsides, often planted, altitude up to 1800 m. Flowering and fruiting: June to January.

Uses — The tree is widely cultivated as an ornamental along streets and around buildings. Wood hard and heavy, used in construction, furniture, boxes, and crates. Pods used as fodder for cattle, pigs, and goats.

Specimens examined: *D'Alleizette* 10 (P); *Chevalier* 31389 (L), 37291, 38060 (P); *Derral* 546 (L); *Evrard* 2507, 2647 (P); *Fleury* 38047 (P); *FRI* 22182 (L); *Heyligers* 1399 (L); *Hiep* 123, 952 (P); *Martin et al.* 4316 (L); *Muller* 1044 (P); *Perohm Kingnate* 8 (L); *Seng Houng* 93 (P); *Smith* 8251 (L); *Sykes et al.* 169773 (L).

7. XYLIA Benth.

Xylia xylocarpa (Roxb.) Taub. var. *kerrii* (Craib & Hutch.) I.C. Nielsen — Fig. 35

Synonyms — *Acacia xylocarpa* (Roxb.) Willd.; *Inga xylocarpa* (Roxb.) DC.; *Mimosa xylocarpa* Roxb.; *Xylia dolabrifomis* Benth.; *X. kerrii* Craib & Hutch.

Vernacular names — Deng (L); Cãm xe (V).

Large deciduous trees up to 30 m high. *Trunk* straight, branches slender, drooping, branchlets unarmed and without spinescent stipules. *Bark* creamy brown or red-brown, thin, peeling in rounded flakes, small lenticels, inner bark pink. *Twigs* densely puberulous, glabrescent. *Stipules* filiform, 3–4 mm long. *Petiole* 3–8 cm long, terete, puberulous to tomentose. *Leaves* bipinnate, paripinnate, glabrescent, gland at the junction of the pinnae 1–3 mm diam.; pinnae 1 pair, 7–30 cm long, slightly sulcate, tomentose, glabrescent; gland just below the junction of the petiolules, 1–2 mm diameter. *Leaflets* 3–7 pairs per pinnae, top ones largest, narrowly ovate or elliptic, 3.8–15 by 2.4–6.7 cm, mature leaves upper surface glabrous, lower surface puberulous to velutinous, petiolules 2–3 mm long. *Young shoots* densely covered with yellowish hairs. *Young leaves* delicate pink. *Inflorescence* peduncles 2.5–9 cm long, bearing heads of c. 90 sessile flowers. *Bracts* spoon-shaped. *Calyx* funnel-shaped, 2.9–4 mm long, tomentose to woolly, teeth 0.8–1 mm long, triangular-ovate, acute. *Petals* narrowly elliptic, 3.5–4.5 mm long, slightly fused at base, hairy outside. *Stamens* 10–12, free, 5–12 mm long, filaments very flat, anthers without glands. *Ovary* 2–3 mm long. *Pod* kitchen knife-shaped, 10–17 by 3.5–6 cm, thick and woody, slightly curved, tapering




Fig. 35. *Xylia xylocarpa* (Roxb.) Taub. var. *kerrii* (Craib & Hutch.) I.C. Nielsen (Mimosaceae).

at base, pale creamy brown at first, later dark brown, splitting suddenly into 2 parts which curl backwards, remaining on the tree for a long time. *Seeds* 6–10, ellipsoid, flat, 11–20 by 7–12 mm, dark brown.

Distribution — Myanmar, Indochina, Thailand. In Laos: Vientiane, Savannakhet, Saravane, and Champassak (Sithandone) provinces. In Vietnam: Dac Lac, Gia Lai, Đông Nai, Hồ Chí Minh, and Ninh Thuận provinces.

Habitat & Ecology — Dry evergreen forest, mixed deciduous forest, dry deciduous Dipterocarp forest, sandy deep soils, altitude up to 700 m. Flowering: February to June; fruiting: April to December.

Uses — Wood extremely hard and durable, resistant to termites, insects, and weather. Usually used for furniture, ships and boats, bridges, railway sleepers, piles, and in construction.

Specimens examined: *Chevalier* 30913, 31381, 36845, 39151 (P); *Dournes* 276 (P); *Elusel* 186 (P); *Fleury* 29929, 30920, 39309 (P); *Harmand* 446 (P); *Hiep* 498 (P); *Lecomte* 2011 (P); *Pierre* 26, 93, 119, 120, s.n. (P); *Poilane* 8786, 28306 (L, P), 5636, 9301, 9349, 9541, 11560, 13777, 40841 (P); *Schmid* 7125 (P); *Thorel* 998 (P); *Vidal* 1079, 2456 (P).

MORACEAE

Evergreen or deciduous trees, monoecious or dioecious, usually with white latex in the bark, leaves, and fruits. *Stipules* present. *Leaves* simple, mostly alternate, rarely opposite, sometimes lobed. *Inflorescences* a cyme, catkin, or head in the leaf axils or along trunk and main branches, (male and female always in separate inflorescences on the same or different trees). *Flowers* minute, pale green, more or less enlarged end of the flower-stalk bearing the flower-parts, rarely interior (*Ficus*). *Male flowers*: densely clustered in dangling catkins near the end of twigs, 5–8 cm long. *Sepals* free or connate. *Petals* absent. *Stamens* 4, longer than sepals and opposite them. *Pistillode* central, small, or absent. *Female flowers*: sepals sometimes different from male flowers. *Petals* absent. *Ovary* usually superior, sometimes adnate to the sepals, ovule solitary, style(s) 1 or 2, simple or forked. *Fruits* simple or compound, very various, often united into large compound structures. *Seed* 1.

Distribution — Almost confined to the tropics: 16 genera, c. 1100 species. In Laos and Vietnam: c. 8 genera, 125 species.

BROUSSONETIA Vent.

Broussonetia papyrifera (L.) Vent.

Synonym — *Morus papyrifera* L.

Vernacular names — Sa lae (Xiang Khouang), Sa (Sayaboury), Por sa (general) (L); Dường, Me day (V).

Deciduous shrubs or small trees up to 20 m high, c. 50 cm diameter. *Trunk* cylindrical. *Crown* broad, spreading. All parts of the tree with milksap. *Bark* cream or pale brown, smooth or shallowly cracked with large lenticels, outer bark thin and fibrous, inner bark with white latex. *Stipules* narrowly triangular or ovate, 6–8 mm long, early caducous. *Petioles* 3–10 cm long, slender, hairy. *Leaves* alternate or subopposite, clustered

near the end of twigs, broadly ovate or triangular, 6–29 by 5–12 cm, 1–5-lobed, base rounded or subcordate, sometimes cuneate, margin serrate, teeth variable in size, apex acuminate, young leaves densely covered with soft hairs, mature leaves thin, rough above, softly hairy below, basal veins 3–5, lateral veins 3–7 pairs. *Flowers* unisexual. *Male flowers* densely clustered in dangling catkins or spikes near the end of twigs, 5–8 cm long, individual flowers intermingled with stalked bracts. *Sepals* free in bud, hairy outside. *Stamens* 4, opposite the calyx lobes and longer than these, curved inwards in bud. *Female flowers* in globular heads, axillary, 1.2–2 cm diam., bracts subulate, pubescent at the apex, 3 or 4 teeth, stalk 0.5–1 cm long, individual flowers intermixed with many scales. *Calyx* tubular with 2–4 minute teeth, sparsely hairy outside. *Ovary* superior, style red, filiform, attached on the side of the ovary, much longer than calyx. Ovaries and scales connate to form a multiple fruit (syncarp), globose, about 3 cm diam., bright orange or red when mature. *Seeds* many, red, flattened.

Distribution — India, Myanmar, China, Indochina, Thailand, Malaysia, Indonesia (Java), Japan. In Laos: Sayaboury (Pak Lai), Louang Prabang, Xieng Khouang, Khammouane (Thakhek), and Attopeu provinces. In Vietnam: Hà Nội, Quang Tri, Thanh Hoa, Khanh Hoa, Nghệ An, Hà Tây, and Dac Lac provinces.

Habitat & Ecology — A light demanding and fast-growing tree, common in open areas, secondary forest, thickets, mountain ravines, along streams in deciduous forest, preferring moisture, altitude up to 1600 m. Natural and coppice-regeneration are strong. Flowering: November to June; fruiting: June to December.

Uses — Fibrous bark used for pulp in paper production. In Laos (Xiengkhouang prov., Phou Koud Distr.) traditionally produced paper is used to make paper altars and umbrellas (khan gnou) needed during religious ceremonies. Bark and fibres are exported. The inner bark is beaten by sticks to free the fibres. These are used in making ‘tapa’ cloth or ‘kappa’ in ancient Polynesia. Leaves are used as food for pigs and rabbits. Fruit, bark, and root are used in medicine as tonic for kidney problems and as diuretic. Seeds contain oil, used in soap and in lacquer oil production.

Specimens examined: *Balansa* 2462, 2463, 4111 (P); *Bon* 2934, 3202, 4992 (P); *Chevalier* 32422 (P); *Clemens* 4128 (P); *Dournes* 378 (P); *Eberhardt* 3430, 4018, 4529 (P); *Fleury* 30187 (P); *Harmand* 1296, 3295 (P); *Ninh & Hoang* 11450 (L); *Pételot* 23 (P); *Poilane* 1143, 1194, 2033, 6293, 7049, 10795, 31531 (P); *Pottier* 811 (P); *Spire* 419 (P); *Vidal* 1265, 2082 (P).

MYRISTICACEAE

Trees or shrubs 3–40 m high, dioecious, sometimes buttresses present. *Bark* usually with copious red or orange sap. *Branches* more or less horizontal, usually whorled, crown often pyramidal. *Stipules* absent. *Leaves* simple, alternate, usually distichous, margin entire, often tomentose on both surfaces when young and later glabrous or sometimes tomentum remaining on the lower surface. *Male inflorescences* axillary amongst or behind the leaves, or on the stems, panicles, or racemes, pedunculate or sessile, with scars of previous pedicels and bracts; flowers usually in racemose umbels; bracts present, often caducous; bracteole absent or present; pedicels usually present. *Female inflorescences* similar to male, usually less branched and shorter; flowers smallish. *Perianth* gamophyllous, 2–5-lobed fused at the base, sometimes split to the base, the female

flowers usually larger than the male ones, inside creamy or red. *Stamens* many, filaments united into a column. *Ovary* single, superior, subglobose, style short or absent, stigma 2-lobed, ovule 1, at the base of the ovary. *Fruits* a more or less woody capsule, elliptic, splitting into 2 sections, pericarp usually thick, coriaceous, glabrous or hairy, yellow or red when mature. *Seed* single, large, aril and seed aromatic, aril red or orange-red, fleshy, covering the seed, variously lobed or entire, endosperm ruminant.

Distribution — In all tropical areas: 20 genera, 500 species. In Laos and Vietnam: c. 2 genera, 20 species.

KEY TO THE GENERA

- 1a. Bracteole absent. Leaves when dry easily breaking into pieces, lower surface brown, upper surface dull. Inflorescences a loose panicle **1. Horsfieldia**
 b. Bracteole present in the apical part of the pedicel. Leaves when dry not easily breaking into pieces, lower surface usually glaucous or whitish, upper surface glossy. Inflorescences short umbel-like racemes **2. Knema**

1. HORSFIELDIA Willd.

Horsfieldia amygdalina (Wall.) Warb. — Fig. 36

Synonyms — *Horsfieldia thorelii* Lecomte; *H. tonkinensis* Lecomte

Vernacular names — Han, Hant (Savannakhet), Lueat nok (Vientiane) (L); Sang máu, Máu chó lá lớn (V).

Small to large trees 5–30 m high. *Twigs* towards apex terete, first hairy, then glabrescent. *Bark* finely striate and with lenticels. *Leaf bud* hairy. *Leaves* broadly elliptic, 9–23 by 2.5–9 cm, when dry easily breaking into pieces, base cuneate, apex acute or acuminate, upper surface dull, olive green or dark brown or blackish when drying, lower surface brown, soon glabrescent, midrib flat to slightly raised above, secondary veins 7–14 pairs, venation faintly reticulate. *Inflorescences* a loose panicle, axillary, thinly pubescent with pale hairs. *Bracteoles* absent. *Male inflorescences* 4–22 cm long. *Female inflorescences* 1–6 cm long. *Male flowers* short-obovoid or globose, 1.5–2.5 by 1.5–2 mm, lobes splitting the bud to c. halfway, androecium globose or obovoid, largely solid, 0.8–2 by 0.8–1.3 mm, anthers 10–15, sessile. *Female flowers* ellipsoid, 2–3 by 2 mm. *Ovary* glabrous. *Fruits* ellipsoid, 2–6 by 1.6–2.6 cm, glabrous, brown on drying, stalk 2–4 mm long. *Seed* ovoid, aril entire.

Distribution — India, Bangladesh, Myanmar, South China, Indochina, Thailand. In Laos: Savannakhet province. In Vietnam: Kiên Giang, Ninh Bình, and Đông Nai provinces.

Habitat & Ecology — Evergreen and deciduous forest, altitude up to 650 m. Flowering: September, October; fruiting: January to May.

Uses — Wood used for temporary construction and general furniture. Seeds contain a solid oil used for industrial purposes.

Specimens examined: *Pierre 56, 1812, 5434* (P); *Poilane 10, 1348, 8699, 18160, 18308, 18615, 22394, 23253* (P).



Fig. 36. *Horsfieldia amygdalina* (Wall.) Warb. (Myristicaceae).

2. KNEMA Lour.

Knema globularia (Lam.) Warb.

Synonyms — *Knema corticosa* Lour.; *Myristica corticosa* (Lour.) Hook.f.; *M. globularia* Lam.

Vernacular names — Thom luat (Savannakhet), Luat ma (Vientiane, Na Hai) (L); Máu cho lá nhỏ, Máu chó lá bé (V).

Trees 5–20 m high, 30–35 cm diameter. *Trunk* straight, fluted. Young branches yellow-brown, first tomentose, latter glabrous. *Bark* usually smooth, rarely cracked, never flaking. *Leaves* narrowly elliptic or narrowly ovate, 6–19 by 2–4 cm, thin, coriaceous, when dry not easily breaking into pieces, base cuneate, margin entire, apex acuminate, upper surface glossy and dark green, usually pale brown with blackish metallic shine when dry, lower surface pale green, when dry usually glaucous or whitish, midrib sunken above, secondary veins 10–15 pairs, faint, flat or sunken above, prominent beneath. *Inflorescences* short umbel-like, racemes, usually at least partly 1–5 mm pedunculate, simple or furcate, to 10 mm long. *Male inflorescences* with 5–20 flowers. *Female inflorescences* with 1–10 flower(s). *Male flowers*: pedicel 2–11 mm long, bracteole above median, bud broadly obovoid, 3–5 mm diam., creamy or reddish inside, lobes splitting the bud to c. 2/3 deep, staminal disc circular or 3-angled, flat, including anthers 1.5–2.5 mm diam., anthers 8–13, free. *Female flowers*: pedicel 1.5–4 mm long, bracteole nearly apically, bud ellipsoid-ovoid, 4–5 by 2.5–3.5 mm. *Ovary* ovoid to globose, tomentose. *Style* short, stigma lobed. *Fruits* ellipsoid, subglobose, or sometimes pear-shaped, 1.2–2 by 1–1.5 cm, stalk 3–20 mm long, pericarp thin. *Seed* smooth or glossy, aril entire.

Distribution — India, Myanmar, China, Indochina, Thailand, Peninsular Malaysia, Singapore, Indonesia (Sumatra, Java). In Laos: Vientiane, Vientiane Municipality, and Savannakhet provinces. In Vietnam: Vinh Phu, Đông Nai, Nghệ An, Quang Tri, Lào Cai, Thanh Hoa, and Khanh Hoa provinces.

Habitat & Ecology — Primary and secondary evergreen forest, along river banks, in hilly forest, altitude 150–1200 m. Flowering: June to December; fruiting: December to June.

Uses — Wood used for general furniture. Seeds used in medicine to cure scabies and other skin-diseases.

Specimens examined: *Balansa* 4198 (P); *Chevalier* 43, 37053, 37054, 38283, 38438, 38566, 39179 (P); *Fleury* 5472, 30009, 30114, 30145, 38000, 38438, 39348 (P); *Harmand* 602 (P); *Hiep* 429 (P); *Muller* 898, 1002 (P); *Pételot* 1546 (P); *Pierre* 260, 5434 (P); *Poilane* 119, 131, 1581, 1741, 6600, 10198, 10486, 17567, 20004, 25188, 29203, 29919 (P); *Vidal* 1756, 2250 (P).

MYRTACEAE

Trees or shrubs. *Stipules* absent. *Leaves* simple, opposite (rarely alternate or whorled), entire, pinnately veined, often with an intramarginal vein, with dots. *Inflorescences* complex cymes, sometimes racemes, many-flowered, rarely solitary. *Flowers* white or pink, regular, bisexual, rarely unisexual, (3- or 4- or 5- (or 6-)merous. *Sepals* and *petals* free or forming a deciduous cup, usually orbicular. *Stamens* usually many, free or united at the base into 4 or 5 bundles, arranged at the edge of the hypanthial cup,

anthers usually attached at the back. *Ovary* usually more or less inferior, 2–5-locular, ovules two to many per locule; style 1, often filiform, stigma minute and head-like. *Fruit* usually a berry or capsule, rarely a drupe, often with persistent sepal lobes. *Seed(s)* 1 to many.

Distribution — Worldwide: Large tropical family: c. 70 genera, 4620 species. In Laos and Vietnam: c. 12 genera, 69 species.

SYZYGIUM Gaertn.

Synonyms — *Caryophyllus* L.; *Eugenia* auct. non L.; *Jambosa* Adans.; *Opa* Lour.

Shrubs to large trees. *Trunk* sometimes buttressed or fluted. *Bark* various, often papery, flaky, or peeling; sapwood pinkish, red, yellow, or white, often turning purplish on exposure. *Leaves* opposite, rarely spirally arranged or whorled, intramarginal veins usually present, sometimes absent. *Inflorescences* many-flowered terminal or axillary panicles or racemes, sometimes few-flowered cymes. *Bracts* and bracteoles usually caducous, sometimes persistent. *Flowers* 4- or 5-merous, sessile or pedicellate, glabrous in bud. *Calyx* lobes broadly funnel-shaped or club-like, rarely cylindrical, persistent or caducous early. *Petals* free, sometimes united in a cap-like structure. *Stamens* free, usually many. *Ovary* 2- (rarely 3- or 4-)locular, ovules few or many per locule, style as long as or shorter than the stamens. *Fruit* a berry, variously shaped, apex usually crowned by the persistent remnants of the calyx tube or by enlarged calyx lobes, or both. *Seed(s)* 1 or 2, rarely more.

KEY TO THE SPECIES

- 1a. Leaf elliptic or ovate-elliptic, about 2 times longer than broad, secondary veins 19–30 pairs, intramarginal vein 1. Inflorescences terminal or axillary panicles. Flowers sessile **1. *S. cumini***
 b. Leaf narrowly ovate, about 4 times longer than broad, secondary veins 11–18 pairs, intramarginal vein(s) 1 or 2. Inflorescences terminal racemes. Pedicel 3–13 mm long **2. *S. jambos***

1. *Syzygium cumini* (L.) Skeels

Synonyms — *Eugenia cumini* L.; *E. jambolan* Lam.; *Myrtus cumini* L.

Vernacular names — Sa (Xiang Khouang), Sa leng, Va (L); Vối rừng (V).

Trees up to 35 m high. *Twigs* angular when young, later terete. *Bark* grey, slightly flaking, inner bark reddish. *Petiole* 6–28 mm long. *Leaves* elliptic or ovate-elliptic, 6–13 by 3–7 cm, index 2, base cuneate, acute, or rounded, apex shortly acuminate, secondary veins 19–30 pairs, intramarginal vein 1. *Inflorescences* terminal or axillary panicles, several together, shorter than the leaves, 4–10 cm long, peduncles 3–10 mm long, branches and branchlets striate. *Bracts* triangular, 7 by 10 mm. *Bracteoles* triangular, 6 by 4 mm. *Flowers* white or pale yellow, sessile. *Hypanthial* cup 2.4–5.1 mm long, funnel-shaped. *Pseudostipe* 1–2 mm long. *Sepals* 4, triangular or rounded, obscure, 2.5–6 mm long. *Petals* 4, joined into a cup, 1.4–2.8 mm long, early caducous, 5–19 gland

dots per petal. *Outer stamens* 4–6 mm long, anthers elliptic, 0.4–0.5 mm long. *Ovary* 2-locular, ovules 10–18 per locule, style 1.8–6.5 mm long. *Fruits* ovoid or ellipsoid, 8–15 mm long, pink turning dark red-purple or black when ripe. *Seed* 1, ellipsoid.

Distribution — Widely distributed in the Indo-Malayan region. In Laos: Sayaboury (Pak Lay Distr.), Xieng Khouang, Vientiane, Savannakhet (Sepone Distr.), and Attopeu provinces. In Vietnam: Vinh Phu and Đông Nai provinces.

Habitat & Ecology — Widespread in many types of forest. Often found in open and semi-open deciduous evergreen forest, conifer-broadleaved forest, or individual trees on grassland, altitude 0–1100 m. Flowering: March, April; fruiting: June, July.

Uses — Wood is pink and hard, used for fuel, construction, and furniture. The fruits are edible. The juice of the leaves is used for treatment of dysentery and for cleaning external wounds. Local people treat mild forms of diabetes.

Specimens examined: *Dieu* 56 (P); *Harmand* 3190, *s.n.* (P); *Hosseus* 436 (P); *Pételot* 4679 (P); *Pierre* 14, 1812 (P); *Poilane* 13480, 22394, 23253 (P).

2. *Syzygium jambos* (L.) Alston

Synonyms — *Eugenia jambos* L.; *Jambosa vulgaris* DC.

Vernacular names — Chieng, Kieng (L); Gioi (V).

Trees up to 15 m high. *Twigs* slender, compressed, or angular. *Bark* whitish grey or brown. *Petiole* 7–14 mm long. *Leaves* narrowly ovate, 8–20 by 1.5–5 cm, index 4, base cuneate or obtuse, apex acute or acuminate, secondary veins 11–18 pairs, intramarginal vein(s) 1 or 2. *Inflorescences* terminal racemes, 3–8-flowered, rarely 1-flowered, up to 6 cm long; peduncles 30–50 mm long, rachis and branches terete. *Bracts* triangular, 7 by 10 mm. *Bracteoles* triangular, 6 by 4 mm. *Flowers* white or pale yellow, pedicel 3–13 mm long. *Hypanthial* cup 12–14 mm long, broadly funnel-shaped. *Pseudostipe* 4–6 mm. *Sepals* 4, semi-orbicular, 5–8 mm long. *Petals* 4, free, orbicular, 15–17 mm long, base thick, more than 200 gland dots per petal. *Outer stamens* 16–37 mm long, anthers elliptic, 1–1.2 mm long. *Ovary* 2-locular, ovules c. 28 per locule, style 27–43 mm long. *Fruits* depressed globose, 3–6 cm diam., pale pink when ripe. *Seed(s)* 1 or 2.

Distribution — Native to the Indo-Malayan region, widely cultivated in tropical regions. In Laos: Vientiane and Champassak (Lakhon Distr.) provinces. In Vietnam: Hà Bac, Hà Nội, Ninh Bình, Sơn La, Thanh Hoa, Nghệ An, Dac Lac, and Đông Nai provinces.

Habitat & Ecology — Usually planted as a fruit-tree in villages or plantations. Sometimes escaping cultivation and growing wild along streams or rivers. Suitable on wet and deep soils near river banks. Light demanding when mature. Altitude up to 300 m. Flowering: April, May; fruiting: July, August.

Uses — Wood is used locally in general construction. Fruit edible, sweet and delicious. The tree is planted as wind shelter.

Note — According to Parnell & Chantaranonthai, *Flora of Thailand* 7 (2002) the width of the leaves should be 3–4.5 cm. We observed some specimens with much narrower, 1.5–2 cm wide leaves (e.g. *Spire* 703, 1025). Other characters, however, are not different.

Specimens examined: *Balansa* 2861 (P); *Bon* 359, 3312, 4191, 30092 (P); *Butreau* 12 (P); *Chevalier* 30092 (P); *Clemens* 3509 (P); *Eberhardt* 2778 (P); *Fleury* 29923, 37551 (P); *Lei* 308 (P); *Molliard* s.n. (P); *Pételot* 1832, s.n. (P); *Pierre* 93, 1962, s.n. (P); *Schmid* 996 (P); *Simond* 184 (P); *Spire* 703, 1025 (P); *Thorel* 2790, s.n. (P).

PAPILIONACEAE

Herbs, shrubs, or trees. *Stipules* usually present. *Leaves* spirally arranged, simple or compound, pinnate or digitate. *Flowers* mostly pea-flower-shaped, in terminal or axillary 1- to numerous-flowered inflorescences, usually bisexual. *Sepals* joined to form a cup or tube with 5 teeth. *Petals* mostly 5, imbricate, free or almost so, 2 lowermost forming together the keel (carina) enclosed by the 2 more or less parallel lateral ones (wings, alae), topmost petal (standard, vexillum) in bud embracing the other ones. *Stamens* mostly 10 (rarely 8 or 9), connate or free, all perfect or partially reduced to staminodes. *Ovary* superior, 1-celled, with 1 to many ovules. *Fruit* a pod. *Seed(s)* 1 to numerous, septate between the seeds or not, or articulate, when mature dehiscent or not, frequently breaking up into 1-seeded articles. Seed coat with complex structure in region of scar (hilum), in transverse section of a mature seed the albumen is visible on either side of the embryo.

Distribution — Worldwide: c. 505 genera, c. 13,800 species. In Laos: 70 genera, 267 species. In Vietnam: 86 genera, 385 species.

KEY TO THE GENERA

- 1a. Trunk and branches unarmed. Leaves imparipinnate. Stipels absent. Flowers white or bright yellow. Fruits flat 2
- b. Trunk and branches armed. Leaves trifoliolate. Stipels present, gland-like. Flowers red or orange. Fruits strap-like, not flat **2. Erythrina**
- 2a. Exudate red. Fruits winged, rounded. Seed usually 1 **3. Pterocarpus**
- b. Exudate absent. Fruits not winged, linear. Seeds usually more than 1 **1. Dalbergia**

1. DALBERGIA L.f.

Trees, erect or scandent shrubs to woody climbers, sometimes spiny. *Stipules* usually small, caducous, rarely persistent. *Leaves* alternate, imparipinnate. *Leaflets* alternate or subopposite, without stipels. *Inflorescences* axillary or terminal panicles, racemes. *Bracts* and bracteoles small, most often caducous. *Flowers* numerous and small to very small, or relatively large, often fragrant. *Calyx* bell-shaped, 5-lobed, the upper ones broader and shorter than the lower 3, connate at the base, the lowest lobe usually longest. *Corolla* usually white, sometimes purple, green or yellowish, or violet. *Standard* with or without basal callosities, claw well-developed. *Wings* free, elliptic, base truncate, auriculate or hastate. *Keel* petal mostly free, obtuse, only joined at the tip. *Stamens* 9 or 10, monodelphous or diadelphous, sometimes connate near the base only. Anthers globular, small, uniform, erect, didymous, dehiscing with a small transverse slit. *Ovary*

stalked, few-ovuled, style short, curved, glabrous, stigma small, capitate. *Fruits* linear, indehiscent, usually thin and flat, 1–4-seeded, more or less in the centre, not thickened or winged at the sutures. *Seeds* compressed laterally, reniform, radicles bent inwards.

Distribution — Worldwide: c. 300 species. In Africa: c. 41 species. In Asia and Australia: 102 species of which 36 occur in Malesia. In Laos: 15 species. In Vietnam: 27 species.

KEY TO THE SPECIES

- 1a. Leaflets ovate, apex not emarginate 2
- b. Leaflets elliptic, apex emarginate 3
- 2a. Leaflets 7–10. Secondary veins 7–9 pairs. Calyx glabrous. Apex of standard emarginate, callosities absent. Stamens monodelphous. Ovary glabrous. Fruits very flat **1. *D. cochinchinensis***
- b. Leaflets 13–17. Secondary veins 9–12 pairs. Calyx hairy on teeth. Apex of standard rounded, callosities present. Stamens diadelphous. Ovary hairy along the margin. Fruit bulging over seeds **4. *D. oliveri***
- 3a. Calyx hairy on teeth. Standard obovate, callosities absent. Stamen 9, monodelphous. Ovary glabrous. Fruit very flat. Seeds near the margin of fruit . . . **2. *D. cultrata***
- b. Calyx hairy all over. Standard rounded or orbicular, callosities present. Stamens 10, diadelphous. Ovary hairy along the margin. Fruit bulging over the seeds. Seeds in the central part of fruit **3. *D. lanceolaria***

1. *Dalbergia cochinchinensis* Pierre — Fig. 37

Synonym — *Dalbergia cambodiana* Pierre

Vernacular names — K(h)anhung (L); Trác, Sura nam bộ (V).

Large, evergreen trees 15–30 m high, c. 60 cm (sometimes up to 120 cm) diam., profusely branched, crown spherical. *Bark* brownish yellow, longitudinally fissured, sometimes peeled off into fragments. *Stipules* caducous. *Petioles* 2.5–5 cm long, rhachis 6.5–15 cm long, glabrous. *Leaves* 13–25 cm long. *Leaflets* 7–9, coriaceous, ovate to narrowly ovate, 3–8(–10) by 1.8–4(–5) cm, apex obtuse or short acuminate, base obtuse or rounded, glabrous, secondary veins 7–9 pairs, venation reticulate below, petiolules 3–4 mm long. *Inflorescences* axillary or terminal panicles, 10–20 cm long; bracts and bracteoles early caducous; pedicels c. 1 mm long. *Flowers* white or whitish, 5–6 mm long, sweet-smelling, stalk c. 2 mm long, sparsely short hairy. *Calyx* c. 5 mm long, glabrous, sparsely hairy. *Petals* with straight claws; standard obovate or rectangular 4–6 by 2.5–3.5 mm (incl. claw c. 1.5 mm long), callosities and auricles absent, apex obtuse or rounded; wings machete-shaped, or irregular elliptic, 5 by 1.5 mm (incl. claw), apex rounded; keel petals 5–6 by 2 mm (incl. claw), auricles present, apex rounded. *Stamens* 10, 5–6 mm long, monodelphous. *Ovary* stipitate, c. 3 mm long, glabrous, stalk c. 3 mm long, sparsely hairy, with 2–4 ovules, style c. 1.5 mm long, glabrous. *Pods* very flat, narrow, straight, 4–7.5 by 0.8–1.2 cm, both margins parallel at the seed areas, wall thin, glabrous, calyx persistent at the base of the stalk. *Seeds* reniform, 6 by 4 mm in the central part, brown or reddish.




Fig. 37. *Dalbergia cochinchinensis* Pierre (Papilionaceae).

Distribution — Thailand, Cambodia. In Laos: Savannakhet (Ban That) province. In Vietnam: Dac Lac, Lâm Đông, Gia Lai, Kon Tum, Đông Nai, Hồ Chí Minh, and Bà Rịa-Vũng Tàu provinces.

Habitat & Ecology — Shade tolerant when young, later inclining to be light demanding and drought tolerant, growth rate rather slow, coppice-regeneration very strong. In mixed deciduous forest, altitude 50–500 m. The tree grows sparsely in open and

semi-deciduous forest, occasionally in pure stands. Preferring deep sandy clay soil and calcareous soil. Flowering: March to August; fruiting: September to December.

Uses — Used for furniture (beds, wardrobes, desks, etc.), wood turnery, fine-art articles, musical instruments, and sewing-machines.

Specimens examined: *Chevalier* 30096, 40927 (P); *Dournes* s.n. (P); *Hoi* 175 (P); *Lefèvre* 138, 320 (P); *Morange* 313345 (P); *Pierre* 1710 (L, P); *Poilane* 22353 (P); *Schmid* s.n. (P); *Talmy* s.n. (P); *Thorel* 398, s.n. (P).

2. *Dalbergia cultrata* Benth. — Fig. 38

Synonym — *Dalbergia fusca* Pierre

Vernacular names — Lamz (Louang Prabang), Pik nhang (Xieng Khouang) (L); Cẩm lai giao, Trắc giao (V).

Deciduous trees 10–30 m high, with straight trunk, crown open, narrow. *Bark* creamy, quite smooth with horizontal wrinkles, inner bark reddish brown; heartwood dark purple. *Young* twigs, branchlets, and shoots hairy. *Petioles* slender, green, pubescent.




Fig. 38. *Dalbergia cultrata* Benth. (Papilionaceae).

Leaves 10–20 cm long. *Leaflets* alternate or subopposite, 3–6 pairs, narrowly elliptic, 2.5–5 by 1.5–2.5 cm, apex obtuse or rounded, emarginate, base obtuse or rounded, upper surface dark green, glabrous, lower surface glabrescent-pubescent when mature, secondary veins 9–12 pairs, smaller veins easily visible; petiolules 2–3 mm long. *Inflorescences* axillary, panicles, 2–8 cm long. *Bracts* and bracteoles filiform, 1 mm long, persistent, tomentose. *Pedicel* 1–2 mm long, *Flowers* white or pink, 5–6 mm long, appearing before young leaves. *Calyx* tube 3–5 mm long, glabrous, all teeth nearly the same length, sparsely hairy. *Standard* obovate or subreniform, 5 by 3 mm (incl. claw 1.5 mm long), callosities absent, apex slightly emarginated. *Wings* and *keel petals* machete-shaped, 4.5 by 1.5 mm (incl. claw 1.2 mm long). *Stamens* 9, monodelphous. *Ovary* glabrous, 10–12 mm long, stipe long, 3 mm long, ovules 2 or 3. *Fruits* stalked, linear often slightly curved, 2.5–8.5 by 1.5–2 cm, coriaceous, glabrous, brown or greyish when dry, very flat, slightly veined over seeds, stalk 0.5–1.5 cm long. *Seed(s)* 1–3, reniform, near the margin of fruit, 8–10 by 2–6 mm.

Distribution — Myanmar, China (Yunnan), Indochina, Thailand. In Laos: Sayabouri (Pak Lai), Louang Prabang (Phou Khouang), Xieng Khouang (Moung Soui, Phou Kabo, and Moung You), and Savannakhet provinces. In Vietnam: Dac Lac, Lâm Đông, and Đông Nai provinces.

Habitat & Ecology — In mixed forest, usually found in open areas, altitude 100–1500 m. Flowering: January to March; fruiting: March to September.

Uses — Wood reddish black, durable, good quality for precious furniture.

Specimens examined: *Poilane* 11527 (L, P), 12093, 20084 (P), 30948, 32382 (L); *Pierre* 1706 (L); *Thorel* s.n. (P); *Vidal* 1659, 2600 (P).

3. *Dalbergia lanceolaria* L.f.

Vernacular names — Kaboz, Kamphiz (Louang Prabang), Pa dong khao (Vientiane) (L); Trắc mui giáo, Bạt ong (V).

Deciduous trees 15–30 m high, 60–100 cm diameter. *Trunk* straight. *Young branchlets* spreading, ferruginous tomentose, later glabrous. *Bark* smooth, black. *Stipules* ovate, slightly falcate, 12–15 mm long, caducous. *Petioles* 1.5–2.5 cm long. *Leaves* 10–20 cm long, rhachises 7–15 cm long. *Petiolules* 2–3 mm long. *Leaflets* alternate or subopposite, 7–13, rarely 15, lateral leaflets ovate, elliptic, terminal ones elliptic or obovate, 2.5–6.5 by 1.5–3 cm, apex rounded, emarginate, base rounded, obtuse, both surfaces glabrous or glabrescent, secondary veins 9–12 pairs, smaller veins reticulate, both surfaces easily visible, at first densely tomentose, later glabrous. *Inflorescences* axillary or terminal panicle, 5–10 cm long, appearing before leaves. *Peduncles* slender, 2–5 cm long, hairy. *Bracts* and bracteoles very small, caducous. *Flowers* greenish, 9–10 mm long. *Pedicels* 2 mm long, hairy. *Calyx* tube, 4–6 mm long, hairy, upper teeth ovate, 1–1.5 mm long, obtuse, hairy. *Standard* rounded, slightly emarginated at apex, 6–8 by 6–8 mm (incl. claw c. 2 mm long), callosities present at base. *Wings* spoon-shaped, 6 by 4 mm (incl. claw 1.5–2 mm long). *Keel petals* machete-shaped, 4 by 3.5 mm (incl. claw 1.8 mm long). *Stamens* 10, 4–6 mm long, diadelphous (5+5). *Ovary* 3–5 mm long, hairy along the margin and all at base toward stalk, style 2 mm long, glabrous, ovules 3 or 4, stalk 2 mm long, hairy. *Fruit* a pod, 6–11 by 1.5 cm, flat,

bulging over the seeds, fruit-stalk 1–1.5 cm long. *Seed(s)* 1–3, reniform or bean-shaped, 6–7 by 4–5 mm in the central part, dark brown or black.

Distribution — Sri Lanka, India, Myanmar, Indochina, Thailand. In Laos: Louang Prabang (Phou Kouay), Vientiane (Ban Pha Khao), and Savannakhet (Savannakhet – Thakhak) provinces. In Vietnam: Hà Nội, Hà Bac, Dac Lac, and Lâm Đông provinces.

Habitat & Ecology — Usually in mixed deciduous forest, evergreen and open forest, along streams, altitude up to 1500 m. Seedling develop well under the tree canopy, when mature they become light demanding. Natural regeneration medium. Flowering: March to June; fruiting: June to December.

Uses — Wood used for tool-handles, boats, rafters, scantlings, packaging, and other general purposes. Bark contains 14% tannin, used in medicine.

Specimens examined: *Balansa* 4996 (P); *Chevalier* 40299 (P); *Schmid* 747 (P); *Tixier s.n.* (P); *Vidal* 1437, 1775, 2601 (P).

4. *Dalbergia oliveri* Prain — Fig. 39

Synonyms — *Dalbergia bariensis* Pierre; *D. dongnaiensis* Pierre; *D. duperreana* Pierre; *D. mammosa* Pierre

Vernacular names — Kham phi leung (Vientiane), Padong deng (Savannakhet) (L); Cầm lai, Cầm lai bong, Cầm lai mât (South), Trắc lai (Ninh Thuận) (V).

Deciduous trees 15–30 m high, 60–90 cm diam., crown spreading when mature. *Bark* grey, rather thick, scaly, flaking in small pieces, inner bark yellow, heartwood dark red. *Branches* stout, slightly pubescent. *Stipules* early caducous. *Leaves* 15–25 cm long, petioles 3–5 cm long, rachises 10–18 cm long. *Petioliules* 3–4 mm long. *Leaflets* alternate, 13–17, rarely 9–11 or 19–21, ovate, 4–8 by 1.2–3 cm, lateral and terminal leaflets similar in shape and size, apex obtuse or acuminate, often mucronate at the top, base rounded, both surfaces glabrous, secondary veins 9–12 pairs, both surfaces reticulate; young leaves pale pink with silky hairs, mature leaves dark grey-green, glabrous. *Inflorescences* axillary or terminal panicle, 10–45 cm long. *Peduncle* 8–10 cm long, slightly pubescent. *Bracts* early caducous. *Bracteoles* ovate, 1.2 by 0.3 mm, hairy, caducous. *Flowers* purple in bud, pale purple or white, in branched clusters at end of twigs, 10–12 mm long. *Pedicels* 1.5 mm long, slightly pubescent. *Calyx* tube, 4–5.5 mm long, glabrous at the base, teeth hairy, the longest tooth ovate-elliptic, 3 mm long, lateral teeth ovate, 2 by 1.2 mm, upper teeth triangular, 1.2 by 2 mm. *Standard* rounded or orbicular, 6–7 by 6–7 mm (incl. claw 1.5–2 mm long), callosities present, apex rounded. *Wings* spoon-shaped, curved toward apex, 6–7.5 by 3.5–4 mm (incl. claw 1.5–2 mm long). *Keel petals* ear-shaped, 4–6 by 2.5–3.5 mm (incl. claw 1.5–2 mm long). *Stamens* 10, diadelphous (5+5), 6–8 mm long. *Ovary* 3–4 mm long, hairy along the margin at the lower part, ovules 2 or 3, style 2–3 mm long, glabrous, stalk 2–3 mm long, hairy. *Pods* elliptic, 10–12 by 2.5–4.5 cm, flat, bulging over the seeds, fruit-stalk 1–1.5 cm long. *Seed(s)* usually 1, sometimes 2 or 3, reniform or bean-shaped, 12.5 by 9 mm, in the central part, flattened, brown or reddish.

Distribution — Myanmar, Indochina, Thailand. In Laos: Savannakhet and Saravane provinces. In Vietnam: Quang Tri, Dac Lac, Lâm Đông, Phu Yên, Ninh Thuận, Bình Thuận, Đông Nai, and Bà Rịa-Vung Tàu provinces.




Fig. 39. *Dalbergia oliveri* Prain (Papilionaceae).

Habitat & Ecology — Usually in primary and secondary forest, mixed deciduous forest, tropical evergreen or semi-deciduous forest, along streams, altitude up to 1200 m. Usually mixed with *Albizia chinensis*, *Dalbergia cochinchinensis*, *Dipterocarpus alatus*, and *Sindora siamensis*. Flowering: February to June; fruiting: April to December.

Uses — Wood red, hard, with an attractive structure and therefore usually used for furniture and handles of agricultural implements.

Specimens examined: *Chevalier* 36628 (P); *Pierre* 1711 (P); *Poilane* 5632, 5844, 10058, 12059, 13745, 13747 (P); *Vidal* 1745 (P).

2. ERYTHRINA L.

Erythrina fusca Lour. — Fig. 40Synonym — *Erythrina ovalifolia* Roxb.

Vernacular names — Thong lang (L); Vông (V).

Deciduous trees up to 35 m high, 25–60 cm diameter. *Bole* gnarled; bark greyish or pale brown. *Trunk* and branches armed with up to 2 cm long spines, crown rounded. *Stipules* orbicular, c. 3 mm diam., caducous. *Leaves* trifoliolate, 20–30 cm long; petioles 4–10 cm long, very hairy, later glabrous. *Stipellae* present or absent below the terminal leaflet, gland-like, persistent and rounded. *Terminal leaflets* elliptic or ovate,

Fig. 40. *Erythrina fusca* Lour. (Papilionaceae).

10–15 by 5–8 cm, petiolules 3–4 cm long, base broad rounded or obtuse, apex obtuse or almost short acuminate, secondary veins 5–9 per side; lateral ones smaller, 9–11 by 4–5 cm, petiolules 0.4–0.5 cm long. *Inflorescences* lax racemes, 8–20 flowers, 8–30 cm long; peduncle 5–20 cm long, densely pubescent, glabrescent. *Bracts* ovate, puberulous, caducous. *Pedicels* 5–15 mm long, puberulous. *Bracteoles* at apex of pedicels, ovate, 2–3 by 1.5–2.5 mm, outside densely pubescent, inside glabrous. *Flowers* 30 mm long. *Calyx* broadly bell-shaped, 12 by 15 mm, asymmetric bilabiate, upper lip 10–14 mm long, lower lip 12–17 mm long, curved downwards, outside densely pubescent, inside glabrous. *Standard* orange to dark red; claw 9–21 mm long; blade cuneiform, or orbicular to transversely elliptic, 25–30 by 30–40 mm, apex usually emarginate, both sides glabrous. *Wings* creamish green, purplish at apex, claw 1–5 mm long; blade obliquely obovate, 17–35 by 10–20 mm, apex rounded or obtuse, both sides glabrous. *Keel petals* adnate, creamish green, often purple at the base or purplish veined, claw 2–3 mm long; blade ovate to obliquely rhomboid, 13–35 by 7–17 mm, both sides glabrous. *Stamens* 10, diadelphous (1+9), greenish white, half as long as the standard, about 30–40 mm long. *Ovary* 15–25 mm long, densely hairy; style 10–12 mm long, pubescent at the base, upwards glabrous. *Pods* strap-like, black, 10–20 by 1–1.5 cm, stipe 1.5–2 cm long, dehiscent, outside densely pubescent, inside glabrous. *Seeds* usually more than 1, dark brown or black, transversely ellipsoid, 1–18 by 5–8 by 6 mm, hilum 6–8 mm long.

Distribution — South America, India, Myanmar, Indochina, Thailand, throughout Malesia, Pacific Islands. In Laos: Vientiane province. In Vietnam: Lâm Đông, Bình Thuận, Ninh Thuận, and Trà Vinh provinces; planted in many cities in the north.

Habitat & Ecology — From tropical dry to wet through subtropical dry to wet forest, secondary or swamp forest, often along coasts or on river banks, usually close to water. Flowering and fruiting probably throughout the year. In Indochina flowering: November to February; fruiting: January to April.

Uses — Planted as a hedge, as a support tree, and as ornamental shade trees, along roads, around buildings, as living fences of rice-fields and gardens, along pastures, coffee, or cacao plantation. Can be used for green manure and animal feeding. Flowers and young leaves are eaten or used to make tea, and seeds are used in folk medicine as remedies for cancer. Wood is very light and used for canoes and carving.

Specimens examined: *Chevalier* 30309, 30343 (P); *Hiep* 508 (P); *Poilane* 31333 (P); *Thorel* 699, 3005 (P); *Vidal* 1 (P).

3. PTEROCARPUS Jacq.

Small to big semi-deciduous trees up to 40 m high, up to 350 cm diam., usually there are big double leader branches which are rather close to the base, with pronounced low plank buttresses and dense wide-spreading drooping dome-shaped crown. *Bark* greyish brown becoming scaly and fissured with age, inner bark with red sap. *Stipules* small linear to triangular, caducous. *Leaves* compound, alternate, imparipinnate, stipels absent. *Leaflets* entire, alternate, sometimes subopposite or opposite. *Inflorescences* axillary or terminal racemes or panicles; bracts early caducous. *Flowers* big, bright yellow, bisexual, irregular. *Pedicels* articulate at bracteoles insertion. *Bracteoles* small, filiform, or linear to narrowly triangular, early caducous. *Calyx* bell-shaped-tubular,

often slightly incurved at the base, 5-lobed, subequal or unequal, upper 2 teeth usually bigger, lower 3 teeth smaller and more or less deeply or rather widely lobed. *Petals* exerted, with long claws. *Standard* orbicular with a well-developed claw. *Wings* free, more or less obovate to sometimes spatulate. *Keel petals* shorter and smaller. *Stamens* 10, monodelphous or diadelphous (5+5). *Ovary* sessile or stalked, ovules 2–4, hairy; style filiform, glabrous. *Pods* flattened, disc-like, more or less rounded or orbicular, sometimes falcate or plainly asymmetrical, winged, seed-bearing portion corky, hairy or glabrous, indehiscent. *Seed(s)* usually 1, sometimes 3 or 4.

Distribution — In all tropical countries of the world: 60–70 species. Almost pan-tropic, from southern Myanmar through Southeast Asia towards the Santa Cruz and Pacific Islands. Cultivated in Africa, Sri Lanka, India, Taiwan, Okinawa, Hawaii, and Central America. In Laos: 1 species. In Vietnam: 2 species.

KEY TO THE SPECIES

- 1a. Leaflets ovate to broad elliptic, 4–15 by 2–9 cm; apex acuminate, acumen emarginate, short apiculate; secondary veins 8–10 pairs. Inflorescences axillary or terminal panicles. Fruit wing smooth, style point (beak) in upper half **1. *P. indicus***
- b. Leaflets narrowly elliptic, 3–9 by 2–4 cm; apex acute or acuminate, acumen not emarginate, long apiculate; secondary veins 11–17 pairs. Inflorescences axillary racemes. Fruits wing wavy, style point (beak) in lower half **2. *P. macrocarpus***

1. *Pterocarpus indicus* Willd. — Fig. 41

Synonyms — *Lingoum indicum* (Willd.) Kuntze; *L. wallichii* (Wight & Arn.) Pierre; *Pterocarpus pallidus* Blanco; *P. wallichii* Wight & Arn.; *P. zollingeri* Miq.

Vernacular names — Chan daeng, Du bai yai (L); Giáng hương (V), Gioc (Kien Gian), Huỳnh bà rừng (Kiên Giang).

Large, deciduous trees c. 40 m high, up to 350 cm diam., with large and high buttresses, in open areas usually with a wide-spreading to dome-like and large crown, lower branches drooping and sometimes touching the ground. *Bark* greyish brown, shallowly fissured, often thin-scaly and peeling off irregularly. *Branchlets* light brown to dark brown, sparsely to densely lenticellate or sometimes not at all, young parts dark brown to blackish, hairy, glabrescent. *Stipules* linear, 7–15 mm long, hairy on both sides, caducous. *Leaves* 12–25 cm long, petiole 2–4 cm long, usually hairy, rhachis 6–18 cm long, sparsely hairy, glabrescent. *Leaflets* 5–11, ovate to widely elliptic, 4–15 by 2–9 cm; chartaceous to subcoriaceous, base generally rounded or sometimes obtuse to acute, apex usually acuminate, sometimes acute, rarely obtuse, acumen emarginate, short apiculate, surfaces concolorous, greyish brown, sometimes greenish, above slightly shiny, glabrous, beneath slightly dull, sparsely hairy, glabrescent, secondary veins 8–10 pairs; petiolules 3–5 mm long. *Inflorescences* laxly branched, axillary or terminal panicles, 10–20 cm long. *Bracts* and bracteoles filiform, 3–6 mm long, often early caducous. *Flowers* bright yellow. *Pedicels* 6–11 mm long, pubescent. *Calyx* 5–6 mm long, glabrous or sparsely hairy, all the lobes inside sparsely hairy towards the top. *Standard* orbicular, 10–15 mm wide, claw 4.5–6 mm long. *Wings* 10–15 by 5–7 mm, claw 4.5 mm long. *Keel petals* 6–9 by 3–4 mm, claw 4.5 mm long. *Stamens* 10,




Fig. 41. *Pterocarpus indicus* Willd. (Papilionaceae).

diadelphous. *Ovary* densely hairy, style 4–5 mm long, 2 or 3 ovules. *Fruit* orbicular or semi-orbicular, 4–6.6 cm diam., brown to blackish, densely hairy, wing smooth, style point (beak) in upper half, stalk 5–9 mm long, seed-bearing part 1.5–3 cm diam., thickened, 6–9 mm thick, more or less woody. *Seed(s)* 1 or 2, widest at or below the hilum, 2–5 by 8–10 mm; testa dark brown, smooth.

Distribution — India, Myanmar, Indochina, Thailand, farther eastward reaching the Solomons (eastern limit) in the Pacific via Sumatra, West Java, Borneo, Philippines, Sunda Islands, Moluccas, New Guinea, the Pacific (Ryukyu, Carolines). In Laos: unknown. In Vietnam: Kiên Giang, Hồ Chí Minh, and Thừa Thiên-Huê provinces.

Habitat & Ecology — Rain forest or evergreen forest, also in dry areas, in the lowlands, altitude up to 600 m. Seems to prefer a seasonal climate. Flowering: February to May; fruiting: December.

Uses — Wood is used for cabinetry, cart wheels, carving, construction, furniture, and musical instruments. Planted occasionally as shade and ornamental tree. The young leaves and flowers are edible. The flowers are a honey source. The leaves are used as shampoo. The beautiful, termite resistant, rose-scented timber is marketed. The wood gives a reddish dye. Leaves and bark used as anti-emetic, folk remedy for bladder ailments, diarrhoea, dropsy, headache, sores, stones, thrush, and tumours of the abdomen.

Note — Vidal (1959) mentions this species occurring in Laos, although all other literature does not list it.

Specimens examined: *Chevalier* 31397, 31410, 38287 (P); *Edouard* 337, 2709 (P); *Fleury* 30088 (P); *Kerr* 15809 (P); *Put* 2222 (P); *Zimmermann* 39 (P).

2. *Pterocarpus macrocarpus* Kurz

Synonyms — *Lingoum cambodianum* Pierre; *L. macrocarpum* (Kurz) Kuntze; *Pterocarpus cambodianus* (Pierre) Gagnep.; *P. cambodianus* var. *glaucinus* (Pierre) Gagnep.; *P. cambodianus* var. *gracilis* (Pierre) Gagnep.; *P. cambodianus* var. *parviflorus* (Pierre) Gagnep.; *P. macrocarpus* Kurz var. *oblongus* (Pierre) Gagnep.

Vernacular names — Mai du (general), Du luad (Sayabouri) (L); Dáng hương, Sông la (Quang Tri) (V).

Large, deciduous trees c. 30 m high, up to 300 cm diam., with large and high buttresses, in open areas usually with a wide-spreading to dome-shaped crown, large and spreading branches, drooping and sometimes touching the ground. *Bark* pale brown or grey, thickened, slightly fissured, becoming darker or blackish and scaly with age; inner bark fibrous, red-brown with sparse drops of red sap. *Stipules* linear, 7–15 mm long, hairy on both sides, caducous. *Leaves* 8–20 cm long, petiole 3–5 cm long, usually short hairy, rachis 5–15 cm long, densely hairy, glabrescent. *Leaflets* 7–13, alternate or rarely subopposite, narrowly elliptic, 3–9 by 2–4 cm, base rounded or obtuse, apex acute or acuminate-obtuse, acumen not emarginate, long apiculate, young leaves densely hairy, mature leaves bright or dark green and smooth above, usually dull green with scattered brown hairs on veins below, secondary veins 11–17 pairs; petiolules 4–5 mm long, with scattered brown hairs. *Inflorescences* usually axillary racemes, 10–15 cm long. *Bracts* early caducous. *Bracteoles* linear, 2–3 by 1 mm, caducous. *Pedicels* 10 mm long, pubescent. *Flowers* bright yellow. *Calyx* bell-shaped, 5–6 mm long, hairy, all the lobes hairy on both sides towards the top. *Standard* obovate-elliptic, 11–13 by 9–12 mm, claw 4–5 mm long. *Wings* falcate, obovate, auriculate, 10–15 by 5–7 mm, claw 4–5 mm long. *Keel petals* auriculate, 9–10 by 4–5 mm, claw 3–5 mm long. *Stamens* 10, monodelphous or diadelphous, unequal. *Ovary* superior, 3–5 mm long, densely brownish hairy, ovules 2 or 3, style 3–5 mm long, glabrous towards the top, stalk 1–2 mm long, glabrous at the base. *Fruits* orbicular or semi-orbicular, 4–6 cm diam., wing wavy, style point (beak) in lower half, densely with pale brown or whitish hairs when young, later glabrescent, seed-bearing part 1.5–3 cm diam., thickened, 6–10 mm thick, more or less woody, margin between the beak and stalk usually concave, rarely straight, stalk 5–10 mm long. *Seed(s)* 1 or 2, 9 mm long, testa brown, smooth.

Distribution — Myanmar, Indochina, Thailand. In Laos: Sayabouri (Phou Sak, Paklay), Louang Prabang (Phou Khouang), Vientiane (Tha Ngon, Hatxiafong, Ban Khuay Daeng), Bolikhamsai (Borikhane Distr.), Savannakhet, and Saravane provinces.

In Vietnam: Hà Nội, Nghệ An, Quang Tri, Dac Lac, Khanh Hoa, Ninh Hoa, Ninh Thuận, Sông Bé, Tây Ninh, Đông Nai, Hồ Chí Minh, and Kiên Giang provinces.

Habitat & Ecology — Occurring in open semi-deciduous or deciduous Dipterocarp forest, on well-drained, light-textured soils, but with shallow depth and poor in humus. Natural stands especially in dry and hot habitats. Light demanding and drought tolerant tree, often mixed with *Azelia xylocarpa*, *Bambusa* spp., *Dalbergia* spp., *Dendrocalamus* spp., *Lagerstroemia calyculata*, *Peltophorum dasyrhachis*, *Pentacme* spp., *Schima crenata*, *Shorea* spp., *Sterculia* spp., *Terminalia* spp., and *Vitex* species. Flowering: January to May; fruiting: April to December.

Uses — The reddish hard wood has a beautiful appearance, is resistant to termites and insects, classified as a first class timber in Laos and Vietnam. Used in construction and for cabinetry, cart wheels, carving, construction, furniture, and musical instruments. Resin can be used as a red dye. Bark and root used for indigenous medicine, folk remedy for bladder ailments, diarrhoea.

Specimens examined: *Chevalier* 2, 12, 31362, 37742, 39160, 39207, 39938, 40570 (P), 30070 (L, P); *Evrard* 845 (P); *Godefroy* 238 (L); *Hiep* 158, 391, 995 (P); *Kerr* 20754 (P); *Lecomte* 2010 (P); *Maxwell* 99-235 (L); *Muller* 190, 245 (P); *Phung Van Dien* 126 (P); *Pierre* 125, 1817, 6545 (P), 6544 (L, P); *Poilane* 1327, 2433, 4979, 8472, 11775, 12383, 13710, 13715, 13769, 13775, 14133, 16357, 17879, 27988, 30477, 32425 (P), 13339 (L, P); *Soejarto* 10724 (L); *Spire* 864 (P); *Vidal* 1305, 2073, 2355, 2613 (P).

ROSACEAE

Trees, shrubs, herbs, or climbers. *Stipules* on the twig or on the base of the petiole, free or adnate to the petiole, rarely absent. *Leaves* simple or compound, usually alternate or spirally arranged, rarely opposite. *Inflorescences* various. *Flowers* usually bisexual and regular. *Hypanthium* (calyx tube) usually very distinct, from saucer-shaped to tubular or bell-shaped, sepals, petals, and stamens inserted on its rim. *Sepals* usually 5, free, overlapping. *Petals* usually 5, free, overlapping, large or small, showy or not. *Stamens* 5 to many, free, filaments slender, anther 2-locular, splitting longitudinally. *Ovary* superior or inferior, styles free, free or connate at terminal, lateral or basal. *Fruits* various, flesh or dry, splitting or not. *Seed(s)* 1 to many, with or without endosperm.

Distribution — Worldwide: c. 100 genera, c. 3000 species. In Laos: c. 18 genera, c. 44 species. In Vietnam: c. 20 genera, 113 species.

PYRUS L.

Pyrus pashia D. Don — Fig. 42

Synonyms — *Pyrus kumaoni* Decne.; *P. variolosa* Wall.; *P. verruculosa* Bertol.

Vernacular names — Mak chong, Mak chong pa (L); San tan ngai (Lai Châu prov.) (V).

Small to medium deciduous trees or shrubs, with spiny trunk and branches. *Stipules* filiform or linear, caducous. *Petioles* 1–4 cm long. *Leaves* ovate, 5–10 by 2–4 cm, base rounded, margin dentate, apex acuminate, secondary veins 7–10 pairs. *Inflorescences* axillary, umbel-like. *Flowers* white, 6–12. *Pedicel* hairy or glabrous, 2–3 cm long. *Sepals* hairy, lobed, triangular, acute. *Petals* 8 by 6 mm. *Stamens* many. *Ovary* inferior,




Fig. 42. *Pyrus pashia* D. Don (Rosaceae).

3–5 mm long, style 2–5 mm long, glabrous or hairy at the base. *Fruit* globose, 1–2 cm diameter.

Distribution — Nepal, China (Himalaya), Indochina. In Laos: Houa Phan, Xieng Khouang, Saravane, and Champassak (Bolaven plateau) provinces. In Vietnam: Lai Châu, Cao Bang, and Lang Son provinces.

Habitat & Ecology — Occurring on moist sandy, loamy and yellow red clay, near small streams or swamp areas, but also at the lower forest edge near rice paddy fields and open shrub land. Flowering: December to January; fruiting: February to April, and fully ripe from September to December.

Uses — Wood can be used as fuel. Fruits are edible. Young shoots, leaves, and flowers are eaten as vegetables.

Specimens examined: *Balansa* 1535 (P); *Chevalier* 16, 29735 (P); *Eberhardt* 5072 (P); *Mieville* 4, 5, 6, 7, 1921, 40910, 41062 (P); *Pételot* 6166 (P); *Poilane* 1920, 1964, 2042, 2043, 2051, 2128, 2134, 2136, 2200, 15904, 16198, 16262, 19066, 20026, 20066, 25660, 28397, 41142, 41144, 41145, 41146, 41193, 41264 (P); *Spire* 465, 1414 (P); *Vidal* 17, 868, 1477 (P).

SAPINDACEAE

Trees, shrubs, or lianas, rarely climbing herbs. *Bark* usually smooth, inner bark without sap except for *Pometia*. *Leaves* spirally arranged, rarely opposite or whorled, simple, pinnate, digitate, or bipinnate. True stipules usually absent, pseudo-stipules sometimes present. *Leaflets* opposite to alternate, entire or dentate. *Inflorescences* axillary or terminal, spike-like to panicles. *Bracts* and bracteoles present. *Flowers* usually unisexual, rarely bisexual. *Sepals* 4 or 5, rarely more, free to united. *Petals* absent or 2–6, free. *Disc* present. *Stamens* 5–10, rarely more, usually 8, always inserted within the disc, filaments glabrous or hairy, anthers basifixed. *Ovary* superior 1–3(–8)-locular, lobed or not, ovule(s) 1 or 2 per locule. *Fruit* capsules, drupes, sometimes with fleshy sarcotesta or aril, pericarp smooth, muricate or with spines or hairs. *Seeds* completely or partially covered by the sarcotesta or aril.

Distribution — Throughout the tropical and subtropical regions of the world, especially well represented in South America: c. 140 genera, c. 1350 species. In Laos and Vietnam: c. 23 genera, c. 63 species.

SAPINDUS L.

***Sapindus saponaria* L.**

Synonyms — *Sapindus microcarpus* Jard.; *S. mukorossi* Gaertn.; *S. vitiensis* A. Gray

Vernacular names — Kho sak, Sak (L); Bò hòn (V).

Trees up to 25 m high, 60 cm diameter. *Trunk* straight and cylindrical. *Bark* smooth, greenish grey, or greyish brown. Young branchlets terete, yellowish grey hairy, glabrescent, with dense, small lenticels and obcordate leaves scars. *Leaves* usually spirally arranged at the tip of the twigs, paripinnate, up to 40 cm long. *Petioles* terete to 3 angular in cross section, 1.5–6 cm long. *Leaflets* 3–5 pairs, subopposite, elliptic, 6–16 by 3–6 cm, base cuneate, oblique, margin entire, apex obtuse to acute, glabrous on both surfaces, shiny above, lateral veins 8–15 pairs, slightly curved to nearly straight, prominent on both surfaces. *Inflorescences* terminal panicles, up to 30 cm long, pubescent. *Flowers* bisexual, cream-coloured. *Sepals* 5, hairy outside or nearly glabrous, ciliate at the margin. *Petals* 5, greenish white, lanceolate, woolly-ciliate. *Disc* annular, glabrous. *Stamens* 8, free, exerted, filaments hairy at base. *Ovary* ovoid, 3-locular, 1 ovule per locule. *Fruit* a globose drupe, 2–3 cm diam., base often with vestiges of 2 aborted carpels, yellowish when mature. *Seeds* globose, dark brown, 1–1.5 cm diam., hilum 5–6 mm long.

Distribution — India, Myanmar, China, Taiwan, Japan, Korea, Indochina, Thailand, Malesia. In Laos: Houa Phan, Louang Prabang, and Xieng Khouang provinces. In Vietnam: in many provinces, such as Cao Bang, Lang Son, Bac Thai, Ninh Binh, Thanh Hoa, and Nghệ An.

Habitat & Ecology — Primary and secondary forest, altitude below 1000 m. Usually distributed around villages or in secondary forests. Preferring deep, fertile, and wet soils. Fast-growing tree. Natural regeneration is good. Flowering: April, May; fruiting: September to November.

Uses — Sapwood yellowish white, heartwood yellowish grey. Used for making furniture, sawing board, plywood, and boards. The drupes, which lather when combined with water, were once used in varnish, floor wash, and soap. In some areas this species is still preferred for washing hair and delicate clothing, even after the introduction of commercial soaps.

Specimens examined: *Averyanov et al.* VH 1513 (P); *Bon* 1561 (P); *Chevalier* 38577, 40982 (L); *Eberhardt* 3274 (P); *Fleury* 37769, 37902, 38006 (L); *Gilbert* 37832 (L); *Hiep* 5 (L); *Poilane* 2045 (L); *Tsang* 29829 (P).

SAPOTACEAE

Trees or shrubs, monoecious or dioecious, with white, rarely yellow latex. *Bark* usually brown, cracked or fissured, inner bark usually soft, fibrous, pink or red. *Twigs* often scurfy when young. *Stipules* paired, generally triangular or ovate and usually soon falling or absent. *Leaves* simple, spirally arranged and frequently crowded at the end of twigs, or alternate and distichous, margin usually entire, venation pinnate. *Inflorescences* fasciculate, rarely solitary, in axils of leaves or leaf scars. *Flowers* bisexual, rarely unisexual, regular. *Sepals* in two whorls 2–5 in a spiral, free. *Corolla* tubular at base, 4–16-lobed, each lobe entire or divided into 3 segments, white, pale yellow or pale green. *Stamens* either of the same number as corolla lobes and opposite them or numerous and arranged in 1–3 whorls, inserted at the base or at the throat or corolla tube; staminodes if present alternating with stamens or fixed in the sinuses of corolla lobes. *Ovary* superior, sessile, small, ovoid, 3–18-celled, each cell with 1 ovule attached to the axis, style simple, stigma small and indistinct, simple or minutely lobed. *Fruit* a fresh berry, rarely a drupe or capsule, with persistent sepals at the base and a persistent style at the apex. *Seed(s)* 1–6, ovoid to ellipsoid, glossy, brown to black outside, hard, scar usually white, large, adaxial, basiventral or basal, elongate, endosperm present or absent.

Distribution — Throughout tropical and subtropical regions: c. 53 genera, 1100 species. In Laos: c. 5 genera, 8 species. In Vietnam: c. 13 genera, c. 36 species.

KEY TO THE GENERA

- 1a. Bark not fissured into squares, usually with many greyish yellow dots. Leaves alternate, lateral veins slightly curved near the margin. Calyx 5-lobed. Corolla 5-lobed. Fruit a capsule **1. Eberhardtia**
 b. Bark usually fissured into squares. Leaves spirally arranged, crowded at the end of twigs, lateral veins diminishing and becoming inconspicuous towards leaf margin. Sepals 4, in two pairs. Corolla 8–12-lobed. Fruit a berry **2. Madhuca**

1. EBERHARDTIA Lecomte

Eberhardtia tonkinensis Lecomte — Fig. 43

Vernacular names — Mắc niêng, Cồng sưa bắc bộ (V).

Trees up to 20 m high, up to 60 cm diameter. *Trunk* straight, cylindrical. *Bark* greyish brown, with many greyish yellow dots, inner bark pink with milky sap. *Twigs* rusty



Fig. 43. *Eberhardia tonkinensis* Lecomte (Sapotaceae).

tomentose. *Stipules* 2, triangular. *Leaves* simple, alternate, elliptic, 10–30 by 3–8 cm, base cuneate, apex with a short point or acuminate, glabrous, green above, densely covered with silvery brown or rusty pubescence beneath, venation pinnate, the lateral veins slightly curved near the margin, prominent beneath. *Petioles* 1.5–3.5 cm long. *Flowers* in axillary clusters. *Calyx* 5-lobed, brownish hairy outside, glabrous inside. *Corolla* tubular, 5-lobed. *Stamens* 5, inserted on the throat of the corolla. *Ovary* ovoid, 5-celled, glabrous. *Fruit* a capsule, ovoid, 3–4 by 2.5–3.5 cm, dehiscent. *Seed* 2–3 cm long, brown and shiny, glabrous, scar adaxial, oblong.

Distribution — Indochina. In Laos: Xieng Khouang province. In Vietnam: Lào Cai, Yên Bái, Lai Châu, Hà Giang, Tuyên Quang, Hoà Bình, Quang Ninh, Vinh Phú, Thanh Hoa, Ninh Bình, Nghệ An, and Hà Tĩnh provinces.

Habitat & Ecology — Primary and secondary forest, altitude below 1000 m, but mainly 500–800 m. Usually mixed with *Canarium album*, *Castanopsis* spp., *Garcinia oblongifolia*, *Girroniera subaequalis*, *Quercus* spp., and *Schima wallichii*. Fast growing and light demanding tree, but shade tolerant when young. Natural and coppice regeneration are good under a forest cover of 40–60%. Flowering: April, May; fruiting: August, September.

Uses — Wood yellowish white, grain fine, resistant to termites. Wood used for construction and making furniture. The oil from the seeds is used for cooking.

Note — Dung (1996) recorded *Eberhardtia tonkinensis* to be endemic in the north of Vietnam. Examining material in Paris, however, revealed that Kerr also collected this species in Xieng Khouang province, Laos.

Specimens examined: *Hance* 5027 (P); *Kerr* 21148 (P); *Pételot s.n.* (P); *Poilane* 2025, 13125, 18807, 18809 (P).

2. MADHUCA J.F. Gmel.

Madhuca pasquieri (Dubard) H.J. Lam

Synonyms — *Bassia pasquieri* Lecomte; *Dasillipe pasquieri* Dubard

Vernacular names — Sên, Sên mật (V).

Large trees up to 40 m high, up to 120 cm diameter. *Trunk* straight, cylindrical. *Bark* blackish brown, fissured into squares, inner bark pink with milky sap. *Stipules* caducous. *Leaves* simple, spirally crowded at the end of twigs, obovate, 6–16 by 3–6 cm, base cuneate, apex with a short point or acuminate, glabrous, green above, pale green beneath, venation pinnate, the lateral veins diminishing and becoming inconspicuous towards leaf margin. Young leaves pink to red. *Petioles* 1–3 cm long. *Flowers* solitary or in axillary clusters, bisexual, yellowish white. *Sepals* 4, in two pairs, outer thick, inner thinner, densely hairy. *Corolla* tubular 8–12-lobed, imbricate. *Stamens* 12–22, in two whorls; staminodes absent. *Ovary* ovoid, densely hairy, 6–8-locular. *Fruit* a berry, ovoid or subglobose, 2.5–3 cm long. *Seeds* elliptic, 2–2.4 cm long, scar adaxial, elliptic.

Distribution — Endemic to Vietnam, in many provinces in the north, such as Lào Cai, Yên Bái, Lai Châu, Hà Giang, Tuyên Quang, Hoà Bình, Hà Bắc, Bắc Giang, Quảng Ninh, Vinh Phú, Thanh Hoá, Ninh Bình, Nghệ An, and Hà Tĩnh.

Habitat & Ecology — Primary and secondary forest, altitude below 1000 m. Usually mixed with *Castanopsis indica*, *Erythrophloeum fordii*, *Garcinia oblongifolia*, *Girroniera subaequalis*, *Hopea mollissima*, and *Vatica odorata*. Slow growing and light demanding tree, but shade tolerant when young. Prefers deep and moist soils. Natural and coppice regeneration are good under a forest cover of 50–60%. Flowering: August to October; fruiting: December to February.

Uses — Wood pinkish red, hard, heavy, grain fine, resistant to termites. Used for construction, railway sleepers, furniture, and ship-building. The oil from the seeds is used for lighting, cooking, and in the chemical industry. Leaves used to treat burns.

Specimens examined: *Balansa* 4336 (L); *Kerr* 21148 (P); *Pételot s.n.* (P); *Poilane* 2025, 13125, 18807, 18809 (P).

SONNERATIACEAE

Evergreen or deciduous trees. *Stipules* present or absent. *Leaves* simple, opposite, biseriate, entire, glabrous, coriaceous. *Flowers* large, showy, bisexual, actinomorphic, 1–3 or more in terminal corymbs, pedicelled, rather large. *Calyx* 4–8, connate into a tube, leathery. *Petals* absent or as many as calyx-segments, in the latter case broad and wrinkled or very narrow and smooth. *Stamens* many, inserted on the calyx-tube, inflexed in bud, filaments filiform-subulate, anthers reniform, 2-celled. *Ovary* superior, sessile with a broad base, up to 10- or more-celled, ovules numerous on thick axil placentas, style 1, robust, stigma 1, head-like, entire or slightly lobed. *Fruit* resting on the calyx-tube, an indehiscent berry or a valvate capsule. *Seeds* many, small, embedded in firm pulp or not.

Distribution — Tropical or subtropical regions: 2 genera, c. 12 species. In Laos: 1 genus, 1 species. In Vietnam: 2 genera, 4 species.

DUABANGA Buch.-Ham.

Duabanga grandiflora (DC.) Walpert — Fig. 44

Synonyms — *Duabanga sonneratioides* Buch.-Ham.; *Lagerstroemia grandiflora* DC.

Vernacular names — Lin ngo (Khammouane prov.), Phay (Xieng Khouang prov.), Ten (Laung Prabang prov.) (L); Phay sùng, Phay, Bảng lảng bản (V).

Big trees up to 40 m high, up to 200 cm diameter. Scarcely buttressed, glabrous in all parts. Young trees have whorls of horizontal branches, but in old trees the branches usually drooping, sometimes horizontal, young branchlets quadrangular. *Bark* pale grey, pimply, not cracked, irregularly flaking in older trees. *Stipules* very small. *Leaves* simple, opposite, ovate to ovate-elliptic, 10–27 by 4–10 cm, base rounded or slightly heart-shaped, margin entire, apex slightly acuminate, upper surface dark green, lower surface grey-green, completely smooth even when young, secondary veins 12–18 pairs, parallel, arched and joined at margin. *Petioles* 2–9 mm long, stout. *Flowers* nocturnal, white, showy, with offensive smell, 4–15 cm across, in heavy branched clusters (corymbs) of 20–30 flowers at the end of twigs. *Calyx tube* pale green, broadly cup-shaped, 1.2–3 cm across, calyx lobes 4–7, fleshy, triangular-ovate, 1.2–3 by 0.7–1.4 cm, spreading. *Petals* 6 or 7, obovate, 4 by 3 cm, apex rounded, very narrow at base, wrinkled and very fragile, falling in the early morning. *Stamens* many, slender, much longer than petals, arranged in two rows or series, anthers curved, very mobile. *Ovary* broadly conical, style 1, slender, stigma lobed, dark green. *Fruit* ovoid-globose, 2.7–4.5 by 3–4 cm, with large star-shaped calyx at the base, green turning brown.

Distribution — India, Myanmar, South China, Indochina, Thailand, Peninsular Malaysia. In Laos: Louang Prabang, Xieng Khouang, Champassak, and Attopeu provinces. In Vietnam: Đông Nai, Nghê An, Lào Cai, Thừa Thiên-Huê, Quang Tri, Kiên Giang, and Lâm Đông provinces.

Habitat & Ecology — Deciduous tree or partially deciduous for a brief period. Light demanding and fast-growing, especially at mountain-foots and along streams and water-ways. Suitable for a cool climate and deep or stony soil. Often in disturbed moist areas, altitude up to 1200 m, more frequent in hilly areas on the moist valley slopes. Flowering: January to June; fruiting: March to November.




Fig. 44. *Duabanga grandiflora* (DC.) Walpert (Sonneratiaceae).

Uses — Wood used for tea-boxes and for house and boat-building, planks, canoes, oars, and umbrellas. Fruits and leaves boiled for traditional black dye. Fruits edible, used boiled as vegetable.

Specimens examined: *Balansa* 3864 (P); *Chevalier* 31325, 39963 (P); *Eberhardt* 2876 (P); *Evrard* s.n. (P); *Harmand* 50 (P); *Hayata* 662 (P); *Hiep* 362 (P); *Jean* 38399 (P); *Pételot* 3942 (P); *Poilane* 1266, 20130, 24766, 25503 (P); *Pierre* 536 (P); *Schmid* s.n. (P); *Spire* 1162 (P); *Thorel* 2688 (P).

STERCULIACEAE

Trees or shrubs, rarely herbs, with stellate hairs or scales. Trees usually with prominent plank buttresses. Inner bark usually prominently radially streaked, or with wedges, generally tough due to long fibres. *Leaves* simple, palmate or digitately compound, alternate or spirally arranged. *Stipules* present. *Petioles* often long and kneeed. *Inflorescences* various, usually axillary and compound, sometimes flowers solitary. *Flowers* unisexual or bisexual, regular, rarely irregular. *Sepals* 3–5, valvate, partly connate or rarely free. *Petals* 5 or absent, contorted or imbricate, free or adnate at the base to the staminal tube, often persistent. *Stamens* 5, 10, 15 or more, free or connate into a narrow tube, and sometimes with staminodes, anthers 2-locular. *Ovary* superior, carpels 2–5, rarely 1 or 10–12, united or free, ovule(s) 1 or 2 or more in each carpel, inserted at the inner angle, style simple or divided into lobes, rarely free to the base. *Fruit* dehiscent or indehiscent, winged or not, fleshy to leathery or woody, often separating as mericarps. *Seeds* sometimes arillate, winged or not.

Distribution — Widespread in the tropics, subtropics, and rarely in temperate regions: c. 60 genera, c. 700 species. In Laos and Vietnam: c. 17 genera, c. 74 species.

PTEROSPERMUM Schreb.

KEY TO THE SPECIES

- 1a. Young leaves palmate, mature leaves elliptic, base oblique, apex pointed. Stipules linear. Fruit a terete, not ridged capsule **1. *P. heterophyllum***
 b. Young leaves reniform, mature leaves obovate, truncate or heart-shaped at base, apex 3–5-lobed. Stipules divided. Fruit a capsule, with 5 straight longitudinal ridges **2. *P. truncatolobatum***

1. *Pterospermum heterophyllum* Hance

Vernacular names — Ham ao (L); Mang xanh, Long mang, Long mang thuong (V).

Trees up to 20 m high, up to 60 cm diameter. *Bark* greyish brown, inner bark pink, soft and fibrous. *Leaves* and young twigs densely covered with brownish yellow stellate hairs. *Stipules* linear, 8–14 mm long. *Leaves* simple, alternate. Young leaves palmate, 17–30 cm diameter. *Petioles* 20–30 cm long. Mature leaves elliptic 6–16 by 3–7 cm, base oblique, margin entire, apex pointed, upper surface yellow brown, lower surface covered with brownish yellow stellate hairs. *Petioles* 1–2.5 cm long. *Inflorescences* axillary, solitary, or in clusters of 2–4 flowers. *Sepals* 5, free, linear, covered with densely stellate hairs outside, whitish brown and long hairs inside. *Petals* 5, strap-shaped, glabrous. *Stamens* 15, connate into 5 bundles, staminodes 5, linear, alternating with stamen bundles. *Ovary* ovoid, densely tomentose, 5-locular, ovules numerous. *Style* slender, hairy. *Fruit* a capsule, narrow elliptic, woody, 4–6 by 2.5–3.5 cm, dark brown tomentose, splitting into 5 pieces when mature. *Seeds* winged, 3–4 cm long.

Distribution — India, China, Indochina, Thailand. In Vietnam: in most provinces in North and Central. In Laos: not yet recorded but presumably occurring.

Habitat & Ecology — Primary and secondary forest, altitude below 700 m. Usually confined to mountainous and limestone regions. Light demanding tree, prefers soils

with medium humidity. Coppice regeneration is good. Flowering: April, May; fruiting: February, March next year.

Uses — The wood is reddish brown, used for making furniture, flooring, plywood, and is suitable for the production of pulp. Leaves and bark are rich in tannins; they are sometimes applied in traditional medicine, such as a poultice against itch and to treat wounds.

Specimens examined: *Cavalerie* 3483 (P); *How* 70856 (P); *Tsiang* 695 (P); *Wang* 695, 34757, 52198 (P).

2. *Pterospermum truncatolobatum* Gagnep.

Vernacular names — Ham ao (L); Mang cụt, Lòng mang cụt (V).

Trees up to 25 m high, up to 80 cm diameter. *Bark* greyish brown, inner bark 1–1.5 cm thick, greyish yellow, with long fibres. *Stipules* 6–10 mm long, divided. *Leaves* and young twigs densely covered with brownish yellow stellate hairs. *Young leaves* reniform, 15–30 cm diameter. *Petioles* 10–20 cm long. *Mature leaves* obovate, 8–17 by 4–8 cm, truncate or heart-shaped at base, margin entire, apex with 3–5 lobes, upper surface yellow brown, lower surface covered with brownish yellow stellate hairs. *Petioles* 8–15 mm long. *Flower* bisexual, axillary, subsessile. *Sepals* 5, free, linear, covered with densely stellate hairs outside, white and long hairs inside. *Petals* 5, bell-shaped, glabrous. *Stamens* 15, connate into 5 bundles, staminodes 5, linear, alternating with the stamen bundles. *Ovary* ovoid, densely tomentose, 5-locular, ovules numerous, style slender, hairy. *Fruit* a capsule, oblong, woody, 4–8 by 2.5–4 cm, with 5 straight longitudinal ridges, dark brown tomentose, splitting into 5 pieces when mature. *Seeds* winged, 3–4 cm long.

Distribution — Confined to Indochina. In Laos: this species is found in Vientiane province. In Vietnam this species is common in many provinces such as Lào Cai, Yên Bái, Tuyên Quang, Hà Giang, Hoà Bình, Ninh Bình, Hà Tây, Thanh Hoa, Nghệ An, Hà Tĩnh, Quang Bình, Đà Nẵng, Kom Tum, and Gia Lai.

Habitat & Ecology — Secondary forest, altitude below 700 m. Light demanding tree. Coppice regeneration is good. Flowering: April to June; fruiting: January to March.

Uses — Wood is reddish brown. Used for making furniture, flooring, plywood, boxes, and is suitable for the production of pulp. Leaves and bark rich in tannins, sometimes applied in traditional medicine, such as a poultice against itch and to treat wounds.

Specimens examined: *Balansa* 1305 (P); *Bon* 4862 (P); *Cavalerie* 3483 (P); *Poilane* 11104 (P).

THEACEAE

Small to large evergreen trees. *Bark* fibrous. *Stipules* absent. *Leaves* simple, alternate or spirally arranged, rarely subopposite, more or less coriaceous, margin entire or serrate. *Flowers* regular, usually bisexual, rarely unisexual, single, rarely fascicles, axillary or extra-axillary, rarely terminal, mostly 5-merous, often showy. *Bracteoles* 2 or 3 at the base of calyx. *Sepals* usually 5, free or slightly connate, overlapping, often unequal. *Petals* usually 5, free or connate at the base, overlapping. *Stamens* numerous adnate to petals, anthers attached to filament at the base (basifixed) or attached to filament at the apex (versatile), opening longitudinal. *Ovary* superior, rarely half inferior, lacking

stalk, usually 2–5-locular, ovules 2 to many in each locule, rarely 1, style(s) 1–5, free or partly united; stigma usually small, head-like (capitate) or entire. *Fruit* a capsule or berry with persistent sepals at the base and style at apex. *Seeds* small, few or many, sometimes winged (*Schima*).

Distribution — Tropical or subtropical regions, mainly in America and Asia, a few in Africa: c. 16 genera, c. 600 species. In Laos and Vietnam: c. 9 genera, 98 species.

SCHIMA Reinw. ex Blume

Schima wallichii (DC.) Korth. — Fig. 45

Synonyms — *Gordonia wallichii* DC.; *Schima crenata* Korth.; *S. noronhae* Reinw. ex Blume

Vernacular names — Mi (Xieng Khouang), Khai so, Thalo in other provinces (L); Vối thuốc (V).

Trees up to 30 m high, up to 70 cm diameter. Buttresses, if present, up to 180 cm high. *Bark* dark grey to black, scaly or cracked in small thick pieces, inner bark pink or dark red, with dark spots, containing fibres which can cause skin irritation. *Sapwood* white.



Fig. 45. *Schima wallichii* (DC.) Korth. (Theaceae).

Young leaves orange to red brown and densely covered with silky hairs. *Leaves* spirally arranged, broadly elliptic, 6–20 by 3–8 cm, base cuneate, apex acute to acuminate, margin more or less toothed, secondary veins 6–8 pairs. *Petioles* 1.5–3 cm long with fine short hairs (slightly pubescent). *Flowers* axillary, solitary or clustered in the axils of leaves near or at the end of the twigs, 17–35 mm diameter. Pedicel 1–2.5 cm long. Bracteoles 2. *Sepals* 5, more or less rounded, glabrous. *Petals* 5, shortly connate at the base, obovate, up to 2 cm long, white, aromatic. *Stamens* many, unequal, filaments glabrous, anthers elliptic. *Ovary* 5-locular, with 2 or 3 ovules in each locule. *Fruits* a woody capsule, almost globose, 1–3 cm diam., silky, opening by 5 valves, stalk 1.3–2 cm long. *Seeds* kidney-shaped, 8 mm long, winged.

Distribution — India, Nepal, Myanmar, South China, Indochina, Thailand, Peninsular Malaysia, Indonesia (West Java, Borneo), Philippines (Palawan). In Laos: Louang Prabang, Houa Phan, Xieng Khouang, Savannakhet, and Saravane provinces. In Vietnam: Lào Cai, Cao Bang, Lai Châu, Ninh Thuận, Kon Tum, and Hà Bac provinces.

Habitat & Ecology — Light demanding when mature, shade tolerant when young, cold-tolerant tree. The trees often grow in primary and secondary forest, well-drained land, altitude up to 3000 m, but most abundant in regions above 400 m. Flowering: March, April; fruits: January to March.

Uses — Wood red-brown, hard, used for timber, house construction, furniture, boats, plywood, agricultural implements. The leaves are used for fodder.

Note — This species is very variable, our material was mainly formerly identified as *S. wallichii* subsp. *noronhae* (leaf mostly quite entire) and some specimens as subsp. *oblata* (leaf slightly crenate, dentate). However, the great overlap in many characters keeps us from accepting subspecies.

Specimens examined: *Averyanov et al. VH 1389* (P); *Chevalier 29576* (P); *Dussaud 4* (P); *Kerr 21116* (P); *Lau 3677* (P); *Mekee 6049* (P); *Pierre 1413* (L); *Pételot 4458* (P); *Poilane 2009, 2339, 5920, 7728, 13725, 18813, 19059, 20035, 20363, 26813, 27092* (P); *Simond 6049* (P); *Thao Kham 4458* (P); *Thorel 2768* (P).

THYMELAEACEAE

Trees, shrubs, or climbers, with a tough and fibrous ‘silvery’ inner bark. *Stipules* absent. *Leaves* simple, opposite, decussate, spirally arranged or alternate, entire. *Petioles* very short. *Inflorescences* terminal, axillary, sometimes on sessile or peduncled, racemose, umbelliform, spicate, capitata, or fascicled. *Flowers* bisexual, rarely unisexual, regular. *Bracts* present, usually fragrant. *Calyx* tubular, often brightly coloured, shallowly or deeply 4- or 5-lobed, imbricate. *Petals* absent or 4–12 scales, inserted on the throat of calyx tube or slightly lower. *Stamens* 2 or many, usually as many as twice the calyx lobes and opposite them. *Disc* scale-like, annular, or cup-shaped, sometimes absent. *Ovary* superior, 1- or 2- or (3–5- in *Gonystylus*)locular, 1 ovule in each locule, style filiform, caducous, sometimes very short or obscure, stigma globose, pyramidal, sometimes papillose. *Fruit* a drupe, a berry, or a capsule. *Seeds* with caruncle or aril.

Distribution — Throughout the tropical and subtropical regions of the world, strongly developed in the drier parts of Africa and Australia: c. 50 genera, 500 species. In Laos: c. 4 genera, c. 4 species. In Vietnam: c. 7 genera, c. 16 species.

AQUILARIA Lam.

Aquilaria crassna Pierre ex Lecomte — Fig. 46

Synonym — *Aquilaria agallocha* Roxb.

Vernacular names — Ketsana (L); Trầm, Trầm hương, Dó, Dó bầu (V).

Trees up to 30 m high, up to 100 cm diameter. Branches drooping. *Bark* brownish grey, shallowly fissured and flaking in thin strips, inner bark pale yellow with patches of fragrant, dark coloured resin in old trees. *Leaves* simple, alternate, spirally arranged, coriaceous, elliptic, 5–11 by 3–6 cm, base broadly cuneate, margin entire but often wavy, apex acuminate or acute, dark or purple-green above, shining, light green beneath, glabrous, veins 15–20 pairs, more conspicuous beneath, veins and veinlets slender. *Petioles* 3–7 mm, pubescent. *Inflorescences* umbellate, axillary or terminal. *Flowers* fragrant, yellowish green. *Pedicels* 5–6 mm, densely yellow-grey-pubescent. *Calyx tube* narrowly bell-shaped, 5–6 mm, densely pubescent on both sides, lobes 5, ovate. *Petals* 10, scale-like, inserted on the throat of the calyx tube, densely pubescent. *Stamens* 10, in 2 rows, filaments c. 1 mm long, anthers oblong. *Ovary* ovoid, densely with greyish white hairs, 2-locular, style absent or very short, stigma capitate. *Fruit* a




Fig. 46. *Aquilaria crassna* Pierre ex Lecomte (Thymelaeaceae).

capsule, ovoid, green and silky hairy when young, 2–4 by 2–3 cm, calyx persistent at base. *Seed(s)* 1, sometimes 2, ovoid, brown, c. 10 by 5 mm.

Distribution — India, Bangladesh, Myanmar, China, Indochina, Thailand, Malaysia. In Laos: in most provinces, such as Houa Phan, Xieng Khouang, Vientiane, Savannakhet, Champassak, and Attapeu. In Vietnam: in many provinces, such as Lang Son, Son La, Lào Cai, Yên Bái, Lai Châu, Tuyên Quang, Hà Giang, Bình Ninh, Hoà Bình, Thanh Hoa, Nghệ An, Hà Tĩnh, Quang Bình, Đắc Lạc, and Kiên Giang.

Habitat & Ecology — Primary and secondary forest, altitude up to 1000 m. This species is well adapted to various habitats, including rocky, sandy, or well-drained slopes, ridges and land near swamps. Light-demanding tree, but tolerant to shade when young. Flowering: April to June; fruiting: July to September.

Uses — Wood soft, used for general furniture, cosmetic preparation, clothing articles, and musical instruments. Bark used for hammock making and paper pulp. Resin used as traditional medical to treat asthma, chest congestion, colic, diarrhoea, diuretic, kidney problems, nausea, thyroid cancer, and lung tumours.

Note — Nowadays this species is very rare, because the resin of the infected and cut trees is much sought after and yields a very high price.

Specimens examined: *Fourdan* 39112 (P); *Kerr* 10043 (L); *Pierre* 1443, 1449, 3619 (P); *Poilane* 1111, 5411, 6680 (P), 6400 (L, P).

TILIACEAE

Trees, shrubs, or herbs, rarely climbers, with stellate hairs or scales. *Bark* fibrous. *Stipules* early caducous. *Leaves* simple, alternate or spirally arranged, lobed or not, base 3–7-veined. *Inflorescences* cymes, racemes or panicles, solitary. *Flowers* regular, 5-merous, bisexual, rarely unisexual. *Sepals* 5, free or united. *Petals* absent, or present and free, alternate with the sepals. *Stamens* many, free or shortly connected at the base or in 5–10 bundles with slender filaments arising from a thickened ring around the ovary, anthers 2-locular, opening by a slit lengthwise or by an apical pore. *Ovary* superior, sessile or stalked, 2–10-locular, ovule 2 to many per locule, style awl-shaped or branched, stigma lobed or pointed. *Fruit* dry or fleshy, splitting or not, winged or not. *Seed(s)* 1 to many in each locule, endosperm present, wings absent.

Distribution — Widely distributed in the tropical and temperate zones: 55 genera, c. 1130 species. In Laos and Vietnam: c. 18 genera, 60 species.

BERRYA Roxb.

Berrya mollis Wall. ex Kurz

Synonym — *Berrya ammonilla* Roxb. var. *mollis* Mast.

Vernacular names — Lieng, Lieng khai (L); Gia ti (V).

Deciduous trees up to 20 m high, up to 25 cm diameter. *Bark* pale brown or grey, slightly cracked, inner bark pale yellow. *Leaves* broadly ovate or circular, 6–15 by 7–18 cm, base heart-shaped, margin entire or irregularly wavy, apex shortly cuspidate or acute,

mature leaves softly hairy on lower surface, secondary veins 3–7 pairs, venation ladder-like, distinct on lower surface. *Petioles* 2–8 cm long, hairy. *Inflorescences* terminal or axillary, 5–20 cm long. *Flowers* globose in bud, c. 5 mm diam., white. *Sepals* sometime 3-merous, ovate. *Petals* slightly spoon-shaped, rather longer than sepals, 4–5 by 1–2 mm. *Stamens* many, smooth, anthers attached at base. *Ovary* globose, 3-locular, hairy, style slender, stigma pointed. *Fruit* a globose capsule, hairy, 3.5–5.5 by 1 cm, splitting in 3 sections, each with a pair of wings, thinly hairy, pinkish brown when ripe.

Distribution — India, Myanmar, Indochina, Thailand. In Laos: Sayaboury, Vientiane, and Savannakhet provinces. In Vietnam: Đông Nai province.

Habitat & Ecology — In moist upper mixed deciduous forest, altitude 100–1000 m. Flowering and fruiting: April to September.

Uses — Wood used in construction of rims, beams, or cartwheels.

Specimens examined: *Beaud 739* (P); *Maxwell 91-898* (P); *Pierre 1492* (P); *Poilane 28014* (P); *Sangkachand 1661* (P); *Spire 188* (P); *Thorel 2111* (P); *Tixier s.n.* (P).

VERBENACEAE

Trees, shrubs, or lianas. *Twigs* often lenticellate, young twigs quadrangular. *Stipules* absent. *Leaves* generally opposite, rarely whorled, simple, palmate, trifoliolate or pinnate. *Inflorescences* in clusters, panicles, cymes, racemes or spikes, axillary or terminal. *Flowers* small or large, bisexual, general bilateral. *Calyx* usually tubular with 4 or 5 lobes or teeth. *Corolla* usually 2-lipped, more or less funnel- or trumpet-shaped with a short or long tube, lobes 4 or 5. *Stamens* 5, 4 or 2, inserted in the corolla tube, projecting out of flower or not, anthers 2-celled. *Ovary* superior, 2–4-celled, often 4-lobed, ovule(s) 1 or 2 in each cell, style one, stigma entire or bifid. *Fruit* a drupe or capsule, seated on or enclosed in the usually enlarged persistent calyx. *Seed(s)* 1–4, small.

Distribution — Widely distributed in tropical and subtropical regions: c. 76 genera, 3000 species in the world. In Vietnam: c. 8 genera, c. 22 species.

KEY TO THE GENERA

- 1a. Petiole terete, not winged. Flower usually large, corolla with 5 unequal lobes, 2 upper lobes and 3 lower lobes, stamens 4, 1 pair longer than the other. Fruit with persistent calyx at base **1. *Gmelina***
- b. Petiole usually narrowly winged. Flower usually small, corolla with 5 or 6 equal lobes, stamens 5 or 6, equal. Fruit enclosed by calyx **2. *Tectona***

1. GMELINA L.

Gmelina arborea Roxb.

Synonyms — *Gmelina rheedii* Hook.; *Premna arborea* (Roxb.) Roth

Vernacular names — So (L); Lòi thọ (V).

Large trees up to 30 m high, up to 100 cm diameter. *Bole* straight, cylindrical. *Bark* smooth, greyish brown, with many black lenticels, inner bark cream-coloured. *Young*

twigs quadrangular, covered with yellow hairs becoming fulvous. *Leaves* simple, opposite, broadly ovate, 10–19 by 7–14 cm, base cuneate or slightly heart-shaped, with 2–6 rounded glands near petiole, margin entire, apex pointed, young leaves densely covered with yellow stellate hairs, mature leaves glabrous or with scattered hairs below, lateral veins 4–7 pairs per side. *Petioles* 4–11 cm long, slender, with two glands at the top of the petiole and 2–6 glands at the base of the leaf blade. *Inflorescences* panicle, axillary or terminal, up to 40 cm long, hairy. *Flowers* bisexual, large. *Calyx* cup-shaped, with 4 or 5 short teeth, densely brown hairy outside. *Corolla* funnel-shaped, creamy-violet or creamy-pink, with 5 unequal lobes, upper 2 lobes and lower 3 lobes with the middle lobe much larger than the side ones, usually hairy outside. *Stamens* 4, 1 pair longer than the other, inserted on corolla tube. *Ovary* smooth, styles 2, stigma divided. *Fruit* a drupe, ovoid, 1.5–2.5 cm long, with persistent calyx at the base. *Seed(s)* 1 or 2.

Distribution — Pakistan, Sri Lanka, India, Myanmar, China, Indochina, Malaysia, and planted in many Asian countries, Africa, Brazil. In Laos: Louang Namtha, Saysabouri, and Attopeu provinces. In Vietnam: Tuyên Quang, Hà Giang, Lang Son, Lai Châu, Son La, Vinh Phu, Quang Ninh, Bac Thai, and Ninh Binh provinces.

Habitat & Ecology — Primary and secondary tropical forest, or along streams or river banks. Usually mixed with *Canarium album*, *Castanopsis fissa*, *Castanopsis indica*, *Endospermum sinensis*, and *Engelhardtia chrysolepis*. Fast-growing and light demanding tree. Flowering: April to June; fruiting: October to December.

Uses — Wood whitish grey, fine grain. Mainly used for light construction, pulping, sawing boards, making furniture, musical instruments, matches, and packing boxes. Fruits and bark have medicine properties against bilious fever. This species is also used as ornamental species in landscaping.

Specimens examined: *D'Alleizette s.n.* (L); *Glutron* 37988 (P); *Harmand* 187, 1377 (P); *Maxwell* 75-318, 76-58, 88-218, 88-431, 88-981, 90-252, 94-294, 94-398 (L); *Pételot* 4536 (P); *Pierre* 1901 (P); *Poilane* 13635 (P).

2. TECTONA L.f.

Tectona grandis L.f.

Synonym — *Tectona theka* Lour.

Vernacular names — Sak (L); Téché (V).

Large trees up to 30 m high, 150 cm diameter. *Bole* straight, cylindrical, usually canalculated and buttressed at the base. *Bark* greyish yellow, longitudinally splitted into small scales, inner bark with red, sticky sap, fibrous. *Twigs* quadrangular, covered with rusty stellate hairs. *Leaves* simple, opposite, broadly obovate, 16–60 by 12–40 cm, base attenuate to the petiole, apex pointed, young leaves densely covered with yellow stellate hairs, mature leaves rough above, softly hairy below, venation pinnate, veins scalariform. *Petioles* 1–5 cm long, narrowly winged. *Inflorescences* panicles, up to 50 cm long. *Bracts* elliptic. *Flowers* bisexual, small. *Calyx tube* bell-shaped, 5-lobed, densely brown hairy, red dots outside. *Corolla* funnel-shaped at base, with 5 or 6 equal lobes, white, hairy outside. *Stamens* 5 or 6, equal, attached on the base of the corolla. *Ovary* conical, densely outside, style as long as stamens, stigma bifid. *Fruit* a globose

drupe, 1–2 cm diam., densely tomentose, enclosed by calyx, green when young, yellow when mature.

Distribution — Naturally occurring in India, Myanmar, Thailand (northern), and Laos. Introduced into Indonesia (especially Java) and other Asian countries. Teak is also found in the Caribbean region, Florida, Virgin Islands and Hawaii, and in China. In Laos: occurs naturally in Sayabouri province and is introduced in many other provinces. In Vietnam: planted from beginning of the 20th century onward, and is now commonly planted in many provinces.

Habitat & Ecology — Teak grows in mixed tropical deciduous forest and develops well on limestone soils, altitude below 1000 m. Light demanding, even when young, seedlings cannot tolerate frost. Natural and coppice regeneration are good, this species can even tolerate regular forest fires. Flowering: May to July; fruiting: November to January.

Uses — Timber dark yellow, grain gross but fine, resistant to termites and insects. Used for construction, ship making, railway sleepers, and furniture. Young leaves serve as wrapper in preparing food. Boiled leaf extract is used to treat bleeding of larynx, trachea, bronchi, or lungs, and sore throat. In traditional medicine, a wood powder paste has been used against bilious headaches and swellings. The charred wood soaked in poppy juice and made into a paste is used to relieve the swelling of eyelids. The bark has been used as an astringent and the wood oil as a hair tonic. This species is also used as an ornamental and for landscaping.

Specimens examined: *Guillemin* 668 (P); *Iwatsuki et al.* 9446 (L); *Kerr* 1297 (L); *Kostermans* 737, 4719 (L); *Koyama* 32178 (L); *Maxwell* 89-1078 (L); *Murata et al.* 17739 (L); *Pierre* 1275 (P); *Plee* 720 (P); *Thomson s.n.* (L).

INDEX TO SCIENTIFIC NAMES

Numbers refer to the pagenumber. Accepted names are in roman type and synonyms in *italics*.

Acacia

- lebbeck* (L.) Willd. 299
- odoratissima* (L.f.) Willd. 301
- procera* (Roxb.) Willd. 302
- xylocarpa* (Roxb.) Willd. 309
- Acrocarpus Wight & Arn. 244
- combretiflorus* Teijsm. & Binn. 244
- fraxinifolius* Wight & Arn. 244
- Adenanthera L. 297
- microsperma* Teijsm. & Binn. 297
- pavonina* L.
- var. *microsperma* (Teijsm. & Binn.) I.C. Nielsen 297
- Afzelia Sm. 245
- cochinchinensis* (Pierre) Leonard 245
- siamica* Craib 245
- xylocarpa* (Kurz) Craib 245
- Aglaia Lour. 293
- gigantea* Pellegr. 293
- spectabilis* (Miq.) S.S. Jain & Bennet 293
- Albizia Durazz. 299
- lebbeck* (L.) Benth. 299
- odoratissima* (L.f.) Benth. 301
- procera* (Roxb.) Benth. 302
- saman* (Jacq.) F. Muell. 307
- Aleurites J.R. Forst. & G. Forst. 275
- ambinux* Pers. 275
- lobata* Blanco 275
- moluccana* (L.) Willd. 275
- Alstonia R.Br. 233
- scholaris* (L.) R.Br. 233
- Altingia Noroña 224
- gracilipes* Hemsl. 224
- siamensis* Craib 224
- takhtajanii* Thai 224
- Altingiaceae 223
- Amoora gigantea* Pierre 293
- spectabilis* Miq. 293
- Anacardiaceae 226

- Anisoptera Korth. 263
 cochinchinensis Pierre 263
 costata Korth. 263
 oblonga Dyer 263
 robusta Pierre 263
 Annonaceae 229
 Anogeissus 258
 acuminata (Roxb. ex DC.) Guill. ex Perr. 258
 harmandii Pierre 258
 Apocynaceae 232
 Aquilaria Lam. 341
 agallocha Roxb. 341
 crassna Pierre ex Lecomte 341
 Archidendron F. Muell. 303
 clypearia (Jack) I.C. Nielsen
 var. *clypearia* 303
 Baccaurea Lour. 277
 cauliflora Lour. 277
 oxycarpa Gagnep. 277
 ramiflora Lour. 277
 sapida (Roxb.) Müll.Arg. 277
Bassia pasquieri Lecomte 334
 Berrya Roxb. 342
 ammonilla Roxb.
 var. *mollis* Mast. 342
 mollis Wall. ex Kurz 342
 Betula L. 236
 acuminata Wall. 236
 alnoides Buch.-Ham. 236
 Betulaceae 236
Betulaster acuminata Spach 236
Bignonia indica L. 238
 pentandra Lour. 238
 stipulata (Wall.) Roxb. 237
 Bignoniaceae 237
 Bischofia Blume 278
 javanica Blume 278
 trifoliata Hook. 278
 Bombacaceae 240
 Bombax L. 240
 ceiba L. 240
 malabaricum DC. 240
 thorelii Gagnep. 240
 Broussonetia Vent. 310
 papyrifera (L.) Vent. 310
 Burseraceae 241
 Caesalpinia L. 246
 dasyrhachis Miq. 251
 sappan L. 246
 Caesalpinaceae 243
Calophyllum nagassarium Burm.f. 256
Calosanthus indica (L.) Blume 238
 Cananga Hook.f. & Thomson 230
 odorata (Lam.) Hook.f. & Thomson 230
 Canarium L. 242
 album (Lour.) Raeusch. 242
 pimela Leenh. 243
 tonkinensis Engl. 242
 tramdenum C.D. Dai & Yakovlev 243
Caryophyllus L. 315
 Cassia L. 248
 fistula L. 248
 siamea Lam. 252
 Castanea Tourm. ex L. 281
 duclouxii Dode 281
 fargesii Dode 281
 hupehensis Dode 281
 indica Roxb. ex Lindl. 283
 mollissima Blume 281
 Castanopsis Spach 282
 cerebrina (Hickel & A. Camus) Barnett 282
 indica (Roxb. ex Lindl.) A.DC. 283
 sinensis A. Chev. 283
 subacuminata Hayata 283
Chickrassia tabularis (A. Juss.) Wight & Arn.
 296
 Chisocheton Blume 295
 balansae C.DC. 295
 chinensis Merr. 295
 cochinchinensis Pierre 295
 cumingianus (C.DC.) Harms 295
 paniculatus Hiern 295
 siamensis Craib 295
 Choerospondias Burt & Hill 226
 axillaris (Roxb.) Burt & Hill 226
 Chukrasia A. Juss. 296
 tabularis A. Juss. 296
 Cinnamomum Schaeff. 286
 aromaticum Nees in Wall. 286
 balansae Lecomte 287
 camphora (L.) T. Nees & C.H. Eberm. 287
 cassia Blume 286
 glaucescens (Nees) Hand.-Mazz. 287
 iners Reinw. ex Blume 288
 nitidum Hook. 288
 paraneuron Miq. 288
 Clusiaceae 256
 Combretaceae 257
Conocarpus acuminata Roxb. ex DC. 258
 hirtus Buch.-Ham. ex Wall. 258
 Cupressaceae 213
 Cupressus L. 213
 funebri Endl. 213
 hodginsii Dunn 215
 tonkinensis Silba 213
 Dacrycarpus (Endl.) de Laub. 220
 imbricatus (Blume) de Laub. 220

- Dacrydium Sol. ex G. Forst. 221
 elatum (Roxb.) Wall. ex Hook. 221
 pierrei Hickel 221
- Dalbergia L.f. 317
 bariensis Pierre 322
 cambodiana Pierre 318
 cochinchinensis Pierre 318
 cultrata Benth. 320
 dongnaiensis Pierre 322
 duperreana Pierre 322
 fusca Pierre 320
 lanceolaria L.f. 321
 mammosa Pierre 322
 oliveri Prain 322
- Dasillipe pasquieri* Dubard 334
- Datiaceae 260
- Decussocarpus fleuryi* (Hickel) de Laub. 221
- Delonix Raf. 248
 regia (Bojer ex Hook.) Raf. 248
- Dialium L. 249
 cochinchinensis Pierre 249
- Dillenia L. 261
 ovata Wall. ex Hook.f. & Thomson 261
- Dilleniaceae 261
- Diospyros L. 274
 mun A. Chev. & Lecomte 274
- Dipterocarpaceae 262
- Dipterocarpus C.F. Gaertn. 264
 alatus Roxb. & G. Don 265
 gonopterus Turcz. 265
 incanus Roxb. 265
 lemeslei Vesque 265
 macrocarpus Vesque 266
 pubescens Koord. & Valetton 266
 retusus Blume 266
 tonkinensis A. Chev. 266
- Doona odorata* (Roxb.) Burck 268
- Dracontomelon Blume 227
 dao (Blanco) Merr. & Rolfe 227
 duperreanum Pierre 227
 mangiferum Blume 227
 sinensis Stapf 227
- Duabanga Buch.-Ham. 335
 sonneratioides Buch.-Ham. 335
 grandiflora (DC.) Walpert 335
- Dysoxylum esquirolli* H. Lév. 296
- Ebenaceae 274
- Eberhardtia Lecomte 332
 tonkinensis Lecomte 332
- Echites scholaris* L. 233
- Endospermum Benth. 280
 chinense Benth. 280
- Enterolobium saman* (Jacq.) Prain 307
- Erythrina L. 324
 fusca Lour. 324
 ovalifolia Roxb. 324
- Eugenia cumini* L. 315
 jambolan Lam. 315
 jambos L. 316
 auct. 315
- Euphorbiaceae 275
- Fagaceae 280
- Fagraea Thunb. 289
 cochinchinensis A. Chev. 289
 fragrans Roxb. 289
 peregrine Blume 289
- Fokienia A. Henry & H.H. Thomas 215
 hodginsii (Dunn) A. Henry & H.H. Thomas 215
 maclurei Merr. 215
- Gmelina L. 343
 arborea Roxb. 343
 rheedii Hook. 343
- Gordonia wallichii* DC. 339
- Hopea Roxb. 267
 chinensis (Merr.) Hand.-Mazz. 267
 hongayensis Tardieu 267
 odorata Roxb. 268
- Horsfieldia Willd. 312
 amygdalina (Wall.) Warb. 312
 thorelii Lecomte 312
 tonkinensis Lecomte 312
- Inga clypearia* Jack 303
 saman (Jacq.) Willd. 307
 xylocarpa (Roxb.) DC. 309
- Irvingia Hook.f. 284
 harmandiana Pierre ex Laness. 284
 malayana Oliv. ex A. Benn. 284
 oliveri Pierre 284
- Irvingiaceae 283
- Jambosa* Adans. 315
 vulgaris DC. 316
- Jatropha moluccana* L. 275
- Juniperus elata* Roxb. 221
- Keteleeria Carrière 216
 davidiana Beissn. 216
 delavayi Tiegh. 216
 dopiana Flous 216
 evelyniana Mast. 216
 rouletii (A. Chev.) Flous 216
- Knema Lour. 314
 corticosa Lour. 314
 globularia (Lam.) Warb. 314
- Lagerstroemia L. 290
 angustifolia Pierre ex Laness. 290
 calyculata Kurz 290
 grandiflora DC. 335

- Lauraceae 285
Laurus camphora L. 287
Lingoum cambodianum Pierre 328
 indicum (Willd.) Kuntze 326
 macrocarpum (Kurz) Kuntze 328
 wallichii (Wight & Arn.) Pierre 326
 Liquidambar L. 225
 acerifolia Maxim. 225
 formosana Hance 225
 maximoviczii Miq. 225
 tonkinensis A. Chev. 225
 Loganiaceae 288
 Lythraceae 290
 Madhuca J.F. Gmel. 334
 pasquieri (Dubard) H.J. Lam 334
Magnolia mediocris (Dandy) Figlar 292
 Magnoliaceae 291
 Markhamia Seem. 237
 indica P.H. H6 237
 stipulata (Wall.) Seem. ex K. Schum. 237
 Meliaceae 293
 Mesua L. 256
 ferrea L. 256
 nagana Gardner 256
 nagassarium (Burm.f.) Kosterm. 256
 Michelia L. 291
 balansae Dandy 291
 baviensis Finet & Gagnep. 291
 mediocris Dandy 292
 tonkinensis A. Chev. 292
Mimosa dulcis Roxb. 306
 lebbeck L. 299
 odoratissima L.f. 301
 procera Roxb. 302
 saman Jacq. 307
 xylocarpa Roxb. 309
 Mimosaceae 296
 Moraceae 310
Morus papyrifera L. 310
Myristica corticosa (Lour.) Hook.f. 314
 globularia Lam. 314
 Myristicaceae 311
Myrobalanus myriocarpa Kuntze 259
 Myrtaceae 314
Myrtus cumini L. 315
 Nageia Gaertn. 221
 fleuryi (Hickel) de Laub. 221
 nerifolia (D. Don) Kuntze 222
Opa Lour. 315
Oroxylum Vent. 238
 indicum (L.) Kurz 238
Pahudia cochinchinensis Pierre 245
 xylocarpa Kurz 245
 Papilionaceae 317
 Parashorea Kurz 269
 chinensis H.C. Wang 269
 Parkia R.Br. 305
 dongnaiensis Pierre 305
 insignis Kurz 305
 macrocarpa Miq. 305
 streptocarpa Hance 305
 sumatrana Miq. 305
Pasania cerebrina Hickel & A. Camus 282
 Peltophorum (Vogel) Benth. 251
 dasyrhachis (Miq.) Kurz 251
 grande Prain 251
Pentacme siamensis (Miq.) Kurz 270
Pentaptera saja Buch.-Ham. 259
Pierardia sapida Roxb. 277
 Pinaceae 216
Pinus L. 217
 finlaysoniana Wall. ex Blume 218
 insularis Endl. 217
 kesiya Royle ex Gordon 217
 khasya Hook.f. 217
 langbianensis A. Chev. 217
 latteri Mason 218
 merkusiana Cooling & Gausson 218
 merkusii Jungh. & de Vriese 218
 var. *tonkinensis* (A. Chev.) Gausson 218
 sumatrana Jungh. 218
 sylvestris auct. 218
 Pithecellobium Mart. 306
 clypearia (Jack) Benth. 303
 dulce (Roxb.) Benth. 306
 saman (Jacq.) Benth. 307
 Podocarpaceae 219
 Podocarpus L'Hér. ex Pers. 222
 annamensis N.E. Gray 222
 fleuryi Hickel 221
 imbricatus Blume 220
 kawaii Hayata 220
 neriifolius D. Don 222
Poinciana regia Bojer ex Hook. 248
 Polyalthia Blume 231
 cerasoides (Roxb.) Bedd. 231
Poupartia fordii Hemsl. 226
Premna arborea (Roxb.) Roth 343
 Pterocarpus Jacq. 325
 cambodianus (Pierre) Gagnep. 328
 var. *glaucinus* (Pierre) Gagnep. 328
 var. *gracilis* (Pierre) Gagnep. 328
 var. *parviflorus* (Pierre) Gagnep. 328
 indicus Willd. 326
 macrocarpus Kurz 328
 var. *oblongus* (Pierre) Gagnep. 328
 pallidus Blanco 326
 wallichii Wight & Arn. 326
 zollingeri Miq. 326

- Pterospermum* Schreb. 337
 heterophyllum Hance 337
 truncatolobatum Gagnep. 338
Pyrus L. 329
 kumaoni Decne. 329
 pashia D. Don 329
 variolosa Wall. 329
 verruculosa Bertol. 329
Rosaceae 329
Samanea Merr. 307
 saman (Jacq.) Merr. 307
Sapindaceae 331
Sapindus L. 331
 microcarpus Jard. 331
 mukorossi Gaertn. 331
 saponaria L. 331
 vitiensis A. Gray 331
Sapotaceae 332
Schima Reinw. ex Blume 339
 crenata Korth. 339
 noronhae Reinw. ex Blume 339
 wallichii (DC.) Korth. 339
Senna Mill. 252
 siamea (Lam.) Irwin & Barneby 252
Shorea Roxb. ex C.F. Gaertn. 270
 bracteata Pierre ex Laness. 270
 chinensis (H.C. Wang) H. Zhu 269
 chinensis Merr. 267
 mekongensis Pierre ex Laness. 270
 siamensis Miq. 270
 wangtianshuea Y.K. Yang & J.K. Wu 269
Sindora Miq. 253
 cochinchinensis Baill. 253
 siamensis Teijsm. 253
 wallichii
 var. *siamensis* (Teijsm. ex Miq.) Baker
 253
Sonneratiaceae 335
Spathodea indica (L.) Pers. 238
 stipulata Wall. 237
 velutina Kurz 237
Spondias acuminata Gamble 226
 axillaris Roxb. 226
Sterculiaceae 337
Sunaptea odorata Griff. 272
Swietenia chickrassa Roxb. 296
Syzygium Gaertn. 315
 cumini (L.) Skeels 315
 jambos (L.) Alston 316
Tamarindus L. 255
 indica L. 255
 occidentalis Gaertn. 255
 officinalis Hook. 255
Tectona L.f. 344
 grandis L.f. 344
 theka Lour. 344
Terminalia 259
 myriocarpa Van Heurck & Müll.Arg. 259
 saja Steud. 259
Tetrameles R.Br. 260
 grahamiana Wight 260
 nudiflora R.Br. 260
Theaceae 338
Thymelaeaceae 340
Tiliaceae 342
Tsuga rouletii A. Chev. 216
Uvaria odorata Lam. 230
Vatica L. 272
 fleuryana Tardieu 272
 odorata (Griff.) Symington 272
 tonkinensis A. Chev. 272
Verbenaceae 343
Wrightia R.Br. 235
 annamensis Eberh. & Dubard 235
 pubescens R.Br. 235
 tomentosa Roem. & Schult.
 var. *cochinchinensis* Pierre ex Pit. 235
Xylia Benth. 309
 dolabriformis Benth. 309
 kerrii Craib & Hutch. 309
 xylocarpa (Roxb.) Taub.
 var. *kerrii* (Craib & Hutch.) I.C. Nielsen
 309
Xylopia L. 232
 vielana Pierre 232