

## ENTOLOMA SUBGENUS NOLANEA—ADDITIONS

M. E. NOORDELOOS

*Rijksherbarium, Leiden*

1. The sectional name *Cosmeoexonema* Largent & Thiers (1972) has priority over the sectional name *Papillata* Romagnesi 1974. This has been corrected in Noordeloos, 1981: 141. As a consequence the following new combination is necessary: **Entoloma subsection Papillata** (Romagn.) Noordeloos *comb. nov.*—Basionym: *Rhodophyllus* section *Papillati* Romagn. in Bull. mens. Soc. linn. Lyon 43: 330. 1974.

2. Subsection *Endochromonema* has been published by me (Noordeloos, 1980: 495) without a Latin diagnosis; no type has been indicated. This is corrected below:

### **Entoloma subsection Endochromonema** Noordeloos, *subsect. nov.*

Pigmentae pallidae, intracellulosae vel membranares, haud incrustantes; spora heterodiametrae vel isodiametrae, haud multiangulatae nodulosae; pileus griseo-brunneus vel ochraceo-fulvus, nunquam atro-brunneus; stipes fibrillosus, saepe striatus, haud citrinus vel flavovirens; odore nullo vel farinaceo, nunquam aromatico.—Typus: *Entoloma cetratum* (Fr.) Moser.

3. Some spelling-corrections: Read *Entoloma cuspidiferum* in stead of *E. cuspidifer* and *E. globuliferum* in stead of *E. globulifer*.

4. In my key to the sections and subsections (Noordeloos, 1980: 431–432) the couplets 6 and 11 have to be changed in the following way:

- 6a. Cheilocystidia tibiiform, sphaeropedunculate or cylindrical to clavate, sometimes subcapitate, spores with dihedral base . . . . . Subsect. *Cosmeoexonema*, p. 472  
11b. Carpophores moderately to strongly pigmented; lamellae always with distinct grey or brown tinge . . . . .  
11'a. Spores with dihedral base . . . . . Subsect. *Cosmeoexonema*, p. 472  
11'b. Spores with basal facet . . . . . Subsect. *Papillata*, p. 553

### REFERENCES

- NOORDELOOS, M. E. (1980). *Entoloma* subgenus *Nolanea* in the Netherlands and adjacent regions with a reconnaissance of its remaining taxa in Europe. *In Persoonia* 10: 427–534.